

Hard exclusive meson production at HERMES, COMPASS and JLAB

Probing Strangeness in Hard Processes

Frascati, Italy


18-21 October, 2010

Ami Rostomyan




exclusive meson production

VMD




GPD

$Q^2 \gg 1 \text{ GeV}^2$




generalized parton distributions



Nucleon Tomography


correlation between longitudinal momentum and transverse position

Elastic Form Factors




transverse position of partons

Parton Distribution Functions (PDFs)



longitudinal momentum of partons

at leading twist: $H E \tilde{H} \tilde{E}$



$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H_q(x, \xi, t) + E_q(x, \xi, t)]$$

$$J_g = \frac{1}{2} \lim_{t \rightarrow 0} \int_0^1 dx [H_g(x, \xi, t) + E_g(x, \xi, t)]$$


factorization for σ_L (and ρ_L, ω_L, ϕ_L) only

 $\sigma_L - \sigma_T$ suppressed by $1/Q$

 σ_T suppressed by $1/Q^2$


generalized parton distributions

$$R = \frac{\sigma_L}{\sigma_T}$$



- ◆ suppression effectively is not working for $Q^2 \sim \text{few GeV}$
- ◆ valuable information on GPDs from higher twist terms

at leading twist: $H \ E \ \tilde{H} \ \tilde{E}$



$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H_q(x, \xi, t) + E_q(x, \xi, t)]$$

$$J_g = \frac{1}{2} \lim_{t \rightarrow 0} \int_0^1 dx [H_g(x, \xi, t) + E_g(x, \xi, t)]$$

factorization for σ_L (and ρ_L, ω_L, ϕ_L) only

 $\sigma_L - \sigma_T$ suppressed by $1/Q$

 σ_T suppressed by $1/Q^2$

given channel probes specific GPD flavor

$$\rho^0 p \quad \frac{1}{\sqrt{2}} [2u + d] + \frac{1}{\sqrt{2}} [2\bar{u} + \bar{d}] + \frac{9}{4} g$$

$$\omega p \quad \frac{1}{\sqrt{2}} [2u - d] + \frac{1}{\sqrt{2}} [2\bar{u} - \bar{d}] + \frac{3}{4} g$$

$$\phi p \quad -[s + \bar{s}] + \frac{3}{4} g$$

$$\rho^+ n \quad 2[u - d] - [\bar{u} - \bar{d}]$$

$$K^{*+} \Lambda \quad -\frac{2}{\sqrt{6}} [2u - d - s]$$

$$+\frac{1}{\sqrt{6}} [2\bar{u} - \bar{d} - \bar{s}]$$

$$K^{*+} \Sigma^0 \quad -\frac{2}{\sqrt{2}} [d - s] + \frac{1}{\sqrt{2}} [\bar{d} - \bar{s}]$$

$$K^{*0} \Sigma^+ \quad [d - s] + [\bar{d} - \bar{s}]$$

pseudoscalar mesons


at leading twist: \tilde{H} \tilde{E}

higher twist: H_T

vector mesons

at leading twist: H E

higher twist: \tilde{H} \tilde{E}



SU(3):

◆ relate nucleon to octet hyperon

◆ relate $p \rightarrow N$ transition GPDs to $p \rightarrow p'$

$$\pi^+ n \quad 2[\Delta u - \Delta d] + [\Delta \bar{u} - \Delta \bar{d}]$$

$$\pi^0 p \quad \frac{1}{\sqrt{2}} [2\Delta u + \Delta d] - \frac{1}{\sqrt{2}} [2\Delta \bar{u} + \Delta \bar{d}]$$

$$K^+ \Lambda \quad -\frac{2}{\sqrt{6}} [2\Delta u - \Delta d - \Delta s]$$

$$-\frac{1}{\sqrt{6}} [2\Delta \bar{u} - \Delta \bar{d} - \Delta \bar{s}]$$

$$K^+ \Sigma^0 \quad -\frac{2}{\sqrt{2}} [\Delta d - \Delta s] - \frac{1}{\sqrt{2}} [\Delta \bar{d} - \Delta \bar{s}]$$

$$K^0 \Sigma^+ \quad [\Delta d - \Delta s] - [\Delta \bar{d} - \Delta \bar{s}]$$

modeling GPDs

- ◆ constraints on the t-behavior of valence quark and gluon GPDs

$$H^q(x, \xi, t) = H^q(x, \xi) F_1^p(t)$$

$$H^g(x, \xi, t) = H^g(x, \xi) F_1^p(t)$$

- ◆ t-behavior of sea quarks is unknown
- ◆ same t-dependence for quarks and gluons

quarks

quarks and gluons

- ➔ measure the t-dependence of cross section (e.g. ρ^+ and ρ^0)

$$\int_0^1 dx E_g + \sum_q \int_{-1}^1 dx x E_q = 0$$


- Diehl (2003) -

- ◆ $E_u \approx -E_d$
- ◆ expectation:

$$\int_0^1 dx E_g = -2 \sum_q \int_0^1 dx x E_{\bar{q}}$$

- ◆ $E_{\bar{q}}$: small sea quark contribution at $x \sim 0.1$
- ◆ small E_g

- Ellinghaus, Nowak, Vinnikov, Ye (2005) -



- VGG code -

vector mesons

t-dependence



$$\frac{d\sigma}{dt} \sim e^{-bt}$$

$$b \sim R_p^2 + R_{q\bar{q}}^2$$

t-dependence

$$\frac{d\sigma}{dt} \sim e^{-bt}$$



$$b \sim R_p^2 + R_{q\bar{q}}^2$$



t-dependence

$$\frac{d\sigma}{dt} \sim e^{-bt}$$




$$b \sim R_p^2 + R_{q\bar{q}}^2$$




t-dependence

$$\frac{d\sigma}{dt} \sim e^{-bt}$$


$$b \sim R_p^2 + R_{q\bar{q}}^2$$



$\gamma^* p \rightarrow \phi p$ cross section










- ◆ similar trend as a function of Q^2
- ◆ monotonically increase with W
- ◆ Regge model calculation
 $W = 2.1, 2.45, 2.9$ GeV
 - *Laget (2003)* -
➔ overestimate the CLAS data



- ◆ applicability of GPD formalism at low W data? probe of gluon field in the nucleon
- ◆ longitudinal cross section is not dominant in low- W kinematics
- ◆ modified perturbative approach might be successful - *Goloskokov Kroll (2007)* -

$\gamma^* p \rightarrow \rho^+ n$ cross section

CLAS preliminary data
 - Fradi (2010) hep-ex/10101198 -



◆ decrease of total cross section with W

Regge model

- Laget (2003) -

◆ exchange of ρ^+ and π^+


◆ good description in all ranges




$\gamma^* p \rightarrow \rho^+ n$ cross section

CLAS preliminary data
- *Fradi (2010) hep-ex/10101198* -


1.00 <math>Q^2(\text{GeV}^2) < 1.50</math>




1.50 <math>Q^2(\text{GeV}^2) < 2.00</math>




2.00 <math>Q^2(\text{GeV}^2) < 2.50</math>




2.50 <math>Q^2(\text{GeV}^2) < 3.00</math>




3.00 <math>Q^2(\text{GeV}^2) < 3.50</math>



3.50 <math>Q^2(\text{GeV}^2) < 4.00</math>



4.00 <math>Q^2(\text{GeV}^2) < 4.50</math>



◆ decrease of longitudinal cross section with W

◆ *GK GPD model*

- *Goloskokov Kroll (2005)* -

◆ *VGG GPD model*

- *Vanderhaeghen, Guichon, Guidal (1999)*

◆ models do not describe the data

▶ GPD formalism is not applicable

▶ missing contribution is GPD parameterizations





cross section ratios

- ◆ cross sections change significantly when varying the nonperturbative input: MRST % CTEQ

- ▶ unpolarized gluon densities at low scales

- Diehl, Kugler, Schaefer, Weiss (2005) -



- ◆ next-to leading order corrections

- ◆ substantial power corrections

➔ cross section ratios for similar channels

➔ cancelation of theoretical uncertainties

$$\frac{\sigma_L(\gamma^* p \rightarrow \phi p)}{\sigma_L(\gamma^* p \rightarrow \rho^0 p)} = \text{const.}$$


$$\frac{\sigma_L(\gamma^* p \rightarrow K^{*+} \Lambda)}{\sigma_L(\gamma^* p \rightarrow \rho^+ n)} \approx \frac{3}{2}$$

ρ^0 transverse target spin asymmetry

- ◆ cross section asymmetry with respect to transverse target polarization

$$A_{\text{target pol.}}^{\gamma^* p \rightarrow \rho_L^0 p} = \frac{|\Delta_{\perp}| \text{Im}(\mathcal{E}^* \mathcal{H})}{(1 - \xi^2)|\mathcal{H}|^2 - (\xi^2 + t/4M^2)|\mathcal{E}|^2 - 2\xi^2 \text{Re}(\mathcal{E}^* \mathcal{H})}$$

- ◆ depends linearly on the helicity-flip GPD E
- ◆ no kinematic suppression of GPD E with respect to GPD H



- HERMES Collaboration (2009) -

average kinematics:

$$\langle -t' \rangle = 0.13 \text{ GeV}^2$$

$$\langle x_B \rangle = 0.09$$

$$\langle Q^2 \rangle = 2.0 \text{ GeV}^2$$

- ◆ L/T separation using the angular distribution
- ◆ leading twist contribution: compatible with 0 overall value

$$A_{\text{target pol.}}^{\gamma^* p \rightarrow \rho_L^0 p} = -0.033 \pm 0.058$$

- ◆ implies that the E^g is small

$$A_{\text{target pol.}}^{\gamma^* p \rightarrow \rho_L^0 p} \propto \frac{E}{H} \propto \frac{E^q + E^g}{H^q + H^g}$$

ρ^0 transverse target spin asymmetry


- ◆ cross section asymmetry with respect to transverse target polarization

$$A_{\text{target pol.}}^{\gamma^* p \rightarrow \rho_L^0 p} = \frac{|\Delta_{\perp}| \text{Im}(\mathcal{E}^* \mathcal{H})}{(1 - \xi^2)|\mathcal{H}|^2 - (\xi^2 + t/4M^2)|\mathcal{E}|^2 - 2\xi^2 \text{Re}(\mathcal{E}^* \mathcal{H})}$$


- ◆ depends linearly on the helicity-flip GPD E
- ◆ no kinematic suppression of GPD E with respect to GPD H

- COMPASS Collaboration -


$\langle Q^2 \rangle \approx 2.2 \text{ (GeV/c)}^2$



$\langle x_{Bj} \rangle \approx 0.04$



$\langle p_t^2 \rangle \approx 0.18 \text{ (GeV/c)}^2$



- ◆ no L/T separation yet
- ◆ compatible with 0 overall value

ρ^0 transverse target spin asymmetry

- ◆ cross section asymmetry with respect to transverse target polarization

$$A_{\text{target pol.}}^{\gamma^* p \rightarrow \rho_L^0 p} = \frac{|\Delta_{\perp}| \text{Im}(\mathcal{E}^* \mathcal{H})}{(1 - \xi^2)|\mathcal{H}|^2 - (\xi^2 + t/4M^2)|\mathcal{E}|^2 - 2\xi^2 \text{Re}(\mathcal{E}^* \mathcal{H})}$$


- ◆ depends linearly on the helicity-flip GPD E
- ◆ no kinematic suppression of GPD E with respect to GPD H

recoil polarization asymmetry

- ◆ cross section asymmetry with respect to recoil polarization

$$A_{\text{recoil pol.}}^{\gamma^* p \rightarrow K^{*+} \Lambda} = \frac{|\Delta_{\perp}| \text{Im}(\mathcal{E}^* \mathcal{H})}{(1 - \xi^2)|\mathcal{H}|^2 - (\xi^2 + t/4M^2)|\mathcal{E}|^2 - 2\xi^2 \text{Re}(\mathcal{E}^* \mathcal{H})}$$




- ◆ access to GPD E from unpolarized target



- Goloskokov, Kroll (2008) -

double spin asymmetry

- ◆ asymmetry arise due to interference between natural and unnatural parity exchange amplitudes
- ◆ non-leading twist: interference between the GPD $H^{s,g}$ and $\tilde{H}^{s,g}$



- Goloskokov, Kroll (2007) -

- $W = 5$ GeV
- .-.- $W = 10$ GeV
- HERMES
- ◇ COMPASS

- ◆ asymmetry is compatible with 0
- ➔ UPE contribution is negligible
- ◆ measurements at different W ranges
- ➔ no W -dependence
- ◆ prediction: relate the A_1^ρ to inclusive asymmetry

$$A_1^\rho = \frac{2A_1}{1 + (A_1)^2}$$

- Fraas -



- ➔ prediction is consistent with data

pseudoscalar mesons

π^+ cross section from Hall C

$$2\pi \frac{d^2\sigma}{dt d\phi} = \frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \sqrt{2\epsilon(1+\epsilon)} \frac{d\sigma_{LT}}{dt} \cos\phi + \epsilon \frac{d\sigma_{TT}}{dt} \cos 2\phi.$$


- The Jefferson Lab $F_{\pi-2}$ Collaboration (2006) -



- ◆ two beam energies
 - ➔ Rosenbluth separation
- ◆ simultaneous fit to the angular distribution

- ◆ L/T separated cross sections
- ◆ longitudinal cross section is well reproduced by the model prediction: pion and rho Regge trajectory exchange
 - Vanderhaeghen, Guidal, Laget (1997) -
- ◆ transverse component undershoot

π^+ cross section from HERMES



◆ no L/T separation - HERMES Collaboration (2007) -

◆ longitudinal component is expected to dominate (large t)

GPD model calculations

◆ GPD E is considered to be dominated by the t -channel pion pole

◆ H is neglected

◆ leading order calculations underestimate the data

◆ power corrections agree with data

Regge model calculations

◆ transverse component of the cross section 6-8% at $-t' < 0.07 \text{ GeV}^2$

◆ the same model underestimate the JLAB data
(holds also for HERMES higher W kinematics?)

π^+ transverse target asymmetry

$$A_{UT,\ell}^{\sin(\phi-\phi_S)} = -\frac{\sqrt{-t'}}{M_p} \times \frac{\xi\sqrt{1-\xi^2}\text{Im}(\tilde{\mathcal{E}}^*\tilde{\mathcal{H}})}{(1-\xi^2)\tilde{\mathcal{H}}^2 - \frac{t\xi^2}{4M_p^2}\tilde{\mathcal{E}}^2 - 2\xi^2\text{Re}(\tilde{\mathcal{E}}^*\tilde{\mathcal{H}})}$$

- HERMES Collaboration (2007) -


average kinematics:

$$\langle -t' \rangle = 0.18 \text{ GeV}^2$$

$$\langle x_B \rangle = 0.13$$

$$\langle Q^2 \rangle = 2.38 \text{ GeV}^2$$


- ◆ 6 azimuthal moments extracted according to - Diehl, Sapeta (2005) -
- ◆ no L/T separation
- ◆ small overall value for leading asymmetry
- ◆ unexpected large overall value for $A_{UT}^{\sin\phi_S}$
- ➡ evidence of contribution from transversely polarized photons



more on K and π mesons

$$A_{\text{UT},\ell}^{\sin(\phi-\phi_S)} = -\frac{\sqrt{-t'}}{M_p} \times \frac{\xi\sqrt{1-\xi^2}\text{Im}(\tilde{\mathcal{E}}^*\tilde{\mathcal{H}})}{(1-\xi^2)\tilde{\mathcal{H}}^2 - \frac{t\xi^2}{4M_p^2}\tilde{\mathcal{E}}^2 - 2\xi^2\text{Re}(\tilde{\mathcal{E}}^*\tilde{\mathcal{H}})}$$

◆ the same GPDs accessible from $\gamma_L^* p \rightarrow K^+ \Lambda, K^0 \Sigma^+$




- Goeke, Polyakov, Vanderhaeghen (1999) -

◆ results from CLAS on Beam-recoil polarization transfer in the nucleon resonance region in exclusive reactions

$$\gamma^* p \rightarrow K^+ \Lambda, K^0 \Sigma^+$$

◆ beyond the resonance region measurements of cross sections and asymmetries might be helpful

◆ separate the pion/kaon pole term contributions




◆ cross section ratios:

$$\frac{\gamma^* p \rightarrow \pi n}{\gamma^* p \rightarrow \pi p} \quad \frac{\gamma^* p \rightarrow K^+ \Sigma^0}{\gamma^* p \rightarrow K^0 \Sigma^+}$$

- Strikman, Weiss (2008) -

- Diehl, Kugler, Schaefer, Weiss (2005) -

summary




BACKUP SLIDES

ρ^0 cross section

CLAS Hall B
GPD model

Regge model



$$1.5 < Q^2 < 3 \text{ GeV}^2, \quad W > 1.75 \text{ GeV}, \quad 0.21 < x_B < 0.62$$

