



Updated study of Had Recoil $B \rightarrow K^* \nu \nu$ vs Bwd EMC

Elisa Manoni

INFN Sez. Perugia



Det + Comp + Physics: FastSim + detector + DGWG - SuperB
workshop, LNF, April 05, 2011



Outline

- Neutral energy smearing algorithm
- Patch validation
- Results on Bwd EMC veto impact



Motivations and analysis strategy

- September production ntuple: neutral energy smearing not applied
- @ Caltech: analysis of September sample to evaluate the impact of Bwd EMC used as veto device
 - without smearing, results can be too optimistic
- Need to re-compute impact of Bwd EMC veto including resolution effects:
 - make a new production for the BBbar sample is too time consuming
 - **apply smearing off-line**
 - validate smearing algorithm on signal and single particle MCs
 - apply off line smearing on both BBbar and signal samples from September production
 - repeat the analysis and compare $S/\sqrt{B+B}$ w and w/o Eextra_bwd veto



FastSim energy smearing algorithm (I)

- BaBar EMC energy resolution:

$$\sigma(E)/E = 2.35\%/E^{(1/4)} + 1.35\%$$

- Energy resolution parameterization in FastSim:

gaussian component: $\sigma(E)/E = f_a/E^{(e_p)} + o f_b$

+ exponential tail: $\tau = c_{exp}/E^{(p_{exp})} + o d_{exp}$

- parameter values :

par	fwd	brr	bwd
fa	0.0102	0.0102	0.14
fb	0.0	0.0	0.03
ep	0.264	0.264	0.5
cexp	0.0165	0.0165	0.0
dexp	0.0284	0.0284	0.0
pexp	0.05	0.050	0.0



FastSim energy smearing algorithm (II)

- Experimental effects accounted in FastSim:
 - global EMC calibration: apply scaling factor to recover part of the cluster not contained
 - background cluster effects: once the “physics” cluster has been reconstructed, switch on random crystals around the cluster
 - energy smearing due to finite resolution



smearing parameter:

$$\delta E = \text{gaussRnd}(0, \sigma(E)) - \text{expRnd}(\tau(E))$$


gaussRnd/expRnd = random numbers generated according to gaussian/exponential distribution (function of resolution params)

→ smeared energy: $E_{\text{meas}} = E_0 * (1 + \delta E)$



FastSim energy smearing algorithm (III)

- September production:
 - global EMC calibration ON
 - background cluster effects ON
 - energy smearing due to finite resolution OFF





Patch validation



Strategy

- Compare “production with smearing turned on” and “production with smearing turned off + off-line smearing”
- samples: $B^0 \rightarrow K^{*0} \nu \nu$ signal, single- π^0 beam, single- γ beam MCs
- Legend in the following plots:

— old prod, off-line smearing

1) Production without energy smearing + patch with smearing applied after reconstruction (algorithm we want to validate)

— old prod, no off-line smearing

2) Production without energy smearing

+ new prod, on-line smearing

3) Production with energy smearing

- 1) is obtained starting from 2) and applying the smearing algorithm, it aims to reproduce 3)

→ black histo and blue + should match




$B^0 \rightarrow K^{*0} \nu \nu$ signal MC: sample and selection

- sample:
 - ~3M events from September production (V0.2.5 FastSim release, smearing OFF)
 - ~10M events with V0.2.6 FastSim release (smearing ON)
- generate $B^0 \rightarrow K^{*0} \nu \nu$ vs $B^0 \text{bar} \rightarrow$ hadronic modes
reconstruct with PacHadRecoilUser package (FastSim package for hadronic Breco analysis)
- selection
 - examine all reconstructed γ (irrespective of their origin, i.e. γ from any decay product of both Breco and Bsig)
 - thruth matching: reco photons associated to one photon in the MC list



$B^0 \rightarrow K^{*0} \nu \bar{\nu}$ signal MC : γ energy (I)


– $E_{\text{true}} - E_{\text{reco}}$ in bins of E_{true} , truth-matching required





$B^0 \rightarrow K^{*0} \nu \nu$ signal MC : π^0 mass

- list-level cuts and fit constraints not reproduced applying smearing after reconstruction, see slide 14 for single π^0 beam results





Single π^0 : sample and selection

- sample:
 - 500K events with V0.2.6 FastSim release and smearing OFF
 - 500K events with NOMINAL V0.2.6 FastSim release (smearing ON)


- π^0 list
 - γ s from CalorNeutral list,
 - $\gamma\gamma$ invariant mass cut: [0.090,0.165] GeV
 - Pmin set 0.05 GeV, Pmax set 4.0 GeV
 - CosThetamin set -1, CosThetamax set 1

- selection
 - 1 reco π^0
 - thruth matching: π^0 reco daughters associated to 2nd and 3rd particle in the MC list (to cut reco γ matched with bremsstrahlung γ produced after $\gamma \rightarrow e^+e^-$ conversion)



Single $\pi^0 : \gamma$ energy

- $E_{\text{true}} - E_{\text{reco}}$ in bins of E_{true} , truth-matching required






Single π^0 : π^0 mass

m_{π^0} , Brr


both γ s in the barrel

hemc0	
Entries	307792
Mean	0.1357
RMS	0.007933
Underflow	0.01243
Overflow	0.00244
Integral	1



both γ s in the forward

hemc1	
Entries	3873
Mean	0.1372
RMS	0.006778
Underflow	0.01203
Overflow	0.0005229
Integral	1






Single γ : sample and selection

- sample:
 - 500K events with V0.2.6 FastSim release and smearing OFF
 - 500K events with NOMINAL V0.2.6 FastSim release (smearing ON)
- γ list
 - CalorNeutral list
 - Pmin set 0.05 GeV , Pmax set 4.0 GeV
 - FULL ANGULAR COVERAGE : CosThetamin set -1, CosThetamax set 1
 - BWD REGIONS: CosThetamin set -1, CosThetamax set -0.9
CosThetamin set -0.9615, CosThetamax set -0.8815
- selection
 - 1 reco photon
 - thruth matching: reco photon associated to 1st particle in the MC list (to cut reco γ matched with breemmstrahlung γ produced after $\gamma \rightarrow e^+e^-$ conversion)



Single γ – full angular coverage : γ energy (I)




– $E_{\text{true}} - E_{\text{reco}}$ in bins of E_{true} , truth-matching required





Single γ – full angular coverage : γ energy (II)


– E_{reco} in different EMC regions





Single γ – BWD angular coverage : γ energy (I)

- $E_{\text{true}} - E_{\text{reco}}$ in bins of E_{true} , truth-matching required







Single γ – BWD angular coverage : γ energy (II)

– Angular coverages:

- CosTheta [-1,-0.9] (BLACK)
- CosTheta [-0.9615,-0.8815] (RED)






- $E_{\text{true}} - E_{\text{reco}}$ vs E_{true}
truth-matching required





Single γ – BWD angular coverage : γ energy resolution

Etrue - Ereco





Physics results





Eextra distributions, before Bsig selection

– one example: $B^0 \rightarrow K^{*0}(K\pi)\nu\nu$
barrel+fwd



bwd

- without smearing
+ with smearing

signal



BB
cocktail







Eextra BWD distributions, after Bsig selection

– $B^+ \rightarrow K^{*+}(K_s \pi) \nu \nu$



bwd

Extra bwd w smearing - Eextra bwd w/o smearing

signal



BB cocktail







Eextra BWD distributions, after Bsig selection

– $B^+ \rightarrow K^{*+}(K\pi^0)\nu\nu$



bwd

Extra bwd w smearing - Eextra bwd w/o smearing

signal



BB cocktail





Extra BWD distributions, after Bsig selection

– $B^0 \rightarrow K^{*0}(K\pi)\nu\nu$



bwd

Extra bwd w smearing - Extra bwd w/o smearing

signal



BB cocktail





Results: satrtegy

- $E_{\text{extra_bwd}} < 50 \text{ MeV}$, $E_{\text{min},\gamma} = 30 \text{ MeV}$
- Changes wrt to Preliminary results shown in <http://agenda.infn.it/getFile.py/access?contribId=2&resId=0&materialId=slides&confId=3464>
 - fixed bug in Bwd-Barrel angular coverage
 - fixed bug in extra-photon finding algorithm (some of the Breco modes where not incorporated in the study)
 - remove cut on $E_{\text{extra_barrel+fwd}}$ (uncorrelated to $E_{\text{extra_bwd}}$) to increase the statistics

Figure of Merit

- Significance = $S/\sqrt{S+B}$
- $\Delta\text{Significance}/\text{Significance} = (\text{Sig_bwd}-\text{Sig_nobwd})/\text{Sig_nobwd}$
- in the limit $S \ll B$:

$$\Delta\text{Significance}/\text{Significance} = (\epsilon_{\text{sig}}/\sqrt{\epsilon_{\text{bb}}}) - 1$$

being ϵ_{sig} (ϵ_{bb}) the marginal efficiency of the $E_{\text{extra_bwd}}$ cut in signal (BBbar) MC sample



Results : relative change in significance

SMEARING OFF			
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$	786	778	$(99.98 \pm 0.36)\%$
B^0 had cocktail	181	143	$(79.0 \pm 3.0)\%$
$\Delta Sign/Sign$	$(11.4 \pm 1.9)\%$		
$B^+ \rightarrow K^{*+}(K_S \pi^+) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	233	232	$(99.57 \pm 0.43)\%$
B^+ had cocktail	136	114	$(83.8 \pm 3.2)\%$
$\Delta Sign/Sign$	$(8.7 \pm 1.9)\%$		
$B^+ \rightarrow K^{*+}(K^+ \pi^0) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	227	222	$(97.8 \pm 1.0)\%$
B^+ had cocktail	75	65	$(86.7 \pm 3.9)\%$
$\Delta Sign/Sign$	$(5.0 \pm 2.4)\%$		

SMEARING ON			
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$	786	778	$(99.98 \pm 0.36)\%$
B^0 had cocktail	181	146	$(80.7 \pm 2.9)\%$
$\Delta Sign/Sign$	$(10.2 \pm 1.8)\%$		
$B^+ \rightarrow K^{*+}(K_S \pi^+) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	233	232	$(99.57 \pm 0.43)\%$
B^+ had cocktail	136	114	$(83.8 \pm 3.2)\%$
$\Delta Sign/Sign$	$(8.7 \pm 1.9)\%$		
$B^+ \rightarrow K^{*+}(K^+ \pi^0) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	227	221	$(97.4 \pm 1.1)\%$
B^+ had cocktail	75	65	$(86.7 \pm 3.9)\%$
$\Delta Sign/Sign$	$(4.6 \pm 2.4)\%$		



Conclusion

- Algorithm to apply off-line neutral energy smearing in place
- algorithm validated on gamma and π^0 single particle beams and signal MC
- still some issue on Bwd EMC simulation in FastSim under study
- Negligible effect of smearing on physics result:
5-10% gain in significance with Eextra,bwd veto




Back-up slides



Caltech results

– E_{extra_bwd}
 < 0.05 GeV:

$B^0 \rightarrow K^{*0} \nu \bar{\nu}$					
Sample	N_{sel}	ϵ_{tot}	$N_{sel,Bwd}$	$\epsilon_{tot,Bwd}$	$\delta\epsilon/\epsilon$
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$	727	$(24.5 \pm 0.9) \times 10^{-5}$	719	$(24.2 \pm 0.9) \times 10^{-5}$	$(-1.1 \pm 0.4)\%$
B^0 had cocktail	76	$(20 \pm 2) \times 10^{-8}$	60	$(16 \pm 2) \times 10^{-8}$	$(-21 \pm 7)\%$
S/\sqrt{B}	83 ± 7		93 ± 9		
$B^+ \rightarrow K^{*+} (K_s^+ \pi^+) \nu \bar{\nu}$					
Sample	N_{sel}	ϵ_{tot}	$N_{sel,Bwd}$	$\epsilon_{tot,Bwd}$	$\delta\epsilon/\epsilon$
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	223	$(7.1 \pm 0.5) \times 10^{-5}$	222	$(7.0 \pm 0.5) \times 10^{-5}$	$(-0.5 \pm 0.4)\%$
B^+ had cocktail	48	$(12.0 \pm 1.7) \times 10^{-8}$	40	$(10.0 \pm 1.7) \times 10^{-8}$	$(-17 \pm 7)\%$
S/\sqrt{B}	32 ± 4		35 ± 5		






$$\delta \left(\frac{S}{\sqrt{(B)}} \right) = \frac{\left(\frac{S}{\sqrt{(B)}} \right)_{bwd} - \left(\frac{S}{\sqrt{(B)}} \right)_{nobwd}}{\left(\frac{S}{\sqrt{(B)}} \right)_{nobwd}} =$$

$K\pi : (10 \pm 3)\%$
 $K_s \pi : (8 \pm 3)\%$



Single γ – BWD angular coverage : γ energy (II)

- Angular coverages:
 - CosTheta [-1,-0.9] (BLACK)
 - CosTheta [-0.9615,-0.8815] (RED)






$B^0 \rightarrow K^{*0} \nu \nu$ signal MC : γ angle

- $\theta_{\text{true}} - \theta_{\text{reco}}$ in bins of E_{true} , truth-matching required
- NO ANGULAR SMEARING APPLIED: red and black matches (same sample)


- large tails due to reco gamma not correctly matched (probably associated to bremsstrahlung - γ produced after $\gamma \rightarrow e^+e^-$ conversion), improvements in single particle tests with more stringent matching requirements (slide 33)





Single π^0 : γ angle




— $\theta_{\text{true}} - \theta_{\text{reco}}$ and $\phi_{\text{true}} - \phi_{\text{reco}}$ in bins of E_{true} , truth-matching required





$B^0 \rightarrow K^{*0} \nu \nu$ signal MC : γ energy (II)

– E_{reco} in different EMC regions





Eextra distributions, before Bsig selection



– $B^+ \rightarrow K^{*+}(K_s \pi) \nu \nu$

barrel+fwd



bwd

- without smearing
+ with smearing

signal



BB cocktail





Eextra distributions, before Bsig selection



– $B^+ \rightarrow K^{*+}(\bar{K}\pi^0)\nu\nu$

barrel+fwd



bwd

- without smearing
+ with smearing

signal




BB
cocktail







Results : significance-flow in $K\pi$



Marginal efficiency, signal



Marginal efficiency, $B\bar{B}$





Results with E_{extra_brrfwd} cut

SMEARING ON, E_{extra}^{brrfwd} cut			
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^0 \rightarrow K^{*0} \nu \bar{\nu}$	735	727	$(98.91 \pm 0.38)\%$
B^0 had cocktail	91	75	$(82.4 \pm 4.0)\%$
$\Delta Sign/Sign$	$(8.9 \pm 2.4)\%$		
$B^+ \rightarrow K^{*+}(K_S \pi^+) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	191	190	$(99.48 \pm 0.52)\%$
B^+ had cocktail	76	66	$(86.8 \pm 3.9)\%$
$\Delta Sign/Sign$	$(6.7 \pm 2.3)\%$		
$B^+ \rightarrow K^{*+}(K^+ \pi^0) \nu \bar{\nu}$			
Sample	N_{sel}	$N_{sel,Bwd}$	ϵ
$B^+ \rightarrow K^{*+} \nu \bar{\nu}$	214	208	$(97.2 \pm 1.1)\%$
B^+ had cocktail	40	33	$(82.5 \pm 6.0)\%$
$\Delta Sign/Sign$	$(7.0 \pm 3.7)\%$		