



The LHC challenge

Andrea Romanino, SISSA

The LHC triumph

- h has $S = 0$ and $P = 1$
- h is $SU(3)_c \times U(1)_{em}$ neutral
- h belongs to $SU(2)_L$ doublet
- h couplings prop. to masses



Understanding the EW scale

- $V = \mu^2 |H|^2 + \lambda |H|^4$

- μ^2 Higgs potential parameter (tree level)


- M^2 scale of superheavy dofs with **coupling** g to H , e.g. $O(10^{16}\text{GeV})^2$

$$m_H^2 \sim -2\mu^2 + \frac{g^2}{(4\pi)^2} M^2$$

The LHC challenge (I)

- No superheavy (coupled) degrees of freedom
(finite naturalness?)
- Large sensitivity to superheavy scales, but cancellation not accidental
(environmental selection? unknown dynamics?)
- New TeV physics taming sensitivity to high scales
(supersymmetry? composite Higgs?)

The scale of “natural” new physics



$$\left. \begin{array}{l} \delta m_h^2 \\ \langle H \rangle = 174 \text{ GeV} \end{array} \right\} \delta m_h^2 \sim \frac{g^2}{(4\pi)^2} m_{\text{NP}}^2 \approx (125 \text{ GeV})^2 \left(\frac{m_{\text{NP}}}{0.5 \text{ TeV}} \right)^2$$

top contribution

$$\Delta \sim \left(\frac{m_{\text{NP}}}{0.5 \text{ TeV}} \right)^2$$

Example: supersymmetry

$$\Delta \sim \left(\frac{m_{\text{NP}}}{0.5 \text{ TeV} / \sqrt{\log}} \right)^2 \quad \log = \log \frac{M^2}{m_{\text{NP}}^2}$$


M = mediation scale

Example: supergravity

$$M = M_{\text{Planck}}$$


$$\log \sim 70$$

$$\Delta \sim 1 \text{ for } m_{\text{NP}} \sim M_Z$$




Message: low M ?

msugra



$M = 100 \text{ TeV} + \text{singlet}$



The LHC challenge (I)

- No superheavy (coupled) degrees of freedom (finite naturalness?)
- Large sensitivity to superheavy scales, but cancellation not accidental (environmental selection? unknown dynamics?)
- New TeV physics taming sensitivity to high scales (supersymmetry? composite Higgs?)

“Hints” of physics MUCH above the EW scale

- $M_{\text{Pl}} \approx 1.2 \times 10^{19} \text{ GeV}$

who knows

- Quantum number unification

give up

- Neutrino masses

The LHC challenge (I)

- No superheavy (coupled) degrees of freedom
(finite naturalness?)
- Large sensitivity to superheavy scales, but cancellation not accidental
(environmental selection? unknown dynamics?)
- New TeV physics taming sensitivity to high scales
(supersymmetry? composite Higgs?)

Environmental selection


- Give up reductionist understanding of EW scale
- Assume cosmology populates a landscape of vacua
- Retain the understanding of SM gauge quantum numbers, neutrino masses, success of gauge coupling unification, WIMP miracle

Cosmological relaxation

- Original proposal:
 - Accept field excursion up to 10^{30} GeV ($M/10\text{TeV}$)²
 - Invoke inflation model with $N \sim 10^{30}$ e-foldings
 - Non trivial low-E inflation dynamics to avoid $\theta_{\text{CD}} \sim 1$
 - Low cutoff anyway $M \approx 30\text{TeV}$
- (a starting point...)

Relaxion

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + (M^2 - \epsilon M \phi) |H|^2 + V(\epsilon \phi) + \Lambda^3 \lambda_q |H| \cos(\phi/f)$$




$$|H| \sim \frac{\epsilon}{\lambda_q} \left(\frac{M}{\Lambda} \right)^3 f \Rightarrow \epsilon \lesssim 10^{-27}$$

$$\Delta\phi \sim \frac{M}{\epsilon} \Rightarrow N_e \gtrsim 10^{30}$$


$$\theta_{\text{QCD}} \sim 1$$

A new resonance in $pp \rightarrow \gamma\gamma$?





led
crum

Basics





pp → X → γγ





→ talk by
Leandro Cieri

Basics







Significant?




ATLAS 13 TeV 3.2 fb⁻¹ 3.9 σ

ATLAS 8 TeV 20.3 fb⁻¹ 1.9 σ

CMS 13 TeV + 8 TeV 3.4 σ



Significant?





ATLAS 13 TeV only

best fit $m \approx 750 \text{ GeV}$, $\Gamma \approx 45 \text{ GeV}$

3.9 σ local

Slight preference for large Γ



CMS 13 TeV + 8 TeV



best fit $m \approx 750 \text{ GeV}$, $\Gamma \approx \text{wide}$

3.4 σ local

Slightly preference for narrow Γ

Compatible ($S = 0$)?

ATLAS 13 TeV vs 8 TeV
 8 TeV has smaller strength
 compatible at 1.2σ with 13 TeV best fit
 for $S = 0$ and gg production




ATLAS vs CMS


What if $pp \rightarrow X \rightarrow \gamma\gamma$

- Celebrate
- $\mathcal{L}_{SM} + ?$
 - which field? (spin? SM quantum numbers?)
 - which interaction? (parity? width? production and decay?)
- Who ordered that?

Spin



$\Rightarrow S = 0, 2, \dots$



ATLAS

$S = 0$ **1.2 σ (gg)**

2.1 σ (qq)

$S = 2$ **2.7 σ (gg)**

3.3 σ (qq)

8 TeV \rightarrow 13 TeV
scaling: 4.7

8 TeV \rightarrow 13 TeV
scaling: 2.7

Production

\sqrt{s}	C_{gg}	$C_{u\bar{u}}$	$C_{d\bar{d}}$	$C_{s\bar{s}}$	$C_{c\bar{c}}$	$C_{b\bar{b}}$	$C_{\gamma\gamma}$
13 TeV	2137	1054	627	83	36	15.3	54
8 TeV	174	158	89	7.2	2.7	1.07	11
σ_{13}/σ_8	4.7	2.5	2.7	4.4	5.0	5.4	1.9

- Large C allows smaller $\Gamma(X \rightarrow \gamma\gamma) \equiv \Gamma_{\gamma\gamma}$ (more plausible?)

$\sigma_{\gamma\gamma} \equiv \sigma(pp \rightarrow X \rightarrow \gamma\gamma)$ determines $C_{\mathcal{P}\mathcal{P}} \frac{\Gamma_{\mathcal{P}\mathcal{P}} \Gamma_{\gamma\gamma}}{M_X \Gamma}$ (if parton \mathcal{P} dominates)

$$\frac{\Gamma_{\gamma\gamma}}{M} \approx \frac{10^{-3}}{C_{\mathcal{P}\mathcal{P}}} \left(\frac{\Gamma}{\Gamma_{\mathcal{P}\mathcal{P}}} \right) \left(\frac{\sigma_{\gamma\gamma}}{6 \text{ fb}} \right) \approx 99 \cdot 0.5 \cdot 10^{-6} \left(\frac{\Gamma}{\Gamma_{gg}} \right) \left(\frac{\sigma_{\gamma\gamma}}{6 \text{ fb}} \right)$$

- Large σ_{13}/σ_8 ratio improves compatibility 13 TeV vs 8 TeV
- Will assume gg production when relevant

Decay and width

- $\Gamma = \Gamma_{\gamma\gamma} + \Gamma_{gg} + \Gamma_{\text{extra}}$

- large ($\Gamma/M \sim 0.06$) or narrow ($\Gamma/M \sim \text{few } 10^{-6}$)?


- $\Gamma/M \sim 0.06$

- $\Gamma/M \sim g^2/(8\pi) \rightarrow g \sim 1$


- accounted for by Γ_{extra} (tree level?)

- experimental bounds $\Gamma_{\text{extra}} / \Gamma_{\gamma\gamma} \text{ drag } \Gamma_{\gamma\gamma}/M \approx 10^{-4}$

- strongly coupled models? \rightarrow talk by Elena Vigiani



OR




exp resolution \sim **10 GeV**

$M_1^2 = M^2 + O(v^2)$

$\Gamma = 0.06 M \approx 45 \text{ GeV}$

$\Delta M_i \sim v^2/2M \approx$ **20 GeV**



$$m_a \ll M \rightarrow \gamma\gamma \sim \gamma$$

- width large because of tree level $X \rightarrow aa$
- signal large because of tree level $X \rightarrow aa$
- $\Gamma(a \rightarrow \gamma\gamma)$ “only” affects lifetime of a

Knapen Melia Papucci Zurek 1512.04928
 Agrawal Fan Heidenreich Reece Strassler 1512.05775
 Chang Cheung Lu 1512.06671
 Bi et al. 1512.08497



- Why $m_a \ll M?$ $\rightarrow a = \text{pNGB of anomalous global } U(1)$

Aparicio Azatov Hardy R 1602.00949

$$\phi = \frac{f+s}{\sqrt{2}} e^{ia/f} \quad \Gamma(s \rightarrow aa) = \frac{M^3}{32\pi f^2}$$



$f \leftrightarrow U(1)$ breaking

$$|D_\mu \phi|^2 \rightarrow \frac{s}{f} (\partial_\mu a)^2 \quad \Gamma(a \rightarrow \gamma\gamma) = \left(\frac{\alpha}{8\pi f}\right)^2 \frac{m_a^3}{\pi} (Q^2 N)^2$$

$$\Gamma = 45 \text{ GeV} \leftrightarrow f \approx 300 \text{ GeV}$$

- $m_a \approx 0.2 \text{ GeV} + L_{\text{lab}} \approx 0.6 \rightarrow \text{need } \sqrt{N}Q \gtrsim 4 \left(\frac{0.2 \text{ GeV}}{m_a}\right)$

Caveat:
mixing with π
 $m_a \approx m_\pi$




Prediction:
NO jj, YZ, ZZ, WW

Discrimination:
photon conversion
(depends on lifetime)
1602.04692

Narrow width

- Smallest $\Gamma_{\gamma\gamma}$ for i) gg production ii) dominant Γ_{gg}
 $\Gamma_{\gamma\gamma}/M \approx 0.5 \cdot 10^{-6} \rightarrow \Gamma/M \approx 10^{-3}$

- Then



($\phi \neq$ SM fields)

- Extra charged, colored degrees of freedom! \rightarrow talk by Dario Buttazzo
 - Who ordered those as well?

Who ordered X? And ψ ?

- Supersymmetry: $\text{SM} \leftrightarrow \widetilde{\text{SM}} \quad \tilde{m} > m$

- Spontaneously broken through $X = M + \frac{s + ia}{\sqrt{2}} + \theta\psi + F\theta^2$
 - sgoldstino $m = \tilde{m}$
 - goldstino $m = 0$ eaten by gravitino
 - breaks supersymmetry

- Gauginos (and sfermions) get mass by coupling to X

$$\frac{M_a}{2F} \int d^2\theta X W_a^\alpha W_\alpha = \frac{M_a}{2} \lambda_a \lambda_a + \frac{M_a}{2\sqrt{2}F} \left(s v_a^{\mu\nu} v_{\mu\nu}^a - a v_a^{\mu\nu} \tilde{v}_{\mu\nu}^a \right) + \dots$$

Bellazzini Franceschini Sala Serra 1512.05330

Petersson Torre 1512.05333

Demidov Gorbunov 1512.05723

Casas Espinosa Moreno 1512.07895

UV completion?

- $W = \lambda X \phi \bar{\phi}$ (minimal gauge mediation)
 - couple F to gluino and photino
 - couple s to gluon and photon

Baratella Elias-Miro Penedo R 1603.05682
Bardhan Byakti Ghosh Sharma 1603.05251

- Numbers: $\Gamma(s \rightarrow \gamma\gamma) = \frac{M_s^3 M_\gamma^2}{32\pi F^2}$ $M_\gamma = c_W^2 M_1 + s_W^2 M_2$


$$\sqrt{F} \lesssim 5 \text{ TeV} \left(\frac{M_\gamma}{200 \text{ GeV}} \right)^{1/2} \left(\frac{4 \text{ fb}}{\sigma_{\gamma\gamma}} \right)^{1/4}$$

- On the other hand: $M_3 = \frac{\alpha_3}{4\pi} N_3 \frac{F}{M} \rightarrow \lambda N_\gamma N_3 \gtrsim 1500 \frac{M_3}{\text{TeV}} \left(\frac{\sigma_{\gamma\gamma}}{4 \text{ fb}} \right)^{1/2}$




However

$$M_\phi = \lambda M$$



$$m_\pm^2 = M_\phi^2 \pm \lambda F$$



$$M_\phi > \sqrt{\lambda F}$$

$$g_{\text{eff}} = \frac{\lambda M_\phi}{m_-}$$

non-perturbative
near the critical point

$$M_\phi \approx \sqrt{\lambda F}$$

Veneziano NPB44 (1972)

$$(10\text{-}100) \text{ TeV} \quad \phi, \tilde{\phi}_+$$

$$O(\text{TeV}) \quad s, \tilde{\phi}_-, \tilde{g}$$

- loop suppressed by $O(\text{TeV})$
- coupling enhanced by g_{eff}

- Supersymmetric effective description fails: $\lambda M \approx F$
- Fine-tuning $\sim (g_{\text{eff}}/\lambda)^2$
- For $g_{\text{eff}} \sim 4\pi$, ϕ - bound states can form (and mix with s)
- IR non-perturbativity - does not spoil nice UV properties of supersymmetry
- $M \sim 100 \text{ TeV}$, as previously argued
- Sfermions heavier than $O(\text{TeV})$ (tree-level) if X has $U(1)_X$ charge

Example


- $\phi + \bar{\phi} \sim (3, 2, -5/6) + (\bar{3}, 2, -5/6)$ from Σ $SU(5)$ adjoint

- $N = 5$ (barely UV perturbative)

$$\frac{\Gamma_{ZZ}}{\Gamma_{\gamma\gamma}} \approx 1.3$$


$$\frac{\Gamma_{Z\gamma}}{\Gamma_{\gamma\gamma}} \approx 0$$

$$\frac{\Gamma_{WW}}{\Gamma_{\gamma\gamma}} \approx 2.8$$



Wild speculations

$$W = W_{\text{MSSM}} + \lambda X \phi \bar{\phi} + F X$$



(and an R-axion with $m_a = O(0.1 \text{ GeV}) \dots$)

Conclusions

- The LHC has confirmed what we thought about the Higgs
- Has provided a puzzle to solve: where is everybody else
- And, IF the diphoton excess turns into a new resonance, an unexpected twist
 - whose interpretation would be tremendously exciting
 - which could be the first of a series of new discoveries