New Frontiers in Theoretical Physics - XXXV Convegno Nazionale di Fisica Teorica and GGI 10th anniversary



Contribution ID: 45

Type: not specified

Susy Scattering amplitudes, Wilson loops and integrability.

Friday, 20 May 2016 11:30 (20 minutes)

The gluon scattering amplitudes and the null polygonal Wilson loops in planar Super Yang-Mills N=4 are believed to be the same. Many tests have been done at the perturbative level, while at strong coupling the scattering amplitudes can be computed solving a minamal area problem, thanks to the AdS/CFT correspondence.

As for the Wilson loops, a variant of the well-known Operator Product Expansion has been developed enabling us to write it as a sum over the excitations on the GKP vacuum. Thanks to the integrability underlying the theory, we have been obtaining many exact results in a series of papers, mainly on the strong coupling limit. Among them, the computation of the

Wilson loop is recast into a system of TBA-like equations, finding agreement with the classical string result. In addition, a purely quantum

contribution of the same order of the classical one comes out of the blue by considering an ostensibly subdominant sector.

Primary authors: BONINI, Alfredo (BO); FIORAVANTI, Davide (BO); ROSSI, Marco (CS); PISCAGLIA, Simone

Presenter: BONINI, Alfredo (BO)

Session Classification: Parallel 20 am