8th International Workshop on Semiconductor Pixel Detectors for Particles and Imaging.



Contribution ID: 59

Type: contributed paper

Design and Development of an Event-driven SOI Pixel Detector for X-ray Astronomy and Light Dark Matter Search

We have been developing monolithic active pixel detectors, named "XRPIX" based on the silicon-on-insulator (SOI) pixel technology, for future X-ray astronomical satellite missions. Our objective is to replace X-ray Charge Coupled Devices (CCD), which are now standard detectors in the field. The XRPIX series offers good time resolution (~1 μ s), fast readout time (~10 μ s), and a wide energy range (0.5–40 keV) in addition to having imaging and spectroscopic capability comparable to CCDs. XRPIX contains a comparator circuit in each pixel for hit trigger (timing) and two-dimensional hit-pattern (position) outputs. Therefore, signals are read out only from selected pixels. X-ray readout by this function is called "event-driven readout".

In our previous studies, we successfully demonstrated X-ray detection by the event-driven readout. We improved the X-ray spectral performance by introducing in-pixel charge-sensitive amplifier circuit in the frame readout mode, which an analog signal from all pixels periodically. We achieved an energy resolution of 320 eV (FWHM) for 5.9 keV X-rays with which Mn-K α and -K β lines are resolved for the first time in the XRPIX series. Recently, we designed the first prototype to achieve a large-area device for satellite loading. The detector is 24.6 mm × 15.3 mm in size and consists of 608 × 384 pixels. The pixel size and the imaging area are 36 µm × 36 µm and 21.9 mm × 13.8 mm, respectively. Moreover, We propose a light dark matter search experiment using the XRPIX. In this presentation, we report on the design and evaluation results of the new device, and the plan about a light dark matter search experiment.

Primary author: Dr TAKEDA, Ayaki (Kyoto University)

Co-authors: Mr MATSUMURA, Hideaki (Kyoto University); Mr HAYASHI, Hideki (Kyoto University); Mr KAMEHAMA, Hiroki (Shizuoka University); Dr UCHIDA, Hiroyuki (Kyoto University); Dr KAGAWA, Keiichiro (Shizuoka University); Dr YASUTOMI, Keita (Shizuoka University); Dr MIUCHI, Kentaro (Kobe University); Dr MORI, Koji (University of Miyazaki); Mr TAMASAWA, Kouki (Tokyo University of Science); Mr ITOU, Makoto (Kyoto University); Mr OKA, Naoya (Kobe University); Mr TAKEBAYASHI, Nobuaki (University of Miyazaki); Dr NAKASHIMA, Shinya (Japan Aerospace Exploration Agency); Prof. KAWAHITO, Shoji (Shizuoka University); Mr SHRESTHA, Sumeet (Shizuoka University); Mr SATO, Syou (Tokyo University of Science); Dr TANAKA, Takaaki (Kyoto University); Dr KOHMURA, Takayoshi (Tokyo University of Science); Prof. TSURU, Takeshi (Kyoto University); Prof. ARAI, Yasuo (High Energy Accelerator Research Organization, KEK); NISHIOKA, Yusuke (University of Miyazaki); Mr OZAWA, Yusuke (Tokyo University of Science)

Presenter: Dr TAKEDA, Ayaki (Kyoto University)