



Recent Results on Rare B

Decays with BaBar

Martino Margoni

Università di Padova and INFN

on behalf of the BaBar Collaboration

- Motivation

- $B \rightarrow K^* l^+ l^-$

- $B^+ \rightarrow K^+ \tau^+ \tau^-$

- $B \rightarrow K \pi \pi \gamma$




Radiative Penguins

Motivation

Rare B decays: New Physics probes

- Search for deviations from Standard Model (SM) predictions due to virtual contributions of new heavy particles in loop processes
 - Compare experimental results with very precise SM expectations
- The most interesting processes are those that are strongly suppressed in the SM: FCNC ($X_s |^{+-}$) [but also $X_s \gamma$, leptonic decays, LFV, CPV in B^0 mixing, c & τ]
 - New Physics (NP) could increase expectations by orders of magnitude [e.g. A. Buras, arXiv:0910.1032]

- Rare B decays can probe high scales potentially sensitive to NP beyond the direct reach of LHC:



$$\Lambda_{\text{NP}} \sim \frac{M_W}{g^2} \sqrt{\frac{16\pi^2}{|V_{ts}^* V_{tb}|}} \sim 10 \text{ TeV}$$

Rare B decays: New Physics probes

Weak decay of hadron M into final state F described via an Effective Hamiltonian expressed by means of Operator Product Expansion:

$$A(M \rightarrow F) = \langle F | H_{eff} | M \rangle = \frac{G_F}{\sqrt{2}} \sum_i V_{CKM}^i C_i(\mu) \langle F | Q_i(\mu) | M \rangle$$

$C_i(\mu)$: Wilson Coefficients (perturbative short distance couplings)

$Q_i(\mu)$: Hadronic Matrix Elements (non-perturbative long distance effects)

NP could modify Wilson Coefficients $C_i(\mu)$
and/or add new $Q_i(\mu)$ operators

$i = 1, 2$	Tree
$i = 3 - 6, 8$	Gluon penguin
$i = 7$	Photon penguin
$i = 9, 10$	EW penguin
$i = S, P$	(Pseudo)scalar penguin

Complementary information
from different rare decays:

$B \rightarrow \mu\mu$: Scalar/Pseudoscalar interactions

$B \rightarrow K^{(*)}\mu\mu$: Vector/axial interactions




“Measurement of Angular Asymmetries in the Decay

$B \rightarrow K^* \ell^+ \ell^-$ ”

[471 M Y(4S) events]

Phys. Rev. D93, 052015 (2016)

$$B \rightarrow K^* l^+ l^-$$




FCNC process forbidden
at tree level, $BR \sim 10^{-6}$: Probe the SM!

- Sensitive to the effects of NP in photon, vector and axial-vector couplings which can enter at the same order as SM contributions

- Complementary information to $B \rightarrow \mu^+ \mu^-$

- Amplitudes expressed using OPE in terms of:

- Hadronic Form Factors (accuracy $\sim 20\%$)
[A. Barucha et al. arXiv 1004.3249]
- Wilson coefficients C_7^{eff} , C_9^{eff} , C_{10}^{eff}
[PRD 61, 074024 (2000), Z. Phys. C 67, 417 (1995)]
- Clean theoretical predictions especially at low $q^2 = m^2(\mu^+ \mu^-)$
- Experimentally clean signature



- Kinematics of the decay $B \rightarrow V l^+ l^-$
($V=K^*$, φ , ρ) determined by three angles:
 - θ_l, θ_K, ϕ
- Event Yields reconstructed in bins of $q^2=m^2(l^+l^-)$


● Differential Amplitudes:

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos \theta_K)} = \frac{3}{2} F_L(q^2) \cos^2 \theta_K + \frac{3}{4} (1 - F_L(q^2)) (1 - \cos^2 \theta_K)$$

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos \theta_l)} = \frac{3}{4} F_L(q^2) (1 - \cos^2 \theta_l) + \frac{3}{8} (1 - F_L(q^2)) (1 + \cos^2 \theta_l) + A_{FB}(q^2) \cos \theta_l.$$

● Observables Include:

- A_{FB} (forward-backward muon asymmetry)
- F_L (fraction of longitudinally polarized K^*)
- $P_2 = \frac{-2}{3} \frac{A_{FB}}{1 - F_L}$ (with lower uncertainty from hadronic Form Factors)



- Kinematics of the decay $B \rightarrow V l^+ l^-$
($V=K^*$, φ , ρ) determined by three angles:
 - θ_l, θ_K, ϕ
- Event Yields reconstructed in bins of $q^2=m^2(l^+l^-)$

● Differential Amplitudes:

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos \theta_K)} = \frac{3}{2} F_L(q^2) \cos^2 \theta_K + \frac{3}{4} (1 - F_L(q^2)) (1 - \cos^2 \theta_K)$$

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos \theta_l)} = \frac{3}{4} F_L(q^2) (1 - \cos^2 \theta_l) + \frac{3}{8} (1 - F_L(q^2)) (1 + \cos^2 \theta_l) + A_{FB}(q^2) \cos \theta_l.$$

- Non-resonant S-wave $B \rightarrow K \pi l^+ l^-$ contribution neglected
 - Reflects in absolute bias ~ 0.01 on F_L & A_{FB} (smaller than statistical & systematic uncertainties)



• Measurement performed using 5 modes:

➤ $B^+ \rightarrow K^{*+} (\rightarrow K_s \pi^+) \mu^+ \mu^-$, $B^+ \rightarrow K^{*+} (\rightarrow K_s \pi^+) e^+ e^-$, $B^+ \rightarrow K^{*+} (\rightarrow K^+ \pi^0) e^+ e^-$

➤ $B^0 \rightarrow K^{*0} (\rightarrow K^+ \pi^-) \mu^+ \mu^-$, $B^0 \rightarrow K^{*0} (\rightarrow K^+ \pi^-) e^+ e^-$

➤ $K^* J/\psi$ and $K^* \psi(2S)$ regions used as control samples to validate fitting procedure

q^2 bin	q^2 min (GeV ² /c ⁴)	q^2 max (GeV ² /c ⁴)
q_1^2	0.10	2.00
q_2^2	2.00	4.30
q_3^2	4.30	8.12
q_4^2	10.11	12.89
q_5^2	14.21	$(m_B - m_{K^*})^2$
q_0^2	1.00	6.00

• Events reconstructed by means of :

$$m_{ES} = \sqrt{E_{Beam}^{*2} - p_B^{*2}}$$

$$\Delta E = E_B^* - E_{Beam}^*$$



* = Υ reference frame

Candidate multiplicity ~ 1.4
(1.1) in dielectron (dimuon) modes.

Best candidate selected based on ΔE



• Measurement performed using 5 modes:









➤ $K^* J/\psi$ and $K^* \psi(2S)$ regions used as control samples to validate fitting procedure

q^2 bin	q^2 min (GeV^2/c^4)	q^2 max (GeV^2/c^4)
q_1^2	0.10	2.00
q_2^2	2.00	4.30
q_3^2	4.30	8.12
q_4^2	10.11	12.89
q_5^2	14.21	$(m_B - m_{K^*})^2$
q_0^2	1.00	6.00

BKG from Continuum and $B\bar{B}$ reduced using a Likelihood Ratio (L_R) defined from outputs of eight BDTs exploiting kinematical and topological quantities



● Yields, PDFs shapes & normalizations in the different q^2 bins extracted by a 3D (m_{ES} , $m(K\pi)$, L_R) fit Example: $q^2 > 14.21 \text{ GeV}^2$







- F_L in the different q^2 bins extracted as only free parameter by a $4D(m_{ES}, m(K\pi), L_R, \cos(\theta_K))$ fit using PDFs defined in the previous step
- Fit model for F_L and A_{FB} validated on $K^* J/\psi$ and $K^* \psi(2S)$
- BKG shapes from m_{ES} side bands (checked on LFV $B \rightarrow K^* e\mu$)

First $B^+ \rightarrow K^{*+} l^+ l^-$ angular analysis

Example: $1 < q^2 < 6 \text{ GeV}^2$

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos\theta_K)} = \frac{3}{2} F_L(q^2) \cos^2\theta_K + \frac{3}{4} (1 - F_L(q^2)) (1 - \cos^2\theta_K)$$







- A_{FB} in the different q^2 bins extracted as only free parameter by a $4D(m_{ES}, m(K\pi), L_R, \cos(\theta_l))$ fit using PDFs defined in the previous step
- Γ_L fixed to previous result

First $B^+ \rightarrow K^{*+} l^+ l^-$ angular analysis



Example: $1 < q^2 < 6 \text{ GeV}^2$

$$\frac{1}{\Gamma(q^2)} \frac{d\Gamma}{d(\cos \theta_\ell)} = \frac{3}{4} F_L(q^2) (1 - \cos^2 \theta_l) + \frac{3}{8} (1 - F_L(q^2)) (1 + \cos^2 \theta_l) + A_{FB}(q^2) \cos \theta_l.$$






Angular Variables Results versus q^2 : Tension with SM in $B^+ F_L$ at low q^2





● Angular Variable Results for $1 < q^2 < 6 \text{ GeV}^2$




- $1 < q^2 < 6 \text{ GeV}^2$: Perturbative window with theory error under good control, away from $q^2 \rightarrow 0$ photon pole and $c\bar{c}$ resonances at higher q^2
- Small F_L value for $B^+ \rightarrow K^{*+} l^+ l^-$ (First Angular Analysis)



- $P_2 = \frac{-2}{3} \frac{A_{FB}}{1 - F_L}$: Reduced theoretical uncertainty & greater sensitivity to non-SM contributions

[Nucl. Phys. B854, 321 (2012); JHEP 1204, 104 (2012); Phys. Rev. D88, 074002 (2013); JHEP 1412, 125 (2014)]




- Theoretical predictions available only at low q^2 [JHEP 1412, 125 (2014)]
- Slight tension observed with SM




- $P_2 = \frac{-2}{3} \frac{A_{FB}}{1 - F_L}$: Reduced theoretical uncertainty & greater sensitivity to non-SM contributions

[Nucl. Phys. B854, 321 (2012); JHEP 1204, 104 (2012); Phys. Rev. D88, 074002 (2013); JHEP 1412, 125 (2014)]



- Result dominated by statistical error
- Systematics from BKG modeling, signal angular efficiency, PDFs parameterization & cross feed from different signal decays



“Search for $B^+ \rightarrow K^+ \tau^+ \tau^-$ at the BaBar Experiment”


[471 M Y(4S) events]

arXiv:1605.09637

Submitted to Phys. Rev. Lett.




- Highly suppressed in the SM: $BR \sim (1-2)10^{-7}$
- Provides additional sensitivity to New Physics due to third-generation couplings & large τ mass



2HDM


J. Phys. G24, 49 (1998)

	m_{h^0} (GeV)	m_{H^\pm} (GeV)	m_{H^0} (GeV)	m_{A^0} (GeV)
Mass set-1	80	200	150	100
Mass set-2	250	300	100	350
Mass set-3	100	400	200	150





- Measurement performed using only leptonic τ decays:
 - $B^+ \rightarrow K^+ \tau^+ \tau^-$, $\tau \rightarrow \mu \nu_\tau \nu_\mu$, $\tau \rightarrow e \nu_\tau \nu_e$
 - Three signal modes: ee , $\mu\mu$, $e\mu$
- Many neutrinos in the final states: lack of kinematic constraints
 - Signal events selected on the recoil of fully reconstructed hadronic $B \rightarrow DX$ decays (B_{tag}) ($D = D^{(*)0}, D^{(*)\pm}, D_s^{(*)}$, J/ψ ; $X < 6 h$ ($h=K, \pi$))





● B_{tag} Reconstruction

● B hadronic decays selected by means of m_{ES} & ΔE

● Best candidate per event retained from the highest purity mode (computed from MC) & ΔE

➤ Only B_{tag} candidates with Purity > 40% used $\rightarrow \epsilon(B_{\text{tag}}) = (0.2 - 0.4)\%$

● Continuum events suppressed by exploiting a Likelihood Selector consisting of six event-shape variables (e.g. Thrust, missing momentum vector, $P(B_{\text{tag}})$, angles between them,...)


➤ $LS > 0.5$ removes > 75% of BKG retaining 80% of the signal



- $B \rightarrow K^+ \tau^+ \tau^-$ Reconstruction

- Signal candidates reconstructed from events with three charged particles, identified as K + two leptons, not belonging to B_{tag}
- Vetos applied against J/ψ , $D^0 \rightarrow K\pi$ ($\rightarrow \mu$), $\gamma \rightarrow e^+e^-$, $\pi^0 \rightarrow \gamma\gamma$

- Dominant BKG from $B \rightarrow D^{(*)} l \nu$, $D^{(*)} \rightarrow K l' \nu$ (same final-state) suppressed by a Neural Network using angles between momenta, $m(K^+ l^-)$ and missing energy





- BR for each of the signal modes:

$$B_i = \frac{N_{\text{obs}}^i - N_{\text{bkg}}^i}{\epsilon_{\text{sig}}^i N_{B\bar{B}}} \quad N_{B\bar{B}} = 471 \times 10^6$$

	$e^+ e^-$	$\mu^+ \mu^-$	$e^+ \mu^-$
N_{bkg}^i	$49.4 \pm 2.4 \pm 2.9$	$45.8 \pm 2.4 \pm 3.2$	$59.2 \pm 2.8 \pm 3.5$
$\epsilon_{\text{sig}}^i (\times 10^{-5})$	$1.1 \pm 0.2 \pm 0.1$	$1.3 \pm 0.2 \pm 0.1$	$2.1 \pm 0.2 \pm 0.2$
N_{obs}^i	45	39	92
Significance (σ)	-0.6	-0.9	3.7

- Signal efficiencies and expected Peaking BKG events (92%) obtained from simulation corrected to reproduce B_{tag} data yield
- Expected combinatorial BKG events (8%) from data m_{ES} Side Band



- BR for each of the signal modes:

$$\mathcal{B}_i = \frac{N_{\text{obs}}^i - N_{\text{bkg}}^i}{\epsilon_{\text{sig}}^i N_{B\bar{B}}}$$

$$N_{B\bar{B}} = 471 \times 10^6$$

	e^+e^-	$\mu^+\mu^-$	$e^+\mu^-$
N_{bkg}^i	$49.4 \pm 2.4 \pm 2.9$	$45.8 \pm 2.4 \pm 3.2$	$59.2 \pm 2.8 \pm 3.5$
$\epsilon_{\text{sig}}^i (\times 10^{-5})$	$1.1 \pm 0.2 \pm 0.1$	$1.3 \pm 0.2 \pm 0.1$	$2.1 \pm 0.2 \pm 0.2$
N_{obs}^i	45	39	92
Significance (σ)	-0.6	-0.9	3.7

- e^+e^- , $\mu^+\mu^-$ yields show consistency with expected BKG events.
- $e\mu$ channel has excess of 3.7 σ :
 - No evident signal-like behaviour or systematic problems from kinematic distributions



- BR for each of the signal modes:

$$B_i = \frac{N_{\text{obs}}^i - N_{\text{bkg}}^i}{\epsilon_{\text{sig}}^i N_{B\bar{B}}} \quad N_{B\bar{B}} = 471 \times 10^6$$

	e^+e^-	$\mu^+\mu^-$	$e^+\mu^-$
N_{bkg}^i	$49.4 \pm 2.4 \pm 2.9$	$45.8 \pm 2.4 \pm 3.2$	$59.2 \pm 2.8 \pm 3.5$
$\epsilon_{\text{sig}}^i (\times 10^{-5})$	$1.1 \pm 0.2 \pm 0.1$	$1.3 \pm 0.2 \pm 0.1$	$2.1 \pm 0.2 \pm 0.2$
N_{obs}^i	45	39	92
Significance (σ)	-0.6	-0.9	3.7

- Overall significance $< 2\sigma$:

$BR(B^+ \rightarrow K^+ \tau^+ \tau^-) < 2.25 \times 10^{-3}$ (90% CL) First Measurement

- Systematics from B_{tag} yield correction, theoretical models for efficiency determination, PID, and Data/MC agreement

$$\mathcal{B} \rightarrow \text{K}\pi\pi\gamma$$


“Time-dependent analysis of $B^0 \rightarrow K_S \pi^- \pi^+ \gamma$ and studies of the $K^+ \pi^- \pi^+$ system in $B^+ \rightarrow K^+ \pi^- \pi^+ \gamma$ decays”

[471 MY(4S) events]

Phys. Rev. D93, 052013 (2016)


Radiative decays and the γ polarization

- $b \rightarrow s\gamma$: described in the SM as an interaction between left-handed quarks and right-handed antiquarks:




$$\lambda = \frac{\vec{S} \cdot \vec{p}}{|\vec{p}|}$$

- $b \rightarrow s\gamma_L$ & $\bar{b} \rightarrow \bar{s}\gamma_R$




- Mixing induced CP Asymmetry = 0

- New heavy particles in the loop could enhance opposite helicity γ contribution



- $b \rightarrow s\gamma_{L/R}$ & $\bar{b} \rightarrow \bar{s}\gamma_{R/L}$



- Mixing induced CP Asymmetry $\neq 0$

Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$

$$A_{CP}(\Delta t) = \frac{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}\gamma) - \Gamma(B^0(\Delta t) \rightarrow f_{CP}\gamma)}{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}\gamma) + \Gamma(B^0(\Delta t) \rightarrow f_{CP}\gamma)}$$

$$= S_{f_{CP}} \sin(\Delta m_d \Delta t) - C_{f_{CP}} \cos(\Delta m_d \Delta t)$$



- SM predicts $S_{f_{CP}} = m_s/m_b = 0.02$
- Look for enhancement due to new-particle exchange

$[\Delta t = t_{\text{Rec}} - t_{\text{Tag}}$ from distance between the two B^0 decay vertices in the event]

- Experimentally: perform a time-dependent analysis of $B^0 \rightarrow K_S \rho \gamma$

- Main Issue: dilution from irreducible BKG from non CP eigenstates:

CP eigenstate $B^0 \rightarrow K_S \rho \gamma$ Non CP eigenstate $B^0 \rightarrow K^*(K_S \pi) \pi \gamma$



Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$

$$A_{CP}(\Delta t) = \frac{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}\gamma) - \Gamma(B^0(\Delta t) \rightarrow f_{CP}\gamma)}{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}\gamma) + \Gamma(B^0(\Delta t) \rightarrow f_{CP}\gamma)}$$

$$= S_{f_{CP}} \sin(\Delta m_d \Delta t) - C_{f_{CP}} \cos(\Delta m_d \Delta t)$$

- SM predicts $S_{f_{CP}} = m_s/m_b = 0.02$
- Look for enhancement due to new-particle exchange

$[\Delta t = t_{\text{Rec}} - t_{\text{Tag}}$ from distance between the two B^0 decay vertices in the event]

- Experimentally: perform a time-dependent analysis of $B^0 \rightarrow K_S \rho \gamma$

- Main Issue: dilution from irreducible BKG from non CP eigenstates:

CP eigenstate $B^0 \rightarrow K_S \rho \gamma$ Non CP eigenstate $B^0 \rightarrow K^*(K_S \pi)\pi \gamma$

Dilution: $D_{K_S^0 \rho \gamma} \equiv \frac{S_{K_S^0 \pi^+ \pi^- \gamma}}{S_{K_S^0 \rho \gamma}}$

Effective value on inclusive $K_S \pi \pi \gamma$ sample

Signal value

Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$

- Dilution expressed in terms of few resonant decay modes:

$\rho^0 K_S$, $K^{*+} \pi^-$, $K^{*-} \pi^+$, $(K\pi)_0^{*+} \pi^-$, $(K\pi)_0^{*-} \pi^+$ S-wave ($K^*_0(1430)$ + NR

component) and their **interference**:

$$D_{K_S^0 \rho \gamma} \equiv \frac{\mathcal{S}_{K_S^0 \pi^+ \pi^- \gamma}}{\mathcal{S}_{K_S^0 \rho \gamma}} = \frac{\int \left[|A_{\rho K_S^0}|^2 - |A_{K^{*+} \pi^-}|^2 - |A_{(K\pi)_0^{*+} \pi^-}|^2 + 2\Re(A_{\rho K_S^0}^* A_{K^{*+} \pi^-}) + 2\Re(A_{\rho K_S^0}^* A_{(K\pi)_0^{*+} \pi^-}) \right] dm^2}{\int \left[|A_{\rho K_S^0}|^2 + |A_{K^{*+} \pi^-}|^2 + |A_{(K\pi)_0^{*+} \pi^-}|^2 + 2\Re(A_{\rho K_S^0}^* A_{K^{*+} \pi^-}) + 2\Re(A_{\rho K_S^0}^* A_{(K\pi)_0^{*+} \pi^-}) \right] dm^2}$$

[LAL-15-75 (2015)]

- Ideal World:** Perform a time-dependent **Amplitude Analysis**
- Real World:** Not enough statistics, dilution computed from the amplitudes of the intermediate resonances from $B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ assuming **Isospin Symmetry**

$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ Selection

● $B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ events selected by means of:





➤ $1.5 < E_\gamma^* < 3.5 \text{ GeV}$

* = Υ reference frame

➤ m_{ES}

➤ ΔE

● Continuum BKG suppressed using a Fisher exploiting topological quantities



$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ Selection


- $B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ signal yield extracted from an unbinned fit to m_{ES} , ΔE and Fisher discriminant:

$$\pm N_{sig} = 2441 \pm 91^{+41}_{-54}$$

$$\pm BF(B^+ \rightarrow K^+ \pi^+ \pi^- \gamma) = (24.5 \pm 0.9 \pm 1.2) 10^{-6}$$


- $m(K\pi\pi)$, $m(K\pi)$ and $m(\pi\pi)$ spectra obtained using s Plot technique

[NIM A 555, 356-369 (2005)]



$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ Analysis


- BFs of the various resonances decaying to $K\pi\pi$ extracted from the $m(K\pi\pi)$ spectrum



Mode	$\mathcal{B}(B^+ \rightarrow \text{Mode}) \times \mathcal{B}(K_{\text{res}} \rightarrow K^+ \pi^+ \pi^-) \times 10^{-6}$	$\mathcal{B}(B^+ \rightarrow \text{Mode}) \times 10^{-6}$	Previous world average [18] ($\times 10^{-6}$)
$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$...	$24.5 \pm 0.9 \pm 1.2$	27.6 ± 2.2
$K_1(1270)^+ \gamma$	$14.5^{+2.1+1.2}_{-1.4-1.2}$	$44.1^{+6.3+3.6}_{-4.4-3.6} \pm 4.6$	43 ± 13
$K_1(1400)^+ \gamma$	$4.1^{+1.9+1.2}_{-1.2-1.0}$	$9.7^{+4.6+2.8}_{-2.9-2.3} \pm 0.6$	<15 at 90% C.L.
$K^*(1410)^+ \gamma$	$11.0^{+2.2+2.1}_{-2.0-1.1}$	$27.1^{+5.4+5.2}_{-4.8-2.6} \pm 2.7$	n/a
$K_2^*(1430)^+ \gamma$	$1.2^{+1.0+1.2}_{-0.7-1.5}$	$8.7^{+7.0+8.7}_{-5.3-10.4} \pm 0.4$	14 ± 4
$K^*(1680)^+ \gamma$	$15.9^{+2.2+3.2}_{-1.9-2.4}$	$66.7^{+9.3+13.3}_{-7.8-10.0} \pm 5.4$	<1900 at 90% C.L.

$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ Analysis


- Extraction of the dilution from amplitudes of intermediate states decaying to $K\pi$ and $\pi\pi$
- Full amplitude analysis in the $m(K\pi)$ - $m(\pi\pi)$ not possible due to small statistics
 - ✚ Perform a 1D fit to $m(K\pi)$ using as inputs the BRs obtained from the $m(K\pi)$ fit



Mode	$\frac{\mathcal{B}(B^+ \rightarrow \text{Mode}) \times \mathcal{B}(R \rightarrow h\pi)}{\mathcal{B}(R \rightarrow h\pi) \times 10^{-6}}$	$\mathcal{B}(B^+ \rightarrow \text{Mode}) \times 10^{-6}$	Previous world average [18] ($\times 10^{-6}$)
$K^*(892)^0 \pi^+ \gamma$	$15.6 \pm 0.6 \pm 0.5$	$23.4 \pm 0.9^{+0.8}_{-0.7}$	20^{+7}_{-6}
$K^+ \rho(770)^0 \gamma$	$8.1 \pm 0.4^{+0.8}_{-0.7}$	$8.2 \pm 0.4 \pm 0.8 \pm 0.02$	< 20 at 90% CL
$(K\pi)_0^{*0} \pi^+ \gamma$	$10.3^{+0.7+1.5}_{-0.8-2.0}$...	n/a
$(K\pi)_0^0 \pi^+ \gamma$ (NR)	...	$9.9 \pm 0.7^{+1.5}_{-1.9}$	< 9.2 at 90% CL
$K_0^*(1430)^0 \pi^+ \gamma$	$0.82 \pm 0.06^{+0.12}_{-0.16}$	$1.32^{+0.09+0.20}_{-0.10-0.26} \pm 0.14$	n/a

$B^+ \rightarrow K^+ \pi^+ \pi^- \gamma$ Analysis

- Extraction of the dilution from amplitudes of intermediate states decaying to $K\pi$ and $\pi\pi$
- Full amplitude analysis in the $m(K\pi)$ - $m(\pi\pi)$ not possible due to small statistics
 - ✦ Perform a 1D fit to $m(K\pi)$ using as inputs the BRs obtained from the $m(K\pi\pi)$ fit



$$D_{K_S^0 \rho \gamma} = F(A_\rho, A_{K^*}, A_{(K\pi)S\text{-wave}}) = -0.78^{+0.19}_{-0.17}$$

Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$




- Time-dependent analysis of $B^0 \rightarrow K_S \rho \gamma$ decays
- Event yield and CP parameters C and S extracted from a fit to m_{ES} , ΔE , Fisher and Δt
- Sample divided in 6 mutually exclusive tagging categories c

$$\mathcal{P}_j^i(m_{ES}, \Delta E, \mathcal{F}, \Delta t, \sigma_{\Delta t}; q_{\text{tag}}, \mathbf{C})$$

$$= \mathcal{P}_j^i(m_{ES}) \mathcal{P}_j^i(\Delta E) \mathcal{P}_j^i(\mathcal{F}) \mathcal{P}_j^i(\Delta t, \sigma_{\Delta t}; q_{\text{tag}}, \mathbf{C})$$


$c = \text{tagging category}$

$q_{\text{tag}} = 1$ ($B_{\text{Tag}} = B^0$)
 -1 ($B_{\text{Tag}} = \overline{B^0}$)



Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$

- Time-dependent analysis of $B^0 \rightarrow K_S \rho \gamma$ decays
- Event yield and CP parameters C and S extracted from a fit to m_{ES} , ΔE , Fisher and Δt
- Sample divided in 6 mutually exclusive tagging categories c



$$\begin{aligned}
 & \mathcal{P}_{\text{sig}}^i(\Delta t, \sigma_{\Delta t}; q_{\text{tag}}, c) \\
 &= \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left[1 + q_{\text{tag}} \frac{\Delta D_c}{2} \right. \\
 & \quad \left. + q_{\text{tag}} \langle D \rangle_c \left(S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t) \right) \right] \\
 & \otimes \mathcal{R}_{\text{sig}}^c(\Delta t, \sigma_{\Delta t}),
 \end{aligned}$$

ΔD : Difference between D for B^0 and $\overline{B^0}$ tags

c = tagging category
 $q_{\text{tag}} = 1$ ($B_{\text{tag}} = B^0$)
 -1 ($B_{\text{tag}} = \overline{B^0}$)

- Tagging imperfection D , ΔD & Δt Resolution \mathcal{R}_{sig} from quarkonium $\sin(2\beta)$ analysis [PRL 99, 171803 (2007)]

Measurement of A_{CP} in $B^0 \rightarrow K_S \rho \gamma$

Results:

$$\text{BF}(B^0 \rightarrow K_S \pi \pi \gamma) = (20.5 \pm 2.0^{+2.6}_{-2.2}) 10^{-6}$$

$$S_{K_S \pi \pi \gamma} = 0.14 \pm 0.25 \pm 0.03$$

$$C_{K_S \pi \pi \gamma} = -0.39 \pm 0.20^{+0.03}_{-0.02}$$

• After correcting for $\mathcal{D}_{K_S^0 \rho \gamma}$:

$$\oplus S_{K_S \rho \gamma} = -0.18 \pm 0.32^{+0.06}_{-0.05}$$

Consistent
with SM

- Systematics from resonance modelling and ΔE , m_{ES} and Fisher distributions shape
- Results consistent with Belle [PRL 101, 251601 (2008)]

Conclusions

Conclusions

- Rare B decays are an excellent laboratory for the search for physics beyond the SM
- In the last few years several new measurements from LHC & B-Factories experiments released with impressive experimental precision
- Almost all the results are in agreement with expectations **but some tension is present in some sectors: BaBar F_L for $B^+ \rightarrow K^{*+}l^+l^-$ (shown today), $B \rightarrow K^{(*)}\mu\mu$ ($P5'$, $BR(B \rightarrow K\mu\mu)/BR(B \rightarrow Kee)$), (but also $B \rightarrow D^{(*)}\tau\nu/B \rightarrow D^{(*)}\mu\nu$)**

- **Strong constraints on NP models from flavor measurements**
- Rich program of measurements is expected from LHC/Belle II experiments in the coming years
 - **Chances to discover/understand NP in the flavor sector in the near future?**

Backup

$$B \rightarrow K^* \ell^+ \ell^-$$



- Event Yields:

Mode	q_0^2	q_1^2	q_2^2	q_3^2	q_4^2	q_5^2
$B \rightarrow K^* \ell^+ \ell^-$	40.8 ± 8.4	31.7 ± 7.1	11.9 ± 5.5	21.3 ± 8.5	31.9 ± 9.2	33.2 ± 7.8
$B^+ \rightarrow K^{*+} \ell^+ \ell^-$	17.7 ± 5.2	8.7 ± 4.1	3.8 ± 4.0	7.7 ± 5.6	9.0 ± 4.8	9.4 ± 4.2
$B^0 \rightarrow K^{*0} \ell^+ \ell^-$	23.1 ± 6.6	22.9 ± 5.8	8.1 ± 3.8	13.7 ± 6.4	22.8 ± 7.8	23.8 ± 6.6

- Systematics:




- PDF shapes and parameter statistical error
- F_L statistical error propagated in A_{FB} fit
- Modeling of BKG PDF shape and Signal efficiency
- Signal crossfeed
- Fit bias
- Stability vs cuts

$B \rightarrow K^* \ell \ell$ Angular Analysis: P_5' parameter



- LHCb full statistics result on P_5' : discrepancy at 3.4σ level [JHEP 02, 104 (2016)]
- Belle confirms the tension at 2.1σ level [arXiv:1604.04042]
- Need to control the charm penguin to disentangle SM from NP in C_7^{eff} and C_9^{eff}

$B \rightarrow K^* \mu^+ \mu^-$: CMS Results




Results consistent with SM




- + Systematics from BKG PDF shapes, efficiency, simulation mismodeling and fit bias.
- + Theoretical predictions:
 - + Light-cone sum rules at low q^2 and extrapolation at high q^2 [JHEP 09 089 (2010), JHEP 02 010 (2013)]
 - + Lattice [Phys. Rev. D89 094501 (2014)]

$B \rightarrow K^*$ II Related quantities


● $K^* \mu^+ \mu^-$ tension motivates studies of differential BRs



[JHEP 06, 133 (2014)]



[JHEP 09, 179 (2015)]




[JHEP 06, 115 (2015)]

● All the results are “consistent” with SM at $< 2.2 \sigma$

● But all of them are lower than the predictions...



● Measurements of related $b \rightarrow d \mu \mu$ channels very useful to reveal information on Minimal Flavor Violation nature of New Physics



LHCb [JHEP 10, 034 (2015)]:

$BR(B^+ \rightarrow \pi^+ \mu^+ \mu^-) = (1.83 \pm 0.24 \pm 0.05) 10^{-8}$ in agreement with MFV

$BR(B^+ \rightarrow \pi^+ \mu^+ \mu^-) / BR(B^+ \rightarrow K^+ \mu^+ \mu^-) = 0.037 \pm 0.008 \pm 0.001$

$|V_{td}| / |V_{ts}| = 0.24^{+0.05}_{-0.04}$ in agreement with box processes ($\Delta m_s / \Delta m_d$) results




- Signal efficiencies and expected Peaking BKG events (92%) from simulation corrected according to Data/MC ratio before NN cut:

$$\left(\frac{N^{Data}}{N^{MC}} \right)_{BKG} = 0.913 \pm 0.020$$

- Expected combinatorial BKG events (8%) from data mEs Side Band

- Data/MC B_{tag} yields cross-checked using $B^+ \rightarrow D^0 | \nu$, $D^0 \rightarrow K\pi$ (before NN cut)





- Cross checks to understand the excess:

- Excess present also in the B_{tag} side band

- Discriminating variable in the NN:

- $s_B = \frac{q^2}{m_B^2} = \frac{(p_{B_{\text{sig}}} - p_K)^2}{m_B^2}$, $m(K^+ \tau^-)$, K - τ angle in the di-tau frame, lepton momentum, missing energy, e.m. energy not associated to B_{tag}

- All of them compatible with BKG statistical fluctuation



Systematics:

- Theory (signal efficiency): 3% from shape of the q^2 distribution (Lattice QCD vs light cone sum rules)
- Btag Yield: 1.5% from MC correction using m_{ES} sideband
- PID: 5% from Data/MC comparison
- π^0 Veto: 3%
- NN cut: 2.6% from Data/MC checked on $B^+ \rightarrow D^0 \nu$ ($D^0 \rightarrow K\pi$)

Measurement of A_{CP} in $B^0 \rightarrow K_S \pi \gamma$

Belle [PRL 101, 251601 (2008)]:

$$S_{K_S \pi \gamma} = 0.11 \pm 0.33^{+0.05}_{-0.09}$$

$$A_{CP}(\text{direct}) = 0.05 \pm 0.18 \pm 0.06$$

LHCb [PRL 112, 161901 (2014)]:



$$A_{ud} \equiv \frac{\int_0^1 d \cos \theta \frac{d\Gamma}{d \cos \theta} - \int_{-1}^0 d \cos \theta \frac{d\Gamma}{d \cos \theta}}{\int_{-1}^1 d \cos \theta \frac{d\Gamma}{d \cos \theta}}$$

θ = angle between photon and $K\pi\pi$ plane normal

$m_{K\pi\pi}$	[1.1,1.3]	[1.3,1.4]	[1.4,1.6]	[1.6,1.9]
A_{ud}	6.9 ± 1.7	4.9 ± 2.0	5.6 ± 1.8	-4.5 ± 1.9

5.2 σ significance for nonzero up-down asymmetry

First measurement



Capri 2016, 11-13 June 2016