


Crystalline condensates in dense quark matter




Stefano Carignano

CSQCD V, May 2016


The QCD phase diagram



The QCD phase diagram




The QCD phase diagram



Compact stars: even more complicated!


Compact stars: even more complicated!

- Charge neutrality



Compact stars: even more complicated!

- Charge neutrality



- Magnetic fields
-


Pairing mechanisms in quark matter

Pairing mechanisms in quark matter

Vacuum: quark-antiquark pairing
→ homogeneous chiral symmetry breaking

Pairing mechanisms in quark matter


Vacuum: quark-antiquark pairing
→ homogeneous chiral symmetry breaking



becomes disfavored at high chemical potentials


Pairing mechanisms in quark matter

High density: diquark pairing
→ color-superconductivity




Pairing mechanisms in quark matter

Intermediate densities: quark-hole pairing (opposite momenta)
→ exciton



Pairing mechanisms in quark matter

Intermediate densities: quark-hole pairing (equal momenta)
→ inhomogeneous chiral symmetry breaking



Why inhomogeneous phases ?

- Popular already for quite some time...
 - Overhauser pairing in nuclear matter
 - Pion condensation
 - (Color-) Superconductivity

- Recently rediscovered and revised
 - Studies of lower-dimensional models (GN_2 , NJL_2 , ...)
 - Quarkyonic chiral spirals
 - ...

- May be relevant for cold dense matter !

Formalism

- Nambu–Jona-Lasinio model

$$\mathcal{L}_{NJL} = \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi + G \left((\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma^5\tau^a\psi)^2 \right)$$

- Chirally invariant four-fermion interaction
- Mean-field approximation \rightarrow Thermodynamic potential

Formalism

- Nambu–Jona-Lasinio model

$$\mathcal{L}_{NJL} = \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi + G \left((\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma^5\tau^a\psi)^2 \right)$$

- Chirally invariant four-fermion interaction
- Mean-field approximation \rightarrow Thermodynamic potential
- Inhomogeneous phases:
retain spatial dependence of the condensates

$$\langle \bar{\psi}\psi \rangle = S(\mathbf{x}), \quad \langle \bar{\psi}i\gamma^5\tau_3\psi \rangle = P(\mathbf{x})$$

- Minimize thermodynamic potential $\Omega(T, \mu; S, P)$


What kind of modulation?

- Tackling an arbitrary spatial modulation is extremely complicated!

What kind of modulation?

- Tackling an arbitrary spatial modulation is extremely complicated!
- In practice: choose a specific ansatz for the spatial dependence
- Typical examples:
 - Chiral density wave: $S(\mathbf{x}) \sim \Delta \cos(\mathbf{q} \cdot \mathbf{x})$, $P(\mathbf{x}) \sim \Delta \sin(\mathbf{q} \cdot \mathbf{x})$
 - Real cosine: $S(\mathbf{x}) \sim \Delta \cos(\mathbf{q} \cdot \mathbf{x})$, $P(\mathbf{x}) = 0$
 - Real kink crystal: $S(\mathbf{x}) \sim \Delta \nu \operatorname{sn}(\Delta z, \nu)$, $P(\mathbf{x}) = 0$
 - 2D solutions: $S(\mathbf{x}) \sim \Delta \cos(qx) \cos(qy)$, $P(\mathbf{x}) = 0$

What kind of modulation?




- Real kink crystal is (usually) the most favored solution
- Qualitatively similar results for order parameters / phase diagrams for all modulations


CDW order parameters at $T = 0$

- Chiral density wave: $S(\mathbf{x}) \sim \Delta \cos(\mathbf{q} \cdot \mathbf{x})$, $P(\mathbf{x}) \sim \Delta \sin(\mathbf{q} \cdot \mathbf{x})$


$$M(\mathbf{x}) = -2G(S(\mathbf{x}) + iP(\mathbf{x})) \equiv \Delta e^{i\mathbf{q} \cdot \mathbf{x}}$$



Phase diagram




Phase diagram



Model dependence?


Model dependence?

Quark-meson model (“renormalized” extended MFA)



Model dependence?

Dyson-Schwinger methods



Extended NJL for realistic compact star description

- Magnetic fields
- Density effects (“vector interactions”)
- Interplay with color-superconductivity
- Charge neutrality


Magnetic fields

- Effective dimensional reduction at the lowest Landau level (LLL)
- “Famous” consequence: Magnetic catalysis


Magnetic fields

- Effective dimensional reduction at the lowest Landau level (LLL)
- “Famous” consequence: Magnetic catalysis
- Dimensional reduction to 1+1D \rightarrow Inhomogeneous χ SB
- Known examples:
 - 1+1-dimensional QCD
 - NJL₂ model
 - Quarkyonic matter (\rightarrow QCD₂)

Order parameters



Order parameters



Vector interactions

- Vector interactions expected to play a role at finite densities
- Extend NJL Lagrangian by adding


$$\mathcal{L}_V = -G_V \sum_a (\bar{\psi} \gamma^\mu \tau_a \psi)^2$$

- Mean-field - amounts to a shift in the chemical potentials


$$\mu_f \rightarrow \tilde{\mu}_f = \mu_f - 4G_V \rho_f$$

- G_V free parameter


Vector interactions




Vector interactions




Phase diagram - vector interactions




Phase diagram - vector interactions



Phase diagram - vector interactions



Interplay with (homogeneous) color-superconductivity




Equation of state

- Inhomogeneous window appears at finite density
- What are its effects on the EoS ?

Equation of state



- Inhomogeneous window appears at finite density
- What are its effects on the EoS ?




Equation of state

- Inhomogeneous window appears at finite density
- What are its effects on the EoS ?

Compared with the effect of vector interactions...



M(R) sequences for inhomogeneous quark matter



Conclusions


- Inhomogeneous phases likely to form in dense quark matter

Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!


Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!




Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!



Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!



Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!

- Experimental signatures ?

Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!
- Experimental signatures ?
 - $M(R)$ sequences

Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!

- Experimental signatures ?
 - $-M(R)$ sequences
 - Neutrino emission
 - ...


Conclusions

- Inhomogeneous phases likely to form in dense quark matter
- Need to rethink the QCD phase diagram!

- Experimental signatures ?
 - ~~M(R)~~ sequences
 - Neutrino emission
 - ...

- Reviews:
 - Prog. Part. Nucl. Phys 81 (2015) , arXiv:1406.1367
 - EPJA Topical Issue "Exotic matter in NS" (2016), arXiv:1508.04361

phase diagram + csc



M(R) sequences - vector interactions

