Contribution ID: **84** Type: **Oral**

Measurements Of Stellar And Big-Bang Nucleosynthesis Reactions Using Inertially-Confined Plasmas

Wednesday, 21 June 2017 10:00 (20 minutes)

%
% Nuclear Physics in Astrophysics 8 template for abstract
%
% Format: LaTeX2e.
%
% Rename this file to name.tex, where 'name' is the family name
% of the first author, and edit it to produce your abstract.
%
\documentstyle[11pt]{article}
%
% PAGE LAYOUT:
%
\textheight=9.9in
\textwidth=6.3in
\voffset -0.85in
\hoffset -0.35in
\topmargin 0.305in
\oddsidemargin +0.35in
\evensidemargin -0.35in

%\renewcommand{\rmdefault}{ptm} % to use Times font

\long\def\TITLE#1{{\Large{\bf#1}}}\long\def\AUTHORS#1{ #1\\[3mm]}
\long\def\AFFILIATION#1#2{[1] #2\\}
\begin{document}
{\small \it Nuclear Physics in Astrophysics 8, NPA8: 18-23 June 2017, Catania, Italy}

\vspace{12pt}

\thispagestyle{empty}

\begin{center}
%%%
%%% Title goes here.
%%%
\TITLE{Measurements Of Stellar And Big-Bang Nucleosynthesis Reactions Using Inertially-Confined Plasmas }\\[3mm]
%%%
%%% Authors and affiliations are next. The presenter should be
%%% underlined as shown below.
%%%
\AUTHORS{A.B. Zylstra[1], H.W. Herrmann[1], M. Gatu Johnson[2], Y.H. Kim[1], J.A. Frenje[2], G. Hale[1], C.K. Li[2], M. Rubery[3], M. Paris[1], A. Bacher[4], C.R. Brune[5], D.T. Casey[7], C. Forrest[6], V. Yu. Glebov[6], R. Janezic[6], D. McNabb[7], A. Nikroo[7], J. Pino[7], T.C. Sangster[6], D.B. Sayre[7], F.H. S{\'e}guin[2], H. Sio[2], C. Stoeckl[6], R.D. Petrasso[2] }

%%%
{\small \it
\AFFILIATION{1}{Los Alamos National Laboratory, Los Alamos, NM 87544 USA}
\AFFILIATION{2}{Massachusetts Institute of Technology, Cambridge, MA 02139, USA}
\AFFILIATION{3}{Plasma Physics Dept, AWE plc, Reading RG7 7PR, UK}
\AFFILIATION{4}{Indiana University, Bloomington, IN 47405, USA}
\AFFILIATION{5}{Ohio University, Athens, OH 45701, USA}
\AFFILIATION{6}{Laboratory for Laser Energetics, University of Rochester, Rochester, NY 14623, USA}
\AFFILIATION{7}{Lawrence Livermore National Laboratory, Livermore, CA 94550, USA}
}
%%%
\vspace{12pt} % Do not modify

% Enter contact e-mail address here.

\centerline{Contact email: {\it zylstra@lanl.gov}}

\vspace{18pt} % Do not modify

\end{center}
%%%
%%% Abstract proper starts here.
%%%
The ^{3}He+^{3}He, T+^{3}He, and p+D reactions directly relevant to either Stellar or Big-Bang Nucleosynthesis (BBN) have been studied at the OMEGA laser facility using inertially-confined plasmas. These high-temperature plasmas are created using shock-driven 'exploding pusher' implosions. The advantage of using these plasmas is that they better mimic astrophysical systems than cold-target accelerator experiments. A new measured S-factor for the T(^{3}He,γ)^{6}Li reaction rules out an anomalously-high ^{6}Li production during the Big Bang as an explanation to the high observed values in metal poor first generation stars. Our value is also inconsistent with values used in previous BBN calculations [1]. In a second experiment, proton spectra from the ^{3}He+^{3}He and T+^{3}He reactions are used to constrain nuclear R-matrix modeling. The spectral shapes disagree with R-matrix calculations using coefficients derived from fits to T+T data at higher or lower center-of-mass energy. Finally, recent experiments have probed the p+D reaction for the first time in a plasma; this reaction is relevant to energy production in protostars, brown dwarfs, and at higher CM energies, to BBN. The first plasma data is consistent with previous accelerator experiments at $E_{cm} \sim 16$ keV, work is ongoing to further reduce our experimental uncertainties.

\bigskip
{\small

\noindent [1] A.B. Zylstra et al., Phys. Rev. Lett. 117, 035002 (2016).

}
%%%
%%% End of abstract.
%%%
\end{document}

Primary author: Dr ZYLSTRA, Alex (Los Alamos National Laboratory)

Presenter: Dr ZYLSTRA, Alex (Los Alamos National Laboratory)

Session Classification: Nuclear astrophysics with lasers

Track Classification: Nuclear astrophysics with lasers