



EMC Simulation Studies: update on geometry effects

SuperB Collaboration Workshop

DGWG parallel session

LNF 2/12/2009

C. Cecchi - S. Germani* Università di Perugial & INFN







- Update on Fwd PID thickness effects
- Clustering Algorithm
 - Cluster Size in Barrel and Fwd
 - Tuning
- Conclusions and Outlook







LNF - 2/12/2009







- Calorimeter with back aligned endcap
- Fwd PID simulated as pasiive material:
 - Variable thickness quartz layer
- Two option for PID positions:
 - Close to the Fwd Endcap surface
 - Far form the Fwd Endcap surface







1 E_{meas}/E_{beam}



Adding the clustering algorithm the • effect of the Forward PID material it is not negligible

0.7

0.8

0.9

Larger impact on low energy • resolution

Fraction of Measured Energy

25 mm

60 mm

³⁰⁰ no longer describes

0.6

Low energy tails

0.5

700

600

500

200

100

8.4

- Fwd PID material distance from • EMC has also an impact
- Need to find a better description for • long tails



- No Clustering:
 - E_{crystal} > 1 MeV
 - Same as slac meetiong
- Clustering :
 - Clustering algorithm as (supposed to be) in BaBar:
 - 1. Start from maximum energtyCrystal
 - 2. Look for crystal arount ME Xtal
 - 3. Sum crystal energy if **E > digi** threshold (0.2 - 0.5 MeV)
 - If a Crystal around the ME one has E > seed threshold (2-3 MeV) look around it too
 - Adapted for LSO
- 5 x 5 3 x 3 Matrix
 - Take maximum energy crystal and a matrix of crystal arount it













•Measured energy in crystals around the maximum energy one (ϕ - θ plane) •Plot centered on the maximum energy crystal



Barrel

LNF - 2/12/2009





•Measured energy in crystals around the maximum energy one (φ-θ plane) •Plot centered on the maximum energy crystal



LNF - 2/12/2009





•Measured energy in crystals around the maximum energy one (φ-θ plane) for crystal above the seed threshold •Plot centered on the maximum energy crystal



LNF - 2/12/2009





•Measured energy in crystals around the amximum energy one (φ-θ plane) for crystal above the seed threshold •Plot centered on the maximum energy crystal

Seed Thr = 2 MeV



Seed Thr = 1 MeV



LNF - 2/12/2009







Crystal distance from Maximum - Projection



Projection (φ) of crystal distance from maximum energy crystal Distribution centered on the maximum energy crystal



The Fwd Endcap has a wider distribution in terms of number of crystals but in term of Moliere Radii thw width is the same for Barrel and Encap



- Energy resolution vs seed
 threshold
- Energy resolution vs distance from seed crystal
 - Include crystals above digi and seed thresholds with $\delta X \le 1$
 - Include crystals above digi and seed thresholds with $\delta X \le 2$

























•Including crystals with $\delta X \le 2$ gives non negligible improvement at loe energy for the Fwd Endcap •Need to check what happens with background





- Adding the clustering algorithm:
 - The PID material seems to have a non negligible effect on the energy resolution
 - Distance of fwd PID material from EMC have additional effect
- Clustering algorithm
 - Looking at 2 ring around the seed crystal seems to have a non negligible improvement for the Fwd Endcap
- Realistic background and signal is necessary for clustering algorithm optimization and poerformance evaluation
- Next steps:
 - Better description for low energy tails
 - Algorithm performance for π^0
 - Adding Fake background (?)
 - Adding other particles in the same event(?)
 - Other algorithm(?)