



The $\gamma\gamma$ Physics Program at **BESIII**

October 27, 2016 | Christoph Florian Redmer
for the BESIII Collaboration

KLOE-2 Workshop on e^+e^- collision physics at 1GeV

$$a_\mu = \frac{g_\mu - 2}{2} = \frac{\alpha}{2\pi} + \dots = 0.001161\dots$$

$$a_\mu^{\text{theo}} = a_\mu^{\text{QED}} + a_\mu^{\text{weak}} + a_\mu^{\text{hadr}}$$


Contribution	Result in 10^{-10} units		
QED(leptons)	11658471.885	± 0.004	Kinoshita et al. (2012)
Weak	15.4	± 0.2	Czarnecki et al. (2003)
HVP(leading order)	692.3	± 4.2	Davier et al. (2011)
HVP(higher order)	-9.84	± 0.07	Hagiwara et al. (2009)
HLBL	11.6	± 4.0	Jegerlehner, Nyffler (2009)
Total	11659181.3	± 5.8	
Experiment	11659208.9	± 6.3	


Prediction completely limited by hadronic contributions !

Challenge:

Perturbative methods cannot be applied in the relevant energy regime

- Exchange of two photons in e^+e^- collisions
- Pseudoscalar, axial, and tensor states accessible
- $M_x \ll \sqrt{s}$
- $\sigma \propto \alpha^2 \ln^2 E$
- $\sigma \propto F^2(Q_1^2, Q_2^2)$, with $Q_i^2 = -q_i^2$
- Forward peaked kinematic
 - Experimentally challenging
 - Special tagging detectors recommended






- Main Drift Chamber (MDC)
 - $\sigma(p)/p = 0.5\%$
 - $\sigma_{dE/dx} = 6.0\%$
- Time-of-flight system (TOF)
 - $\sigma(t) = 90\text{ps}$ (barrel)
 - $\sigma(t) = 110\text{ps}$ (endcap)
- EMC
 - 6240 CsI(Tl) crystals
 - $\sigma(E)/E = 2.5\%$
 - $\sigma_{Z,\phi}(E) = 0.5 - 0.7 \text{ cm}$

- Operated at BEPCII collider

- $2.0 \leq \sqrt{s} [\text{GeV}] \leq 4.6$
- Design luminosity achieved
 - $\mathcal{L} = 1.0 \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$ at $\psi(3770)$

- Large data sets for

- Charmonium Spectroscopy
- Charm Physics
- Light hadrons
- τ and R-Scan



- Muon Chambers



- 8 – 9 layers of RPC
- $p > 400 \text{ MeV}/c$
- $\delta R\Phi = 1.4 \sim 1.7 \text{ cm}$

- Superconducting Magnet

- 1 T magnetic field


Single-Tag Measurements

- Reconstruct
 - only one scattered lepton
 - Produced system
- Unmeasured lepton from momentum conservation
 - Require scattering angle to be small
 - Small virtuality
- $F(q_1^2, q_2^2) \rightarrow F(q_1^2, 0) \rightarrow F(q^2)$




Example: π^0, η
 BESIII Monte Carlo,
 $L_{int} : 2.93 \text{ fb}^{-1} @ \Psi(3770)$
 Tagged Lepton: e^+


zoom




$$e^+e^- \rightarrow e^+e^- \pi^0$$



$$e^+e^- \rightarrow e^+e^- \eta$$



$$e^+e^- \rightarrow e^+e^- \eta'$$







- Recent results from B-factories cover only large Q^2 ($5 < Q^2 [\text{GeV}^2] < 40$)
- Discrepancy for π^0 between BaBar and Belle
- Data scarce at lowest Q^2
- Region of relevance for $(g-2)_\mu$

CELLO: Z.Phys.C49 (1991) 401
 CLEO: Phys.Rev.D57 (1998) 33
 BaBar: Phys.Rev.D80 (2009) 052002
 Phys.Rev.D84 (2011) 052001
 Belle: Phys.Rev.D86 (2012) 092007

2D integral representation for pion-pole contribution by Knecht, Nyffeler (2002):


$$a_{\mu}^{\text{HLbL};\pi^0} = \int_0^{\infty} dQ_1 \int_0^{\infty} dQ_2 \sum_i w_i(Q_1, Q_2) f_i(Q_1, Q_2)$$



- Universal weight functions w_i
- Form factor dependence f_i


Relevant momentum regions:

0.25 – 1.25 GeV

BESIII Monte Carlo, $\Psi(3770)$ $L_{\text{int}} : 2.93 \text{ fb}^{-1}$, Tagged Lepton: e^+ 



- Exactly one lepton
- Two to four photons
- $\cos\theta_{\text{untagged}} < -0.99$
- $\cos\theta_{\text{Helicity}} < 0.8$

- Angle between γ in π^0 rest frame and π^0 in lab
- Flat for signal
- Peaked for background
- Reject events with $\cos(\theta_H) > 0.8$


BESIII Monte Carlo, $\Psi(3770)$ $L_{\text{int}} : 2.93 \text{ fb}^{-1}$, Tagged Lepton: e^+ 

- Exactly one lepton
- Two to four photons
- $\cos\theta_{\text{untagged}} < -0.99$
- $\cos\theta_{\text{Helicity}} < 0.8$
- Reject hadronic background

- ISR results in wrong Q^2
- Useful observable:
$$R_y = \frac{\sqrt{s} - E_{e^\pm \pi^0 \eta}^{\text{CMS}} - p_{e^\pm \pi^0 \eta}^{\text{CMS}}}{\sqrt{s}}$$
- If ISR,
$$R_y = \frac{2 E_\gamma}{\sqrt{s}}$$
- Reject events with $R_y > 0.05$

BESIII Monte Carlo, $\Psi(3770)$ $L_{\text{int}} : 2.93 \text{ fb}^{-1}$, Tagged Lepton: e^+ 

- Analysis useful for π^0 and η
- Monte Carlo description of background incomplete
- Bkg subtr. by counting π^0 yield per Q^2 bin
- $|F(Q^2)|^2$ extracted by division by WZW-MC



- Exactly one lepton
- Two to four photons
- $\cos\theta_{\text{untagged}} < -0.99$
- $\cos\theta_{\text{Helicity}} < 0.8$
- Reject hadronic background
- Bkg subtr. by counting π^0 yield per Q^2 bin
- $|F(Q^2)|^2$ extracted by division by WZW-MC
- Full BESIII $\Psi(3770)$ data set analyzed
- Competitive accuracy up to 3.1 GeV^2
- Unprecedented accuracy below $Q^2 = 1.5 \text{ GeV}^2$

- Only statistical errors shown
- Systematic uncertainty dominated by background subtraction


- Current accuracy of a_μ : $\sim 6.3 \times 10^{-10}$
- Contribution of π^0 : $\sim 7 \times 10^{-10}$ Knecht, Nyffeler
Phys.Rev.D65 (2002) 073034
- Expected accuracy of new experiments at FNAL and J-PARC: $\sim 1.6 \times 10^{-10}$

- Contributions of η and η' relevant!


$$\eta \sim 1.5 \times 10^{-10}$$

$$\eta' \sim 1.5 \times 10^{-10}$$

Knecht, Nyffeler
Phys.Rev.D65 (2002) 073034




BESIII Simulation: $2.9 fb^{-1}$ @ 3.773 GeV




- $\eta \rightarrow \pi^+ \pi^- \pi^0$
- $\eta' \rightarrow \pi^+ \pi^- \eta$
- Any combination of
 - one positron
 - two charged pions
 - two photons
- $\cos\theta_{\text{untagged}} > 0.99$
- reject hadronic background
- Mass window cuts on $\gamma\gamma$ invariant mass
- Kinematic fit
- Relatively low background conditions

$$F_{\eta, \gamma, \gamma^*}(Q^2)$$



$$F_{\eta', \gamma, \gamma^*}(Q^2)$$



- Full BESIII $\Psi(3770)$ data set analyzed
- statistics compatible to previous measurements
 - only one decay channel of η and η' analyzed at BESIII
 - More ($\times 3.5$) data available
- Systematic studies to be done


JGU Space-like $\pi^+\pi^-$ Transition Form Factor

- Additional Motivations:

- Resonance parameters
- Pion polarizabilities, pion structure
- Essential for dispersive frameworks
- Rescattering effects in low mass region

Collangelo, Hoferichter, Procura, Stoffer
JHEP 1409,091; JHEP1509,074

- Until recently only untagged measurements:



MarkII, Phys. Rev. D42 (1990) 5

CELLO, Z. Phys. C56 (1992) 381

Belle, Phys. Rev D75 (2007) 051101


- First single-tagged result on $\pi^0\pi^0$ by Belle

Phys. Rev. D93 (2016) 032003

JG|U Space-like $\pi^+\pi^-$ Transition Form Factor

- Additional Motivations:
 - Resonance parameters
 - Pion polarizabilities, pion structure
 - Essential for dispersive frameworks
 - Rescattering effects in low mass region
- Until recently only untagged measurements:

Collangelo, Hoferichter, Procura, Stoffer
JHEP 1409,091; JHEP1509,074



Calculations by
Assmussen, Masjuan, and Vanderhaeghen:

Untagged

Single-Tag ($Q_1^2 = 0.5 \text{ GeV}^2$)

Double-Tag ($Q_1^2 = Q_2^2 = 0.5 \text{ GeV}^2$)

- First single-tagged result on $\pi^0\pi^0$ by Belle

Phys. Rev. D93 (2016) 032003

- At BESIII: Single-Tag measurement


- Event selection analogous to single pseudoscalar analysis

- Major Backgrounds:

- $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

- Radiative Bhabha scattering coupling to ρ (s + t channel)

- MC generators missing




- $\gamma\gamma^{(*)} \rightarrow \mu^+\mu^-$

- Two-photon production of muon pairs

- Precise MC generators available from LEP era (BdkRC + Diag36ABC)

- Train ANN to suppress muon background




- Subtract muon background using MC distributions
 - To be replaced with ANN trained for muon suppression
- Subtract ρ contribution
 - Fit with Kühn-Santamaria parameterization
- Study $\pi^+\pi^-$ invariant mass in bins of Q^2 and $\cos\theta^*$
- First single-tag measurement of $\pi^+\pi^-$!
- Access to:
 - low momentum transfers $0.2 < Q^2 [\text{GeV}^2] < 2.0$
 - low invariant masses $m_{\pi^+\pi^-} < M [\text{GeV}] < 2.0$
 - full coverage of $\cos\theta^*$

- More than 10 fb^{-1} on disk at $3.77 < \sqrt{s} [\text{GeV}] < 4.6$

- Double-tag measurement possible

- Measure $F_{\gamma^* \gamma^* \pi^0}(Q_1^2, Q_2^2)$
- 1st Step: Test TFF models
 - e.g. VMD vs. LMD+V



Calculations: A. Nyffeler
hep-ph:1602.03398

- Test polarization in $\gamma\gamma$ production
 - General two-photon cross section:



$$d\sigma = F \left\{ v_{TT} \sigma_{TT} + v'_{TT} \cos(2\tilde{\phi}) (\sigma_{\parallel} - \sigma_{\perp}) + h_1 h_2 v''_{TT} \frac{1}{2} (\sigma_0 - \sigma_2) \right. \\ \left. + v_{LL} \sigma_{LL} + v_{TL} \sigma_{TL} + v_{LT} \sigma_{LT} + v'_{TL} \cos(\tilde{\phi}) \tau_{TL} + h_1 h_2 v''_{TL} \cos(\tilde{\phi}) \tau_{TL}^a \right\}.$$

- $\tilde{\phi}$: azimuthal angle between lepton planes in $\gamma^* \gamma^*$ CMS
 - Allows to disentangle form factor contributions of multi-meson and tensor states
 - Requires precise measurement of angles


Outlook: Zero Degree Detector



- Tagging of photons and electrons at small angles
 - Polar angle range: 1 – 10 mrad
- Current design: Pb-SciFi, one sided
- Upgrade: Arrays of 48 crystals (PbWO, LYSO) on both sides



In-beam tests at MAMI (Mainz)



- Two-photon physics program established at BESIII
 - Single-tag measurements of π^0 , η , and η' transition form factors
 - Unprecedented accuracy for $Q^2 < 1.5 \text{ GeV}^2$
 - Single-tag measurement of $\pi^+\pi^-$
 - First measurement at low Q^2 , low mass
 - To be extended to neutral final states
 - First double-tagged measurement $\gamma^*\gamma^* \rightarrow \pi^0$ started
 - New prospects from tagging detectors
 - Recent addition to the $\gamma\gamma$ program:
 - Untagged measurements for light hadron spectroscopy