




# Neutrino Physics and Astrophysics with Borexino





The choice of the liquid scintillation technique was dictated by the high light-yield of the scintillator (50 times more than in the Cherenkov technique), and then a good energy resolution – in Bx 500 pe/ MeV

Borexino data taking history in four lines: start up-May 2007

phase 1-2007-2010


further purification 2010-2011


since Dec. 2011- phase 2

Radio-Isotope		Concentration or Flux		Achieved	Achieved
Name	Source	Typical	Required	phase 1	Phase 2
$^{14}\text{C}$	<i>intrinsic scintillator</i>	$\sim 10^{-12} \text{ }^{14}\text{C}/^{12}\text{C}$	$\sim 10^{-18} \text{ }^{14}\text{C}/^{12}\text{C}$	$\sim 2 \times 10^{-18} \text{ }^{14}\text{C}/^{12}\text{C}$	
$^{238}\text{U}$ $^{232}\text{Th}$ equiv.	<i>dust, particulate, all materials</i>	$10^{-5} - 10^{-6} \text{ g/g}$	$< 10^{-16} \text{ g/g}$	$(5.0 \pm 0.9) 10^{-18} \text{ g/g}$ $(3.0 \pm 1.0) 10^{-18} \text{ g/g}$	$< 9.5 \cdot 10^{-20} \text{ g/g}$ $< 7.2 \cdot 10^{-19} \text{ g/g}$
$^7\text{Be}$	<i>cosmogenic</i>	$\sim 3 \times 10^{-2} \text{ Bq/t}$	$< 10^{-6} \text{ Bq/t}$	<i>not observed</i>	
$^{40}\text{K}$	<i>dust, PPO</i>	$\sim 2 \times 10^{-6} \text{ g/g (dust)}$	$< 10^{-18} \text{ g/g}$	<i>not observed</i>	
$^{210}\text{Po}$	<i>surface contamination.</i>	<i>Decaying with a half time of <math>\sim 138</math> days</i>	$< 700 \text{ cpd/100t}$	$500.-20.\text{cpd/100t}$	$\sim 20 \text{ cpd/100t}$
$^{222}\text{Rn}$	<i>emanation from materials, rock</i>	$10 \text{ Bq/l air, water}$ $100-1000 \text{ Bq/kg}$ <i>rock</i>	$< 10 \text{ cpd/100t}$	$< 1 \text{ cpd /100 t}$	
$^{39}\text{Ar}$	<i>air, cosmogenic</i>	$17 \text{ mBq/m}^3 \text{ (air)}$	$< 1 \text{ cpd/100 t}$	$<< 1 \text{ cpd/100 t}$	
$^{85}\text{Kr}$	<i>air, nuclear weapons</i>	$\sim 1 \text{ Bq/m}^3 \text{ (air)}$	$< 1 \text{ cpd/100 t}$	$30 \pm 5 \text{ cpd/100 t}$	$< 6.4 \text{ cpd/100 t}$ <i>fit consistent with 0</i>


unprecedented radio-purity, never reached by any other experiment until now

The Sun functioning is based on the pp chain:





pp



## Physics World: Top 10 Physics Breakthroughs of 2014

astr


Main problem  $^{14}\text{C}$  (end point: 156keV)  
pp- end point: 264 keV

pile up:  $^{14}\text{C}$ - $^{14}\text{C}$  and  $^{14}\text{C}$ -pp ( data driven method)


fit range: 165-590 keV

$$\Phi_{pp} = (6.6 \pm 0.7) 10^{10} \text{ cm}^{-2} \text{ s}^{-1}$$

$^7\text{Be}$



Compton-like shoulder at 660 keV  
fit range: 200-1270 keV



$$\Phi_{^7\text{Be}} = (4.43 \pm 0.22) 10^9 \text{ cm}^{-2} \text{ s}^{-1}$$




# pep/ CNO

pep signal is ten times lower than  $^7\text{Be}$

- About 3cpd/100t
- CNO: rate similar to pep

## Most important background

- cosmogenic  $^{11}\text{C}$  :
- $^{210}\text{Bi}$  :
- External background ( $\gamma$  from PMT):



TFC: incident muon

$e^+$  annihilation from  $^{11}\text{C}$  decay (prompt signal)

p capture of the thermalized neutron (delayed signal- $\sim 240 \mu\text{s}$ )

Pulse shape discrimination: in 50% of the cases the formation of ortho-positronium the  $e^+$  annihilation is delayed

Fit simultaneously

- 1) Energy spectra after  $^{11}\text{C}$  subtraction
- 2) Energy spectra of the subtracted events
- 3) Radial distribution of the events

CNO energy distribution shows a shape very similar to the  $^{210}\text{Bi}$  spectrum ( $\sim 15\text{-}20 \text{ cpd}/100 \text{ t}$ ).

$$\Phi_{\text{pep}} = (1.63 \pm 0.35) 10^8 \text{ cm}^{-2} \text{ s}^{-1}$$

$$\Phi_{\text{CNO}} < 7.7 \times 10^8 \text{ cm}^{-2} \text{ s}^{-1}.$$

$^8\text{B}$ 


not monochromatic, then no fit on the energy spectrum, but on the radial distribution- main background  $^{208}\text{Tl}$ , subtracted via  $^{212}\text{Bi}$ - $^{212}\text{Po}$  coincidence  
-threshold down to 3.2 MeV  $\nu$  energy

$$\Phi_{8\text{B}} = (2.4 \pm 0.4 \pm 0.1) 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

good agreement with SNO and SK

## seasonal modulation

eccentricity of the Earth orbit- 6.7%



@ Borexino phase 2 until December 2015-

1456 astronomical days of data

@ counting in the energy range: 215-715 keV ( $^7\text{Be}$  region)

@ very efficient  $\alpha/\beta$  discrimination: 99.98% - scintillator p. sh.- learning machine method


@ high detector stability- background stability

### Modulation analysis


- *sinusoidal fit*
- *Lomb-Scargle method*- an extension of the Fourier Transform Approach- can treat data sets not evenly distributed in time



## Sinusoidal fit




30 day bins- 4 years data cumulated




20 days bins-

**Null hypothesis rejected at  $3.91\sigma$  (99.99% C.L.)**  
**modulation amplitude  $(6.8 \pm 1.9)\%$**   
**best-t period is  $T = 369.0 \pm 11$  days.**  
**phase =  $-12 \pm 28$  days** ( to be compared with January 3th).

## Lomb-Scargle analysis



L-S Normalized Spectrum Power Density ( $f=1y^{-1}$ )  $\approx 8$



**From Lomb Scargle analysis:**  
**Period 1 year**  
**Absence of modulation rejected**  
**at 99.99% C.L ( $>3.5 \sigma$ ).**

Solar neutrino flux	GS98 (cm <sup>-2</sup> s <sup>-1</sup> ) High metallicity Z/X=0.0229***	AGS09 (cm <sup>-2</sup> s <sup>-1</sup> ) Low metallicity Z/X=0.0178***	Experimental results (Borexino)	Global fit including solar, reactor, accel. data (cm <sup>-2</sup> s <sup>-1</sup> )**
pp	5.99 (1±0.006) x 10 <sup>10</sup>	6.04 (1±0.005) x 10 <sup>10</sup>	6.6 (1±0.106) x 10 <sup>10</sup>	5.97 <sup>+0.037</sup> <sub>-0.033</sub> x 10 <sup>10</sup>
<sup>7</sup> Be	4.80 (1±0.06) x 10 <sup>9</sup> [4.93] <sup>^*</sup> x 10 <sup>9</sup>	4.38 (1±0.06) x 10 <sup>9</sup> [4.50] <sup>^*</sup> x 10 <sup>9</sup>	4.94±0.22 x 10 <sup>9</sup> *	4.80 <sup>+0.24</sup> <sub>-0.22</sub> x 10 <sup>9</sup>
pep	1.44 (1±0.01) x 10 <sup>8</sup>	1.47 (1±0.009) x 10 <sup>8</sup>	1.63±0.35 x 10 <sup>8</sup>	1.448±0.013 x 10 <sup>8</sup>
<sup>13</sup> N <sup>15</sup> O <sup>17</sup> F	2.78 (1±0.15) x 10 <sup>8</sup> 2.05 (1±0.17) x 10 <sup>8</sup> 5.30 (1±0.20) x 10 <sup>8</sup>	2.05 (1±0.14) x 10 <sup>8</sup> 1.44 (1±0.16) x 10 <sup>8</sup> 3.26 (1±0.18) x 10 <sup>8</sup>	 <7.7 x 10 <sup>8</sup> total CNO	≤ 13.7 x 10 <sup>8</sup> ≤ 2.8 x 10 <sup>8</sup> ≤ 8.5 x 10 <sup>7</sup>
<sup>8</sup> B	5.32 (1±0.12) x 10 <sup>6</sup> [5.46] <sup>^*</sup> x 10 <sup>6</sup>	4.37 (1±0.12) x 10 <sup>6</sup> [4.50] <sup>^*</sup> x 10 <sup>6</sup>	5.2±0.3 x 10 <sup>6</sup>	5.16 <sup>+0.13</sup> <sub>-0.09</sub> <sup>+0.30</sup> <sub>-0.26</sub> x 10 <sup>6</sup>

- This is the <sup>7</sup>Be flux for both the <sup>7</sup>Be lines, extrapolated from the higher energy line flux measured by Borexino (96% of the total) in phase 1; the preliminary results of the phase 2 are in agreement with this quoted value and with a total error reduced to 3-3.5%

\*\* from J. Bergstroem et al., arXiv: 1601.00972v1 [hep-ph] 5 jan 2016


\*\*\* from A. Serenelli, F. Villante et al. –paper in preparation- They review the uncertainties of the opacities and up to date the cross section factors: S<sub>11</sub> (pp), S<sub>34</sub> (<sup>3</sup>He+<sup>4</sup>He), S<sub>17</sub> (<sup>7</sup>Be+p), S<sub>114</sub> (<sup>14</sup>N+p)-in these calculations S<sub>34</sub> is assumed by de Boer et al. 2014

<sup>^\*</sup> These different numbers are obtained assuming for S<sub>34</sub> what Iliadis et al. (very recent) and Adelberger et al. 2011 have quoted (F. Villante-private communication)----

heliosismology


## CNO cycle in the Sun only 1%

Reactions of the CNO Cycle



hypothesized as the primary cycle  
in the massive stars and then for  
hydrogen burning in the Universe


astr




**strategy:** need of an independent constraint on  $^{210}\text{Bi}$  rate (remove degeneracy with CNO spectrum)

from  $^{210}\text{Po}$ : two components:

- embedded on the lines (138.376 days half time)
- $^{210}\text{Pb} \rightarrow ^{210}\text{Bi} (\beta) \rightarrow ^{210}\text{Po} (\alpha)$
- easily identified via pulse shape discrimination




## Borexino reached fundamental achievements in Sun Physics




Borexino has demonstrated experimentally that the Sun shines via the proton-proton nuclear reaction chain in the solar interior:  $pp^{**}$ ,  $pep$ ,  ${}^7\text{Be}$ ,  ${}^8\text{B}^*$

\* already measured also by SNO and SuperK

\*\* the radiochemical exp. measured the whole integrated solar flux above 233 keV



The agreement between the solar luminosity in neutrinos, as measure by Borexino, and in photons demonstrates the Sun stability on  $\sim 10^5$  years time scale



Borexino has the chance to give the first experimental evidence of the CNO cycle and to provide a contribution to the solution of the metallicity puzzle in the SSM

# NEUTRINO PHYSICS

$\nu$

## Neutrino oscillation

In two neutrino scenario

IN VACUUM

$$P(\nu_e \rightarrow \nu_\mu) = \sin^2 2\theta \cdot \sin^2 \left( \Delta m^2 \frac{L}{4E} \right)$$

IN MATTER

$$P(\nu_e \rightarrow \nu_\mu) = \sin^2 2\theta_M \cdot \sin^2 \left( \Delta m_M^2 \frac{L}{4E} \right)$$

If  $X = \cos 2\theta$  - maximum mixing


$$\sin^2 2\theta_M = \frac{\sin^2 2\theta}{\sin^2 2\theta + (\cos 2\theta - X)^2}$$

$$L_M = \frac{4\pi E}{\Delta m_M^2} = \frac{L_V}{\sqrt{\sin^2 2\theta + (\cos 2\theta - X)^2}}$$


$4\pi E / \Delta m^2$

$$X = \frac{2\sqrt{2}G_F n_e E}{\Delta m^2}$$

The best way to understand the Borexino contribution to the study of the neutrino oscillation is to compare **the electron-neutrino survival probability** before and after Borexino.



before Borexino



after Borexino

Borexino succeeded to observe for the first time the oscillation in vacuum measuring the  $pp$  neutrino survival probability ( $0.64 \pm 0.12$ ) and to determine the ratio between the survival probabilities in vacuum and in matter, using  $pp$  and  ${}^8\text{B}$ , this last measured by Borexino above 3 MeV. This ratio turns out to be  $=1.75 \pm 0.86$ .




## Study of the day/night asymmetry


- regeneration of the electron neutrinos crossing the Earth
- D/N effect is a consequence of MSW
- not expected for  ${}^7\text{Be}$  in the LMA-MSW model
- large effect expected in the “LOW” solution (excluded by solar exp + Kamland)
- no contradiction with the recent SK results

$$A_{DN} = \frac{N - D}{(N + D)/2} = 0.001 \pm 0.012 (stat) \pm 0.007 (sys)$$

Solar data alone select the LMA-MSW if one includes the Borexino D/N result- $8.5 \sigma$  C.L. (no use of CPT invariance)



## Transition region between the vacuum and matter regime



Recent SuperK paper (June 2016) does not change substantially the situation—measurements down to  $\sim 4$  MeV

Subleading effects (beyond the S.M.):

1-very light sterile mixes very weakly with active neutrinos, which have the smallest mass splitting (solar) (Smirnov & De Holland, Phys. Rev. 2011).

The green curve is calculated for


$$\Delta m_{21}^2 = 7.5 \cdot 10^{-5} eV^2 \quad \Delta m_{01}^2 = 1.2 \cdot 10^{-5} eV^2 \quad \sin^2 \alpha = 0.005 \quad \sin^2 \theta_{12} = 0.31$$

2. NSI- The NSI can be described at low energy by effective four fermion interactions

$\ell_{NSI} = -2\sqrt{2}G_F \epsilon_{\alpha\beta}^{e,u,d} (\bar{\nu}_\alpha \gamma^\mu P_L \nu_\beta) (\bar{f} \gamma_\mu P_C f')$  where  $G_F$  is the Fermi constant,  $\alpha$  and  $\beta$  are the neutrino flavors,  $f$  and  $f'$  the electron or the light quarks,  $L$  and  $R$  indicate the projection of the operator  $P$  (two chiralities), and finally  $\epsilon$  parameterizes the strength of the NS interaction. The curves are calculated for  $\epsilon_D^u = -0.22$ ,  $\epsilon_N^u = -0.30$ ;  $\epsilon_D^d = -0.12$ ,  $\epsilon_N^d = -0.16$  ( $\epsilon_D$  and  $\epsilon_N$  are linear combination of  $\epsilon$ )

A test of these subleading effects can be successful if a measurement is carried out in the range 2.-5. MeV. **Borexino phase 2 is measuring an experimental point for  ${}^8\text{B}$  between 3.-5. MeV.**

The study of a possible NSI can be done more easily analyzing the energy spectrum of the recoiled electron from the  $\nu$ -e scattering, in particular for the  ${}^7\text{Be}$  neutrinos, which is monoenergetic and then does not need of convolution with the incident neutrino spectrum, as for instance is the case of  ${}^8\text{B}$ .




$$\frac{d\bar{\sigma}(T)}{dT} = \frac{2G_F^2 m_e}{\pi} \left[ \bar{g}_{eL}^2 + \bar{g}_{eR}^2 \left( 1 - \frac{T}{E_{\nu_e}} \right) - \bar{g}_{eL} \bar{g}_{eR} \frac{m_e T}{E_{\nu_e}} \right]$$


$$T_{\text{max}} = 0.665 \text{ MeV for } {}^7\text{Be}$$

$$\text{----- } \epsilon_{eL}; \epsilon_{eR} = 0.01, 0.2; = -0.02, 0.05$$

*Bx phase 1*



*Bx phase 2 (foreseen)*



Analysis of the phase 1 data-the contaminants  ${}^{85}\text{Kr}$  and  ${}^{210}\text{Bi}$  influence the analysis as far as the uncertainty in the  ${}^7\text{Be}$  flux

Phase 2:

Kr from  $10 \pm 3$  to  $\sim 0$ . cpd/100 t

Bi from  $\sim 40$  to  $\sim 15$  cpd/100t

${}^7\text{Be}$  flux uncert. from 5% to  $< 3.5\%$


## Search for sterile $\nu$ in Borexino: SOX

$\nu$

### SOX: Short distance neutrino Oscillations with BoreXino

Artificial external neutrino source is allocated in a tunnel present under the Borexino detector, at 8.25 m from the detector center. The source will be a  $^{144}\text{Ce}$ - $^{144}\text{Pr}$  activated at  $\sim 5$  PBq; it emits antineutrinos with a continuous energy distribution up to 3 MeV.

Borexino can study the very short distance neutrino oscillations in the standard *disappearance* technique, but it is possible to observe directly the oscillation waves in the hypothesis of very short baseline oscillations. E/L allows to explore the region around  $\Delta m^2 \sim 1\text{eV}^2$  and the typical oscillation length of a few m (detector diameter  $\sim 6$  m.)



see talk of Lea Di Noto

### WHY:

## LNSD and MiniBooNE :both observe excess of  $\bar{\nu}_e$  from  $\bar{\nu}_\mu$  beam and MiniBooNE also of  $\nu_e$  from  $\nu_\mu$  beam  $\Delta m^2 \sim 1\text{eV}^2$  --  $3.8\sigma$ : a fourth  $\nu$  needed.

E/L agrees with SOX

# claimed reactor problem :  $\sim 3\%$  deficit --  $2.5\sigma$   
Gallex and Sage source anomaly -Deficit of the detected  $\nu_e$  -  $R = 0.76 \pm 0.09$  -  $2.8\sigma$

# Geoneutrinos

geo

## Detection method

$$\bar{\nu} + p \rightarrow n + e^+ \quad E_{\nu} > 1.8 \text{ MeV}$$

- “prompt signal”  
e+: energy loss + annihilation  
(2  $\gamma$ , 511 KeV each)
- “delayed signal”  
n capture after thermalization, 2.2  $\gamma$
- Space and time coincidence

Fit on the selected antineutrino spectrum:

black points = data

----- = best fit U+Th with fixed  
mass chondritic ratio (Th/U=3.9):



$$N_{geo}^{events} = 23.7_{-5.7}^{+6.5} (stat.)_{-0.6}^{+0.9} (syst.)$$

$$N_{reactor}^{events} = 52.7_{-7.7}^{+8.5} (stat.)_{-0.9}^{+0.7} (syst.)$$

blue area=U free parameter

light blue area= U+Th free parameters

----- = Reactor neutrinos



$$S_{geo} = 43.5_{-10.4}^{+11.8} (stat.)_{-2.4}^{+2.7} (syst) \text{ TNU}$$

TNU= Terrestrial neutrino Unit= 1 event/year/10<sup>32</sup>  
protons, 100% efficiency

**Borexino reached an evidence at  $5.9 \sigma$  C.L. and a null hypothesis with a probability of  $\sim 3.6 \cdot 10^{-9}$**

geo

Some hints already reached ( also with the Kamland data):

1. The evaluation of the geo-neutrino signal from the mantle is obtained by subtracting the crust contribution from the measured signal. At the Gran Sasso site the crust signal has been evaluated to be  $S_{\text{crust}} = 23.4 \pm 2.8 \text{ TNU}$  and then  
 $S_{\text{mantle}} = 20.9 + 15.1 - 10.3 \text{ TNU}$
2. The radiogenic heat represents an important part of the total Earth's energy budget. From the Borexino data, If the chondritic mass ratio and the ratio  $m(\text{K})/m(\text{U}) = 10^4$  are assumed, the **radiogenic heat is  $= 33^{+28}_{-20} \text{ TW}$** . The large range is due to the various assumptions on the radioactive nuclide distribution in the mantle: either homogeneously diffused or accumulated close the border mantle-core.. The radiogenic contribution has to be compared with the evaluated **total Earth's heat, which ranges from 31 to 47 TW**.
3. For the first time it is possible to observe two well **separated peaks** in the geo-neutrinos energy range, **produced by  $^{232}\text{Th} + ^{238}\text{U}$  and by  $^{238}\text{U}$** .




In the mean time Borexino reached the **following limits**:

**neutrino effective magnetic moment.** Study of the shape of the electron recoil, especially at low energy where it is enhance at  $\sim 1/E_e$   $\mu_{eff} \leq 5.4 \cdot 10^{-10} B_\mu$  (90% C. L.)


**Test of Pauli exclusion principle.** New limits on non paullian transitions of nucleons from  $1P_{3/2}$  shell to  $1S_{1/2}$  shell in  $^{12}\text{C}$  with the emission of  $\gamma$ , n, p and  $\beta^\pm$  particles:  $\tau(^{12}\text{C} \rightarrow ^{12}\text{C}^{\text{NP}} + \gamma) \geq 5.0 \cdot 10^{31} \text{y}$ ,  $\tau(^{12}\text{C} \rightarrow ^{11}\text{B}^{\text{NP}} + \text{p}) \geq 8.9 \cdot 10^{29} \text{y}$ ,  $\tau(^{12}\text{C} \rightarrow ^{11}\text{C}^{\text{NP}} + \text{n}) \geq 3.4 \cdot 10^{30} \text{y}$ ,  $\tau(^{12}\text{C} \rightarrow ^{12}\text{N}^{\text{NP}} + \text{e} + \nu) \geq 3.1 \cdot 10^{30} \text{y}$ ,  $\tau(^{12}\text{C} \rightarrow ^{12}\text{B}^{\text{NP}} + \text{e} + \nu) \geq 2.1 \cdot 10^{30} \text{y}$ , **BEST LIMITS**

**Stability of the electron:**  $e^- \rightarrow \gamma + \nu^-$  search for 256 keV  $\gamma$  peak – consistent with zero-  $\tau \geq 6.4 \cdot 10^{28} \text{years}$  (90% C.L.) **BEST LIMIT**

**Low energy neutrino and antineutrino signals correlated with gamma-ray bursts.** Dec. 2007-Nov. 1015 (2302 live days)- 2350 GRBs from the IceCube database-  $\Delta t$  (sign)  $\pm 1000$  s for the elastic scattering on electron and 5000 s for antinu events range 1.-15. MeV-separate analysis for electron  $\bar{\nu}$  over 1.8 MeV (inv. beta decay events-well tagged) and the elastic scattering at any energy ( neutrinos and antineutrinos)



electron antineutrinos IBD



(anti) neutrino elastic scattering on electron

## Conclusions:

1. Borexino is still a **unique** detector to study the low energy neutrinos
2. Its results found the **direct experimental evidence** that the nuclear reactions supporting the Sun shining belong to the **pp cycle**
3. For the first time the **oscillation in vacuum** has been observed experimentally and the related  $\nu_e$  **survival probability** has been measured
4. The Borexino data reached ( **5.9  $\sigma$**  ) evidence of the existence of **geo-neutrinos**
5. As byproduct several best **limits** have been obtained
6. Borexino will take data until 2020-2022 doing an effort to measure the **CNO flux** (very challenging) and to check the possible existence of VSB oscillation and of a sterile neutrino



## Borexino Collaboration



UNIVERSITÀ  
DEGLI STUDI  
DI MILANO



PRINCETON  
UNIVERSITY



UNIVERSITÀ DEGLI STUDI  
DI GENOVA



NATIONAL RESEARCH CENTER  
"KURCHATOV INSTITUTE"



Technische Universität  
München



JOHANNES GUTENBERG  
UNIVERSITÄT MAINZ



University of  
Houston



JAGIELLONIAN  
UNIVERSITY  
IN KRAKÓW



JÜLICH  
FORSCHUNGSZENTRUM



Universität  
Hamburg

Virginia  
Tech



SKOBELOTSYN INSTITUTE OF NUCLEAR PHYSICS  
LOMONOSOV MOSCOW STATE UNIVERSITY



Joint Institute for  
Nuclear Research



GRAN SASSO  
SCIENCE INSTITUTE  
CENTRO PER LO STUDIO  
DELLA FISICA NUCLEARE



TECHNISCHE  
UNIVERSITÄT  
DRESDEN



# Thank you for your attention