




UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386



Complex Langevin simulations of a finite density matrix model for QCD

Savvas Zafeiropoulos


Universität Heidelberg

14.12.2017

SM & FT 2017

In collaboration with J. Bloch (Regensburg U.), J. Glesaaen (Swansea U.), J. Verbaarschot (Stony Brook U.)

Phase diagram from a theorist's viewpoint



Stochastic quantization as an alternative

- consider the trivial "QFT" given by the partition function
- $\mathcal{Z} = \int e^{-S(x)} dx$
- in the real Langevin formulation
- $x(t + \delta t) = x(t) - \partial_x S(x(t)) \delta t + \delta \xi$
- stochastic variable $\delta \xi$ with zero mean and variance given by $2\delta t$
- generalization to complex actions Parisi (1983), Klauder (1983)
- $x \rightarrow z = x + iy$
- $z(t + \delta t) = z(t) - \partial_z S(z(t)) \delta t + \delta \xi$
- one can study gauge theories with complex actions Aarts, James, Seiler, Sexty, Stamatescu, ...

Stochastic quantization as an alternative

- consider the trivial "QFT" given by the partition function
- $\mathcal{Z} = \int e^{-S(x)} dx$
- in the real Langevin formulation
- $x(t + \delta t) = x(t) - \partial_x S(x(t)) \delta t + \delta \xi$
- stochastic variable $\delta \xi$ with zero mean and variance given by $2\delta t$
- generalization to complex actions Parisi (1983), Klauder (1983)
- $x \rightarrow z = x + iy$
- $z(t + \delta t) = z(t) - \partial_z S(z(t)) \delta t + \delta \xi$
- one can study gauge theories with complex actions Aarts, James, Seiler, Sexty, Stamatescu, ...

Stochastic quantization as an alternative

- consider the trivial "QFT" given by the partition function
- $\mathcal{Z} = \int e^{-S(x)} dx$
- in the real Langevin formulation
- $x(t + \delta t) = x(t) - \partial_x S(x(t)) \delta t + \delta \xi$
- stochastic variable $\delta \xi$ with zero mean and variance given by $2\delta t$
- generalization to complex actions Parisi (1983), Klauder (1983)
- $x \rightarrow z = x + iy$
- $z(t + \delta t) = z(t) - \partial_z S(z(t)) \delta t + \delta \xi$
- one can study gauge theories with complex actions Aarts, James, Seiler, Sexty, Stamatescu, ...

Stochastic quantization as an alternative

- consider the trivial "QFT" given by the partition function
- $\mathcal{Z} = \int e^{-S(x)} dx$
- in the real Langevin formulation
- $x(t + \delta t) = x(t) - \partial_x S(x(t)) \delta t + \delta \xi$
- stochastic variable $\delta \xi$ with zero mean and variance given by $2\delta t$
- generalization to complex actions Parisi (1983), Klauder (1983)
- $x \rightarrow z = x + iy$
- $z(t + \delta t) = z(t) - \partial_z S(z(t)) \delta t + \delta \xi$
- one can study gauge theories with complex actions Aarts, James, Seiler, Sexty, Stamatescu, ...

Stochastic quantization as an alternative

- consider the trivial "QFT" given by the partition function
- $\mathcal{Z} = \int e^{-S(x)} dx$
- in the real Langevin formulation
- $x(t + \delta t) = x(t) - \partial_x S(x(t)) \delta t + \delta \xi$
- stochastic variable $\delta \xi$ with zero mean and variance given by $2\delta t$
- generalization to complex actions [Parisi \(1983\)](#), [Klauder \(1983\)](#)
- $x \rightarrow z = x + iy$
- $z(t + \delta t) = z(t) - \partial_z S(z(t)) \delta t + \delta \xi$
- one can study gauge theories with complex actions [Aarts, James, Seiler, Sexty, Stamatescu, ...](#)

Is this "the" solution to the sign problem?

- proof relating Langevin dynamics to the path integral quantization-no longer holds
- simulations are not guaranteed to converge to "the correct solution"
- criteria of convergence not fulfilled in practical simulations (at least for some range of the parameters)

Is this "the" solution to the sign problem?

- proof relating Langevin dynamics to the path integral quantization-no longer holds
- simulations are not guaranteed to converge to "the correct solution"
- criteria of convergence not fulfilled in practical simulations (at least for some range of the parameters)

Is this "the" solution to the sign problem?

- proof relating Langevin dynamics to the path integral quantization-no longer holds
- simulations are not guaranteed to converge to "the correct solution"
- criteria of convergence not fulfilled in practical simulations (at least for some range of the parameters)

Validity criteria

- action S and observables \mathcal{O} should be holomorphic in the complexified variables (up to singularities)
- the probability distribution of the complexified variables z along the CL trajectories should be suppressed close to the singularities of the drift term and of the observable
- sufficiently fast decay of the probability distribution in the imaginary direction

Validity criteria

- action S and observables \mathcal{O} should be holomorphic in the complexified variables (up to singularities)
- the probability distribution of the complexified variables z along the CL trajectories should be suppressed close to the singularities of the drift term and of the observable
- sufficiently fast decay of the probability distribution in the imaginary direction

Validity criteria

- action S and observables \mathcal{O} should be holomorphic in the complexified variables (up to singularities)
- the probability distribution of the complexified variables z along the CL trajectories should be suppressed close to the singularities of the drift term and of the observable
- sufficiently fast decay of the probability distribution in the imaginary direction

- focus on a much simpler theory than QCD. Random Matrix Theory
- same flavor symmetries with QCD which uniquely determine (in the ϵ -regime)
 - mass dependence of the chiral condensate $\langle \bar{\eta} \eta \rangle = \partial_m \log Z$
 - the baryon number density $\langle \eta^\dagger \eta \rangle = \partial_\mu \log Z$

- focus on a much simpler theory than QCD. Random Matrix Theory
- same flavor symmetries with QCD which uniquely determine (in the ϵ -regime)
 - mass dependence of the chiral condensate $\langle \bar{\eta} \eta \rangle = \partial_m \log Z$
 - the baryon number density $\langle \eta^\dagger \eta \rangle = \partial_\mu \log Z$

The Stephanov Model

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr} WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$

Stephanov (1996) and Halasz, Jackson, Verbaarschot(1997)

- solve via bosonization

- $\mathcal{Z}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- where σ is an $N_f \times N_f$ matrix

- $\mathcal{Z}^{N_f=1}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- $\mathcal{Z}^{N_f=1}(m, \mu) = \pi e^{-nm^2} \int_0^\infty du (u - \mu^2)^n I_0(2mn\sqrt{u}) e^{-nu}$

The Stephanov Model

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr} WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$

Stephanov (1996) and Halasz, Jackson, Verbaarschot(1997)

- solve via bosonization

- $\mathcal{Z}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- where σ is an $N_f \times N_f$ matrix

- $\mathcal{Z}^{N_f=1}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- $\mathcal{Z}^{N_f=1}(m, \mu) = \pi e^{-nm^2} \int_0^\infty du (u - \mu^2)^n I_0(2mn\sqrt{u}) e^{-nu}$

The Stephanov Model

- $\mathcal{Z} = \int DW e^{-n \Sigma^2 \text{Tr} W W^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$

Stephanov (1996) and Halasz, Jackson, Verbaarschot(1997)

- solve via bosonization

- $\mathcal{Z}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- where σ is an $N_f \times N_f$ matrix

- $\mathcal{Z}^{N_f=1}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- $\mathcal{Z}^{N_f=1}(m, \mu) = \pi e^{-nm^2} \int_0^\infty du (u - \mu^2)^n I_0(2mn\sqrt{u}) e^{-nu}$

The Stephanov Model

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr} WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$

Stephanov (1996) and Halasz, Jackson, Verbaarschot(1997)

- solve via bosonization

- $\mathcal{Z}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- where σ is an $N_f \times N_f$ matrix

- $\mathcal{Z}^{N_f=1}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$

- $\mathcal{Z}^{N_f=1}(m, \mu) = \pi e^{-nm^2} \int_0^\infty du (u - \mu^2)^n I_0(2mn\sqrt{u}) e^{-nu}$

The phase transition

- there is a phase transition separating a phase with zero and non-zero baryon density
- in the chiral limit $\mu_c = 0.527$ for $\mu \in \mathbb{R}$
- we can compute $\Sigma(m, \mu)$ and $n_B(m, \mu)$ and compare it with the Complex Langevin simulation
- first attempts in the Osborn model

$$\mathcal{Z} = \int D[W, W'] e^{-n \Sigma^2 \text{Tr}(WW^\dagger + W'W'^\dagger)} \det^{N_f} \begin{pmatrix} m & iW + \mu W' \\ iW^\dagger + \mu W'^\dagger & m \end{pmatrix}$$

Mollgaard and Splittorff (2013-2014), Nagata, Nishimura, Shimasaki (2015-2016)

However, $\mathcal{Z}(m, \mu) = (1 - \mu^2)^{nN_f} \mathcal{Z}(\frac{m}{\sqrt{1-\mu^2}}, 0)$

The phase transition

- there is a phase transition separating a phase with zero and non-zero baryon density
- in the chiral limit $\mu_c = 0.527$ for $\mu \in \mathbb{R}$
- we can compute $\Sigma(m, \mu)$ and $n_B(m, \mu)$ and compare it with the Complex Langevin simulation
- first attempts in the Osborn model

$$\mathcal{Z} = \int D[W, W'] e^{-n \Sigma^2 \text{Tr}(WW^\dagger + W'W'^\dagger)} \det^{N_f} \begin{pmatrix} m & iW + \mu W' \\ iW^\dagger + \mu W'^\dagger & m \end{pmatrix}$$

Mollgaard and Splittorff (2013-2014), Nagata, Nishimura, Shimasaki (2015-2016)

However, $\mathcal{Z}(m, \mu) = (1 - \mu^2)^{nN_f} \mathcal{Z}(\frac{m}{\sqrt{1-\mu^2}}, 0)$

The phase transition

- there is a phase transition separating a phase with zero and non-zero baryon density
- in the chiral limit $\mu_c = 0.527$ for $\mu \in \mathbb{R}$
- we can compute $\Sigma(m, \mu)$ and $n_B(m, \mu)$ and compare it with the Complex Langevin simulation
- first attempts in the Osborn model

$$\mathcal{Z} = \int D[W, W'] e^{-n \Sigma^2 \text{Tr}(WW^\dagger + W'W'^\dagger)} \det^{N_f} \begin{pmatrix} m & iW + \mu W' \\ iW^\dagger + \mu W' & m \end{pmatrix}$$

Mollgaard and Splittorff (2013-2014), Nagata, Nishimura, Shimasaki (2015-2016)

However, $\mathcal{Z}(m, \mu) = (1 - \mu^2)^{nN_f} \mathcal{Z}(\frac{m}{\sqrt{1-\mu^2}}, 0)$

Complex Langevin for RMT

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr}WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$
- $W = A + iB$
- compute the drift terms $\partial S/\partial A_{ij}$ and $\partial S/\partial B_{ij}$
- complexify the dof $A, B \in \mathbb{R} \rightarrow a, b \in \mathbb{C}$
- $A_{ij}(t + \delta t) = A_{ij}(t) - \partial_{A_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $B_{ij}(t + \delta t) = B_{ij}(t) - \partial_{B_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $\langle \xi_{ij} \rangle = 0$ and $\langle \xi_{ij}(t) \xi_{kl}(t') \rangle = 2\delta t \delta(t - t') \delta_{ik} \delta_{jl}$

Complex Langevin for RMT

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr}WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$
- $W = A + iB$
- compute the drift terms $\partial S / \partial A_{ij}$ and $\partial S / \partial B_{ij}$
- complexify the dof $A, B \in \mathbb{R} \rightarrow a, b \in \mathbb{C}$
- $A_{ij}(t + \delta t) = A_{ij}(t) - \partial_{A_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $B_{ij}(t + \delta t) = B_{ij}(t) - \partial_{B_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $\langle \xi_{ij} \rangle = 0$ and $\langle \xi_{ij}(t) \xi_{kl}(t') \rangle = 2\delta t \delta(t - t') \delta_{ik} \delta_{jl}$


Complex Langevin for RMT

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr}WW^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$
- $W = A + iB$
- compute the drift terms $\partial S / \partial A_{ij}$ and $\partial S / \partial B_{ij}$
- complexify the dof $A, B \in \mathbb{R} \rightarrow a, b \in \mathbb{C}$
- $A_{ij}(t + \delta t) = A_{ij}(t) - \partial_{A_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $B_{ij}(t + \delta t) = B_{ij}(t) - \partial_{B_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $\langle \xi_{ij} \rangle = 0$ and $\langle \xi_{ij}(t) \xi_{kl}(t') \rangle = 2\delta t \delta(t - t') \delta_{ik} \delta_{jl}$


Complex Langevin for RMT

- $\mathcal{Z} = \int DW e^{-n\Sigma^2 \text{Tr} W W^\dagger} \det^{N_f} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix}$
- $W = A + iB$
- compute the drift terms $\partial S / \partial A_{ij}$ and $\partial S / \partial B_{ij}$
- complexify the dof $A, B \in \mathbb{R} \rightarrow a, b \in \mathbb{C}$
- $A_{ij}(t + \delta t) = A_{ij}(t) - \partial_{A_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $B_{ij}(t + \delta t) = B_{ij}(t) - \partial_{B_{ij}} S(x(t)) \delta t + \delta \xi_{ij}$
- $\langle \xi_{ij} \rangle = 0$ and $\langle \xi_{ij}(t) \xi_{kl}(t') \rangle = 2\delta t \delta(t - t') \delta_{ik} \delta_{jl}$

m -scan for $\mu = 1$




$\langle\eta^\dagger\eta\rangle$ for $\mu = 1$




$\langle\bar{\eta}\eta\rangle$ for $\mu = 1$

analytical pq results by Glesaaen, Verbaarschot and SZ (2016)

μ -scan for $m = 0$




$\langle\eta^\dagger\eta\rangle$ for $m = 0$




$\langle\bar{\eta}\eta\rangle$ for $m = 0$

μ -scan for $m = 0.2$




$\langle\eta^\dagger\eta\rangle$ for $m = 0.2$




$\langle\bar{\eta}\eta\rangle$ for $m = 0.2$

μ -scan for $m = 1$



$\langle\eta^\dagger\eta\rangle$ for $m = 1$



$\langle\bar{\eta}\eta\rangle$ for $m = 1$

Gauge Cooling

Originally suggested for QCD by Seiler, Sexty and Stamatescu(2012)

implemented for RMT by Nagata, Nishimura, Shimasaki (2015) in the Osborn model

- complexified action invariant under an enhanced gauge transformation
- steer the evolution in Langevin time towards more physical solutions
- successful application to the Osborn model
- complexified action is invariant $GL(N, \mathbb{C})$
- $A' = \frac{1}{2} \left(hAh^{-1} + (hA^T h^{-1})^T + i(hBh^{-1} - (hB^T h^{-1})^T) \right)$
- $B' = \frac{1}{2} \left(hBh^{-1} + (hB^T h^{-1})^T - i(hAh^{-1} - (hA^T h^{-1})^T) \right)$

Gauge Cooling

Originally suggested for QCD by Seiler, Sexty and Stamatescu(2012)

implemented for RMT by Nagata, Nishimura, Shimasaki (2015) in the Osborn model

- complexified action invariant under an enhanced gauge trafo
- steer the evolution in Langevin time towards more physical solutions
- successful application to the Osborn model
- complexified action is invariant $GL(N, \mathbb{C})$
- $A' = \frac{1}{2} \left(hAh^{-1} + (hA^T h^{-1})^T + i(hBh^{-1} - (hB^T h^{-1})^T) \right)$
- $B' = \frac{1}{2} \left(hBh^{-1} + (hB^T h^{-1})^T - i(hAh^{-1} - (hA^T h^{-1})^T) \right)$

Gauge Cooling

Originally suggested for QCD by Seiler, Sexty and Stamatescu(2012)

implemented for RMT by Nagata, Nishimura, Shimasaki (2015) in the Osborn model

- complexified action invariant under an enhanced gauge trafo
- steer the evolution in Langevin time towards more physical solutions
- successful application to the Osborn model
- complexified action is invariant $GL(N, \mathbb{C})$
- $A' = \frac{1}{2} \left(hAh^{-1} + (hA^T h^{-1})^T + i(hBh^{-1} - (hB^T h^{-1})^T) \right)$
- $B' = \frac{1}{2} \left(hBh^{-1} + (hB^T h^{-1})^T - i(hAh^{-1} - (hA^T h^{-1})^T) \right)$

Gauge Cooling

Originally suggested for QCD by Seiler, Sexty and Stamatescu(2012)

implemented for RMT by Nagata, Nishimura, Shimasaki (2015) in the Osborn model

- complexified action invariant under an enhanced gauge trafo
- steer the evolution in Langevin time towards more physical solutions
- successful application to the Osborn model
- complexified action is invariant $GL(N, \mathbb{C})$
- $A' = \frac{1}{2} \left(hAh^{-1} + (hA^T h^{-1})^T + i(hBh^{-1} - (hB^T h^{-1})^T) \right)$
- $B' = \frac{1}{2} \left(hBh^{-1} + (hB^T h^{-1})^T - i(hAh^{-1} - (hA^T h^{-1})^T) \right)$

Gauge Cooling

Originally suggested for QCD by Seiler, Sexty and Stamatescu(2012)

implemented for RMT by Nagata, Nishimura, Shimasaki (2015) in the Osborn model

- complexified action invariant under an enhanced gauge trafo
- steer the evolution in Langevin time towards more physical solutions
- successful application to the Osborn model
- complexified action is invariant $GL(N, \mathbb{C})$
- $A' = \frac{1}{2} \left(hAh^{-1} + (hA^T h^{-1})^T + i(hBh^{-1} - (hB^T h^{-1})^T) \right)$
- $B' = \frac{1}{2} \left(hBh^{-1} + (hB^T h^{-1})^T - i(hAh^{-1} - (hA^T h^{-1})^T) \right)$

The Cooling norms

- $\mathcal{N}_H = \frac{1}{N} \text{tr} \left[(X - Y^\dagger)^\dagger (X - Y^\dagger) \right]$ (N.B. $X(0) = W$ and $Y(0) = W^\dagger$)

$X = A + iB$ and $Y = A^T - B^T$ when A, B are complexified)

- $\mathcal{N}_{AH} = \frac{1}{N} \text{tr} \left[\left((\phi + \psi^\dagger)^\dagger (\phi + \psi^\dagger) \right)^2 \right]$

- ψ and ϕ are the off-diagonal elements of D : $\psi = iX + \mu$,
 $\phi = iY + \mu$

- $\mathcal{N}_{\text{ev}} = \sum_{i=1}^{n_{\text{ev}}} e^{-\xi \gamma_i}$

- $\mathcal{N}_{\text{agg}} = (1 - s)N_{AH/\text{ev}} + sN_H$, where $s \in [0, 1]$

The Cooling norms

- $\mathcal{N}_H = \frac{1}{N} \text{tr} \left[(X - Y^\dagger)^\dagger (X - Y^\dagger) \right]$ (N.B. $X(0) = W$ and $Y(0) = W^\dagger$)

$X = A + iB$ and $Y = A^T - B^T$ when A, B are complexified)

- $\mathcal{N}_{AH} = \frac{1}{N} \text{tr} \left[\left((\phi + \psi^\dagger)^\dagger (\phi + \psi^\dagger) \right)^2 \right]$

- ψ and ϕ are the off-diagonal elements of D : $\psi = iX + \mu$,
 $\phi = iY + \mu$

- $\mathcal{N}_{\text{ev}} = \sum_{i=1}^{n_{\text{ev}}} e^{-\xi \gamma_i}$

- $\mathcal{N}_{\text{agg}} = (1 - s)N_{AH/\text{ev}} + sN_H$, where $s \in [0, 1]$

The Cooling norms

- $\mathcal{N}_H = \frac{1}{N} \text{tr} \left[(X - Y^\dagger)^\dagger (X - Y^\dagger) \right]$ (N.B. $X(0) = W$ and $Y(0) = W^\dagger$)

$X = A + iB$ and $Y = A^T - B^T$ when A, B are complexified)


- $\mathcal{N}_{AH} = \frac{1}{N} \text{tr} \left[\left((\phi + \psi^\dagger)^\dagger (\phi + \psi^\dagger) \right)^2 \right]$

- ψ and ϕ are the off-diagonal elements of D : $\psi = iX + \mu$,
 $\phi = iY + \mu$

- $\mathcal{N}_{\text{ev}} = \sum_{i=1}^{n_{\text{ev}}} e^{-\xi \gamma_i}$


- $\mathcal{N}_{\text{agg}} = (1 - s)N_{AH/\text{ev}} + sN_H$, where $s \in [0, 1]$

Cooling and the Dirac Spectrum





Scatter plots of the eigenvalues of the fermion matrix for a standard CL run together with the ones from a run cooled with the \mathcal{N}_{AH} cooling norm. The plots show the eigenvalues from the last 60 trajectories, separated by 100 updates. The left hand plot shows the Stephanov model, while the Osborn model is shown to the right.

Cooling and the Dirac Spectrum





Scatter plots of the eigenvalues of the fermion matrix for a standard CL run together with the ones from a gauge cooled run. We chose the parameters $\{\xi = 100, n_{\text{ev}} = 2\}$ for \mathcal{N}_{ev} . The plots show the eigenvalues from the last 60 trajectories, separated by 100 updates. The left hand plot shows the Stephanov model, while the Osborn model is shown to the right

The anti-hermiticity norm



The eigenvalue norm



Value of N_{ev} as a function of Langevin time. Stephanov model to the left, Osborn model to the right

The shifted representation

- shift μ away from the fermionic term by a COV

- initially $D = \begin{pmatrix} m & iA - B + \mu \\ iA^T + B^T + \mu & m \end{pmatrix}$

- Absorb μ into A with a COV $A' = A - i\mu$. The action in terms of A' and B is

$$S = N \text{tr} (A'^T A' - 2i\mu A' + \mu^2 + B^T B) - N_f \text{tr} \log (m^2 + X' Y')$$

$X' = A' + iB$ and $Y' = A'^T - iB^T$. Now, the μ dependence has shifted from the fermionic to the bosonic term.

- Computing again the CL force term ...
- Advantage of the shifted representation is that it starts in an anti-Hermitian state, and since CL is non-deterministic, the configurations could evolve to a different minimum.

The shifted representation

- shift μ away from the fermionic term by a COV

- initially $D = \begin{pmatrix} m & iA - B + \mu \\ iA^T + B^T + \mu & m \end{pmatrix}$

- Absorb μ into A with a COV $A' = A - i\mu$. The action in terms of A' and B is

$$S = N \text{tr} (A'^T A' - 2i\mu A' + \mu^2 + B^T B) - N_f \text{tr} \log (m^2 + X' Y')$$

$X' = A' + iB$ and $Y' = A'^T - iB^T$. Now, the μ dependence has shifted from the fermionic to the bosonic term.

- Computing again the CL force term ...
- Advantage of the shifted representation is that it starts in an anti-Hermitian state, and since CL is non-deterministic, the configurations could evolve to a different minimum.

The shifted representation

- shift μ away from the fermionic term by a COV

- initially $D = \begin{pmatrix} m & iA - B + \mu \\ iA^T + B^T + \mu & m \end{pmatrix}$

- Absorb μ into A with a COV $A' = A - i\mu$. The action in terms of A' and B is

$$S = N \text{tr} (A'^T A' - 2i\mu A' + \mu^2 + B^T B) - N_f \text{tr} \log (m^2 + X' Y')$$

$X' = A' + iB$ and $Y' = A'^T - iB^T$. Now, the μ dependence has shifted from the fermionic to the bosonic term.

- Computing again the CL force term ...
- Advantage of the shifted representation is that it starts in an anti-Hermitian state, and since CL is non-deterministic, the configurations could evolve to a different minimum.

The shifted representation

- shift μ away from the fermionic term by a COV

- initially $D = \begin{pmatrix} m & iA - B + \mu \\ iA^T + B^T + \mu & m \end{pmatrix}$

- Absorb μ into A with a COV $A' = A - i\mu$. The action in terms of A' and B is

$$S = N \text{tr} (A'^T A' - 2i\mu A' + \mu^2 + B^T B) - N_f \text{tr} \log (m^2 + X' Y')$$

$X' = A' + iB$ and $Y' = A'^T - iB^T$. Now, the μ dependence has shifted from the fermionic to the bosonic term.

- Computing again the CL force term ...
- Advantage of the shifted representation is that it starts in an anti-Hermitian state, and since CL is non-deterministic, the configurations could evolve to a different minimum.

The shifted representation

- shift μ away from the fermionic term by a COV

- initially $D = \begin{pmatrix} m & iA - B + \mu \\ iA^T + B^T + \mu & m \end{pmatrix}$


- Absorb μ into A with a COV $A' = A - i\mu$. The action in terms of A' and B is

$$S = N \text{tr} (A'^T A' - 2i\mu A' + \mu^2 + B^T B) - N_f \text{tr} \log (m^2 + X' Y')$$

$X' = A' + iB$ and $Y' = A'^T - iB^T$. Now, the μ dependence has shifted from the fermionic to the bosonic term.


- Computing again the CL force term ...
- Advantage of the shifted representation is that it starts in an anti-Hermitian state, and since CL is non-deterministic, the configurations could evolve to a different minimum.

Shifted representation and cooling



\mathcal{N}_{AH} as a function of Langevin time. Stephanov model (RHS), Osborn model (LHS). The Stephanov plot also includes the values from the shifted representation. These start at 0 for $t = 0$, but quickly shoot up to meet the unshifted curves. Although A and A' start out very different, they coincide after thermalization. This means that $\langle A' \rangle_{CL, shifted} = \langle A \rangle_{CL, standard} - i\mu$, and thus they converge to the same solution. **The advantage of the shifted representation is that it starts in an anti-Hermitian state, and due to the fact that CL is non-deterministic, the configurations could evolve to a different minimum.**

Shifted representation and cooling



\mathcal{N}_{AH} for the shifted representation of the Stephanov model as a function of Langevin time for 8×8 block matrices. The zoomed in plots show the evolution of \mathcal{N}_{AH} with the application of the cooling step. There are 50 gauge cooling transformations between each Langevin step.

The deformation technique

- another idea Ito and Nishimura (2017) is to deform the Dirac operator, to "move" its smallest eigenvalues and then extrapolate to zero deformation parameter at the end of the calculation.
- deformation is achieved by a finite temperature term given by the two lowest Matsubara frequencies $\pm\pi T$

$$Z(m, \mu; \alpha) = \int D[X, Y] \det \begin{pmatrix} m & X + \mu + i\Theta(\alpha) \\ Y + \mu + i\Theta(\alpha) & m \end{pmatrix}$$

where $D[X, Y] = d[X, Y]P(X, Y)$ and $\Theta(\alpha)$ is itself a block-matrix

$$\Theta(\alpha) = \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha \end{pmatrix}$$

where α can be thought as the lowest Matsubara frequency.

The deformation technique

- another idea [Ito and Nishimura \(2017\)](#) is to deform the Dirac operator, to "move" its smallest eigenvalues and then extrapolate to zero deformation parameter at the end of the calculation.
- deformation is achieved by a finite temperature term given by the two lowest Matsubara frequencies $\pm\pi T$

$$Z(m, \mu; \alpha) = \int D[X, Y] \det \begin{pmatrix} m & X + \mu + i\Theta(\alpha) \\ Y + \mu + i\Theta(\alpha) & m \end{pmatrix}$$

where $D[X, Y] = d[X, Y]P(X, Y)$ and $\Theta(\alpha)$ is itself a block-matrix

$$\Theta(\alpha) = \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha \end{pmatrix}$$

where α can be thought as the lowest Matsubara frequency.


The deformation technique

- Beyond a critical value of α the eigenvalue spectrum opens up and at this point chiral symmetry is restored. We can thus extrapolate from higher values in α for which there are no eigenvalues at the origin. In our studies we will see a gap opening at $\alpha \approx 1.0$
- But let's see this in practice...

The deformation technique



- Beyond a critical value of α the eigenvalue spectrum opens up and at this point chiral symmetry is restored. We can thus extrapolate from higher values in α for which there are no eigenvalues at the origin. In our studies we will see a gap opening at $\alpha \approx 1.0$
- But let's see this in practice...

The deformation technique





Scatter plots of eigenvalues from simulating with $N = 24$, $dt = 5 \times 10^{-5}$, $t_{\text{end}} = 5.0$, measured every 400 steps after thermalization. Both plots show $m = 0.2$ and $\mu = 0.5$ for varying values of the "temperature" α .

The deformation technique



The density of the CL forces for $\mu = 0.5$ (LHS) and $\mu = 0.35$ (RHS). Data gathered with a $t_{\text{final}} = 100$ run using $dt = 5 \times 10^{-5}$. If the decay fall is exponential (or faster) the CL will give the correct result but not if it decays as power law (or slower). Define α_c as the first α value that gives power law decay.

The deformation technique



Physical observables as a function of the parameter α . Analytic solutions (lines) for various values of the mass in addition to values from a simulation at $N = 48$ and $(m, \mu) = (0.2, 0.5)$. The parameters of the simulation correspond to the solid analytic line. Chiral condensate (LHS), baryon number density (RHS).

Natural questions to ask: Which points do you include in your extrapolation and which fitting ansatz do you use and what about the phase transition for small masses?



Reweighted Complex Langevin

- generate a CL trajectory for parameter values where complex Langevin is correct
- perform a reweighting of the trajectory to compute observables in an extended range of the parameters where CL used to fail

Reweighting

- **target** ensemble with parameters $\xi = (m, \mu, \beta)$ (for the general QCD case-drop β for RMT)
- **auxiliary** ensemble with parameters $\xi_0 = (m_0, \mu_0, \beta_0)$
- Reweighting from **auxiliary** to **target**

$$\begin{aligned}\langle O \rangle_\xi &= \frac{\int dx w(x; \xi) \mathcal{O}(x; \xi)}{\int dx w(x; \xi)} = \frac{\int dx w(x; \xi_0) \left[\frac{w(x; \xi)}{w(x; \xi_0)} \mathcal{O}(x; \xi) \right]}{\int dx w(x; \xi_0) \left[\frac{w(x; \xi)}{w(x; \xi_0)} \right]} \\ &= \frac{\langle \frac{w(x; \xi)}{w(x; \xi_0)} \mathcal{O}(x; \xi) \rangle_{\xi_0}}{\langle \frac{w(x; \xi)}{w(x; \xi_0)} \rangle_{\xi_0}}\end{aligned}$$

- but now we have a **complex weight** $w(x; \xi_0) = e^{-S(x; \xi_0)}$ so we need CL for this too!

Reweighting


- **target** ensemble with parameters $\xi = (m, \mu, \beta)$ (for the general QCD case-drop β for RMT)
- **auxiliary** ensemble with parameters $\xi_0 = (m_0, \mu_0, \beta_0)$
- Reweighting from **auxiliary** to **target**

$$\begin{aligned}\langle O \rangle_\xi &= \frac{\int dx w(x; \xi) \mathcal{O}(x; \xi)}{\int dx w(x; \xi)} = \frac{\int dx w(x; \xi_0) \left[\frac{w(x; \xi)}{w(x; \xi_0)} \mathcal{O}(x; \xi) \right]}{\int dx w(x; \xi_0) \left[\frac{w(x; \xi)}{w(x; \xi_0)} \right]} \\ &= \frac{\langle \frac{w(x; \xi)}{w(x; \xi_0)} \mathcal{O}(x; \xi) \rangle_{\xi_0}}{\langle \frac{w(x; \xi)}{w(x; \xi_0)} \rangle_{\xi_0}}\end{aligned}$$


- but now we have a **complex weight** $w(x; \xi_0) = e^{-S(x; \xi_0)}$ so we need CL for this too!

m -scan for $\mu = 1$

reweighted from an auxiliary ensemble with $m_0 = 4$ and $\mu_0 = 1$
using $\mathcal{O}(15000)$ confs




$\langle \eta^\dagger \eta \rangle$ for $\mu = 1$ and $n = 6$




$\langle \bar{\eta} \eta \rangle$ for $\mu = 1$ and $n = 6$

m -scan for $\mu = 1$

Auxiliary ensemble $m_0 = 1.3, \mu_0 = \mu = 1.0$ using $\mathcal{O}(560000)$ confs



$\langle \eta^\dagger \eta \rangle$ for $\mu = 1$ and $n = 24$




$\langle \bar{\eta} \eta \rangle$ for $\mu = 1$ and $n = 24$


The number of confs needed, builds up very rapidly. So one has to "just" to generate an auxiliary trajectory long enough to overcome the sign problem.

m -scan for $\mu = 1$

Auxiliary ensemble $m_0 = 1.3, \mu_0 = \mu = 1.0$ using $\mathcal{O}(122000000)$ confs



$\langle \eta^\dagger \eta \rangle$ for $\mu = 1$ and $n = 48$



$\langle \bar{\eta} \eta \rangle$ for $\mu = 1$ and $n = 48$

if you (or the computer) work(s) hard enough you can also deal with the matrices of the original size that we were dealing with $n = 48$

Conclusions and outlook

- studied the CL algorithm for an RMT model for QCD/
comparing numerical with exact analytical results for all the
range of parameters (m, μ)
- compared to previous similar studies this model possesses a
phase transition to a phase with non-zero baryon density
- fails to converge to QCD and it converges to $|QCD|$
- partial attempt to fix the issues via RCL procedure \rightarrow correct
results, even when CL does not work for the target ensemble
- gauge cooling or the deformation technique seem unable to
overcome the pathologies of this model

*Thanks a lot for your attention and many thanks to the organizers
for their warm hospitality!!!*

Stay Tuned!



for upcoming results ...

The analytical solution

Halasz, Jackson, Verbaarschot(1997)

$$\int \mathcal{D}[\dots] \exp \left[-i \sum_{k=1}^{N_f} \psi^{k*} \begin{pmatrix} m & iW + \mu \\ iW^\dagger + \mu & m \end{pmatrix} \psi^k \right] e^{-n\Sigma^2 \text{Tr} W W^\dagger}$$

perform the W integration

$$\int \mathcal{D}[\dots] e \left[-\frac{1}{n\Sigma^2} \psi_{Lk}^{f*} \psi_{Ri}^f \psi_{Ri}^{g*} \psi_{Lk}^g + m \left(\psi_{Ri}^{f*} \psi_{Ri}^f + \psi_{Lk}^{f*} \psi_{Lk}^g \right) + \mu \left(\psi_{Ri}^{f*} \psi_{Li}^f + \psi_{Lk}^{f*} \psi_{Rk}^g \right) \right]$$

write the four-fermion term as a difference of two squares and

linearize via a **Hubbard-Stratonovich trafo**

$$\exp(-AQ^2) \sim \int d\sigma \exp\left(-\frac{\sigma^2}{4A} - iQ\sigma\right)$$

The analytical solution

then one carries out the Grassmann integrals

$$Z(m, \mu) = \int \mathcal{D}\sigma \exp[-n\Sigma^2 \text{Tr}\sigma\sigma^\dagger] \det^n \begin{pmatrix} \sigma + m & \mu \\ \mu & \sigma^\dagger + m \end{pmatrix}$$

which for **one-flavor** and $\Sigma = 1$ becomes

$$\mathcal{Z}^{N_f=1}(m, \mu) = \int d\sigma d\sigma^* e^{-n\sigma^2} (\sigma\sigma^* + m(\sigma + \sigma^*) + m^2 - \mu^2)^n$$

The analytical solution

in polar coordinates after the angular integration

$\mathcal{Z}^{N_f=1}(m, \mu) = \pi e^{-nm^2} \int_0^\infty du (u - \mu^2)^n I_0(2mn\sqrt{u}) e^{-nu}$ in the thermodynamic limit one can perform a **saddle point analysis**

$$I_0(z) \sim e^z / \sqrt{2\pi z}$$

and the saddle point equation takes the form

$$\frac{1}{u - \mu^2} = 1 - \frac{m}{\sqrt{u}}$$

The analytical solution

A 1st order phase transition takes place at the points where $|Z_{u=u_b}| = |Z_{u=u_r}|$, with u_b and u_r two different solutions of the saddle-point equation give the same free-energy. This condition can be rewritten as $|(u_b - \mu^2)e^{2m\sqrt{u_b}-u_b}| = |(\mu^2 - u_r)e^{2m\sqrt{u_r}-u_r}|$. A general solution is quite cumbersome, but for $m \rightarrow 0$ we find that $u_r = 0$ and $u_b = 1 + \mu^2$. This leads to the critical curve $\text{Re}[1 + \mu^2 + \log \mu^2] = 0$ which for **real** μ , $\mu_c = 0.527\dots$

Phase Quenched QCD

- ignore the complex phase of the fermion determinant
- $\mathcal{Z}_{pq} = \mathcal{Z}_{iso} = \int dA |\det(D(\mu))|^2 e^{-S_g}$
- $|\det(D(\mu))|^2 = \det D(\mu) (\det(D(\mu)))^* = \det D(\mu) \det D(-\mu)$
- since $\gamma_5(\gamma_\mu D^\mu + m - \mu\gamma_0)\gamma_5 = (\gamma_\mu D^\mu + m + \mu\gamma_0)^\dagger$
- $\langle O \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int dA O |\det D(\mu)|^2 e^{-S_g}$
- this theory has a different phase diagram and different properties than QCD
- for $T \ll$ and $\mu \gg \rightarrow$ Bose condensation of charged pions

Phase Quenched QCD

- ignore the complex phase of the fermion determinant
- $\mathcal{Z}_{pq} = \mathcal{Z}_{iso} = \int dA |\det(D(\mu))|^2 e^{-S_g}$
- $|\det(D(\mu))|^2 = \det D(\mu) (\det(D(\mu)))^* = \det D(\mu) \det D(-\mu)$
- since $\gamma_5(\gamma_\mu D^\mu + m - \mu\gamma_0)\gamma_5 = (\gamma_\mu D^\mu + m + \mu\gamma_0)^\dagger$
- $\langle O \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int dA O |\det D(\mu)|^2 e^{-S_g}$
- this theory has a different phase diagram and different properties than QCD
- for $T \ll$ and $\mu \gg \rightarrow$ Bose condensation of charged pions

Phase Quenched QCD

- ignore the complex phase of the fermion determinant
- $\mathcal{Z}_{pq} = \mathcal{Z}_{iso} = \int dA |\det(D(\mu))|^2 e^{-S_g}$
- $|\det(D(\mu))|^2 = \det D(\mu) (\det(D(\mu)))^* = \det D(\mu) \det D(-\mu)$
- since $\gamma_5(\gamma_\mu D^\mu + m - \mu\gamma_0)\gamma_5 = (\gamma_\mu D^\mu + m + \mu\gamma_0)^\dagger$
- $\langle O \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int dA O |\det D(\mu)|^2 e^{-S_g}$
- this theory has a different phase diagram and different properties than QCD
- for $T \ll$ and $\mu \gg \rightarrow$ Bose condensation of charged pions

Phase Quenched QCD

- ignore the complex phase of the fermion determinant
- $\mathcal{Z}_{pq} = \mathcal{Z}_{iso} = \int dA |\det(D(\mu))|^2 e^{-S_g}$
- $|\det(D(\mu))|^2 = \det D(\mu) (\det(D(\mu)))^* = \det D(\mu) \det D(-\mu)$
- since $\gamma_5(\gamma_\mu D^\mu + m - \mu\gamma_0)\gamma_5 = (\gamma_\mu D^\mu + m + \mu\gamma_0)^\dagger$
- $\langle O \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int dA O |\det D(\mu)|^2 e^{-S_g}$
- this theory has a different phase diagram and different properties than QCD
- for $T \ll$ and $\mu \gg \rightarrow$ Bose condensation of charged pions

Phase Quenched QCD

- ignore the complex phase of the fermion determinant
- $\mathcal{Z}_{pq} = \mathcal{Z}_{iso} = \int dA |\det(D(\mu))|^2 e^{-S_g}$
- $|\det(D(\mu))|^2 = \det D(\mu) (\det(D(\mu)))^* = \det D(\mu) \det D(-\mu)$
- since $\gamma_5(\gamma_\mu D^\mu + m - \mu\gamma_0)\gamma_5 = (\gamma_\mu D^\mu + m + \mu\gamma_0)^\dagger$
- $\langle O \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int dA O |\det D(\mu)|^2 e^{-S_g}$
- this theory has a different phase diagram and different properties than QCD
- for $T \ll$ and $\mu \gg \rightarrow$ Bose condensation of charged pions