UNIVERSITY OF Southampton

Andreas Schmitt

Mathematical Sciences and STAG Research Centre
University of Southampton
Southampton SO17 1BJ, United Kingdom

STAG RESEARCH CENTRE

Dense nuclear and quark matter in holographic QCD

S.-w. Li, A. Schmitt, Q. Wang, PRD 92, 026006 (2015)
F. Preis, A. Schmitt, JHEP 1607, 001 (2016); EPJ Web Conf. 137, 09009 (2017)

- motivation: theoretical challenges in dense matter and relevance for compact stars
- the Sakai-Sugimoto model: holography as close to QCD as currently possible
- phase diagram including nuclear matter and quark matter

- **Dense QCD matter: what we know**



- first-order onset of nuclear matter at $\mu = 308\,\mathrm{MeV}$
- weakly interacting quark matter at asymptotically large μ
- as a consequence: must be chiral/deconfinement transition in between (presumably in strongly coupled regime)

- **Dense QCD matter: rigorous methods**



- QCD on the lattice: sign problem at nonzero μ, but recent progress
- perturbative QCD: restricted to ultra-high densities
- "standard" nuclear physics: input from experiment, restricted to nuclear saturation density

- **Dense QCD matter in compact stars**





equation of state over wide density regime highly desired!

- **Dense QCD matter: models**



- Nambu–Jona-Lasinio (usually no nuclear matter)
- quark-meson (no nucleons), nucleon-meson (no quarks)
- nucleon-quark-meson (patched together, many parameters)
- extrapolations from nuclear to weakly interacting quark matter

→ even without rigor: models for compact stars hard to construct!

- **Can holography help?**

J. M. Maldacena, Int. J. Theor. Phys. 38, 1113 (1999) [Adv. Theor. Math. Phys. 2, 231 (1998)]

- dual of QCD: probably exists, but currently out of reach
- reliable strong-coupling calculation (usually infinite coupling)

- Sakai-Sugimoto model:
  T. Sakai and S. Sugimoto, Prog. Theor. Phys. 113, 843 (2005)
  - top-down approach with only 3 parameters
  - dual to large-N_c QCD, however in inaccessible limit
  - contains all necessary ingredients:
    baryons, quark matter, chiral/deconfinement phase transitions

• Phases in the Sakai-Sugimoto model (page 1/3)



- in probe brane ("quenched") approximation: phase transition unaffected by quantities on flavor branes (μ, B, ...)
- not unlike expectation from large-N_c QCD

• Phases in the Sakai-Sugimoto model (page 2/3)

- less "rigid" behavior for smaller L
- deconfined, chirally broken phase for $L < 0.3\,\pi/M_{\rm KK}$

O. Aharony, J. Sonnenschein, S. Yankielowicz, Annals Phys. 322, 1420 (2007)

N. Horigome, Y. Tanii, JHEP 0701, 072 (2007)



• Phases in the Sakai-Sugimoto model (page 3/3)



- "decompactified" limit → gluon dynamics decouple
- "NJL-like" dual field theory

  E. Antonyan, J. A. Harvey, S. Jensen, D. Kutasov, hep-th/0604017

  J. L. Davis, M. Gutperle, P. Kraus, I. Sachs, JHEP 0710, 049 (2007)

  F. Preis, A. Rebhan and A. Schmitt, Lect. Notes Phys. 871, 51 (2013)

• Adding baryons (page 1/2)

- baryons in AdS/CFT: wrapped D-branes with N_c string endpoints
  E. Witten, JHEP 9807, 006 (1998); D. J. Gross, H. Ooguri, PRD 58, 106002 (1998)
- baryons in Sakai-Sugimoto:
  - D4-branes wrapped on S^4
  - equivalently: instantons on D8-branes ($\rightarrow$ skyrmions)

  T. Sakai, S. Sugimoto, Prog. Theor. Phys. 113, 843-882 (2005)

  H. Hata, T. Sakai, S. Sugimoto, S. Yamato, Prog. Theor. Phys. 117, 1157 (2007)



• Adding baryons (page 2/2)

- instanton solution for $SU(2)$ gauge fields in the bulk

$$F^2 \sim \frac{\rho^4}{[(\vec{x}-\vec{x}_0)^2+\frac{(z-z_0)^2}{\gamma^2}+\frac{\rho^2}{\gamma^2}]^4} \rightarrow \sum_{n=1}^{N_z}\sum_{i=1}^{i_n} \frac{\rho^4}{[(\vec{x}-\vec{x}_{in})^2+\frac{(z-z_n)^2}{\gamma^2}+\frac{\rho^2}{\gamma^2}]^4}$$

- solve EOMs for $U(1)$ gauge field $\hat{A}_0(u)$ and embedding $x_4(u)$
- minimize free energy wrt u_c, instanton shape ρ, γ, instanton number N_I, instanton layers N_z, z_n



- compare free energy of all three phases

• Phase diagrams with holographic baryons (page 1/4)

- first: without baryons

  N. Horigome and Y. Tanii,
  JHEP 0701, 072 (2007)



- first-order chiral phase transition

(dimensionless temperature $T \to TL$ and quark chemical potential $\mu \to 4\pi\lambda^{-1}L^2 M_{\rm KK}\mu$)

• Phase diagrams with holographic baryons (page 2/4)

- pointlike approximation: delta function instead of instanton profile

  O. Bergman, G. Lifschytz,

  M. Lippert, JHEP 0711, 056 (2007)



- second-order baryon onset
- no chiral restoration of dense matter at small T

• Phase diagrams with holographic baryons (page 3/4)

- finite-size instantons with dynamically determined width ρ

  S.-w. Li, A. Schmitt, Q. Wang, PRD 92, 026006 (2015)



- second-order baryon onset
- chiral restoration of dense matter at (very!) large μ

- **Further improving the instanton approximation**

F. Preis, A. Schmitt, JHEP 1607, 001 (2016)

- include instanton deformation γ and dynamic distribution of layers
- motivated by vacuum results: constraints $\rho = \rho_0 u_c$, $\gamma = \gamma_0 u_c^{3/2}$



- agreement with complementary approximations

"baryonic popcorn": V. Kaplunovsky, D. Melnikov, J. Sonnenschein, JHEP 1211, 047 (2012)
M. Elliot-Ripley, P. Sutcliffe and M. Zamaklar, JHEP 1610, 088 (2016)

• Phase diagrams with holographic baryons (page 4/4)

- deformed instanton layers
  F. Preis, A. Schmitt, JHEP 1607, 001 (2016)



- first-order baryon onset and chiral restoration
- but:
  - chiral transition very sensitive to ρ_0, γ_0
  - (relative) binding energy unrealistically large: $\sim 75\%$,
    with the QCD value $(939 - 16)/939 \simeq 98\%$ no chiral restoration
  - improvements necessary!

• Phase diagrams with holographic baryons (page 4/4)

- deformed instanton layers
  F. Preis, A. Schmitt, JHEP 1607, 001 (2016)



- first-order baryon onset and chiral restoration
- but:
  - chiral transition very sensitive to ρ_0, γ_0
  - (relative) binding energy unrealistically large: $\sim 75\%$,
    with the QCD value $(939 - 16)/939 \simeq 98\%$ no chiral restoration
  - improvements necessary!

• Phase diagrams with holographic baryons (page 4/4)

- deformed instanton layers
  F. Preis, A. Schmitt, JHEP 1607, 001 (2016)



- first-order baryon onset and chiral restoration
- but:
  - chiral transition very sensitive to ρ_0, γ_0
  - (relative) binding energy unrealistically large: $\sim 75\%$,
    with the QCD value $(939 - 16)/939 \simeq 98\%$ no chiral restoration
  - improvements necessary!

• Phase diagrams with holographic baryons (page 4/4)

- deformed instanton layers
  F. Preis, A. Schmitt, JHEP 1607, 001 (2016)



- first-order baryon onset and chiral restoration
- but:
  - chiral transition very sensitive to ρ_0, γ_0
  - (relative) binding energy unrealistically large: $\sim 75\%$,
    with the QCD value $(939 - 16)/939 \simeq 98\%$ no chiral restoration
  - improvements necessary!

• Outlook

- alternative, homogeneous ansatz (not based on instanton solutions)
  M. Rozali, H. H. Shieh, M. Van Raamsdonk and J. Wu, JHEP 0801, 053 (2008)
  S.-w. Li, A. Schmitt, Q. Wang, PRD 92, 026006 (2015)
  M. Elliot-Ripley, P. Sutcliffe and M. Zamaklar, JHEP 1610, 088 (2016)

- improve instanton approximation
  A. Schmitt, F. Preis, work in progress
  exact two-instanton solution K. Hashimoto *et al.*, Prog. Theor. Phys. 122, 427 (2009)
  approximation for instanton interactions S. Baldino *et al.*, arXiv:1703.08695 [hep-th]

- include magnetic field
  pointlike baryons: F. Preis, A. Rebhan, A. Schmitt, JPG 39, 054006 (2012)

- fit parameters to nuclear saturation properties,
  compute high-density EoS, mass/radius curve for hybrid stars
  holographic quark matter: C. Hoyos *et al.* PRL 117, 032501 (2016)