

# Chiral edge modes and crystalline phases in atomic synthetic ladders

*Friday, 27 October 2017 16:20 (25 minutes)*

Cold atomic gases endowed with a synthetic dimension are emerging as an ideal platform to address the interplay between interactions and static gauge fields. A fundamental question is whether these setups can give access to pristine two-dimensional phenomena, such as the fractional quantum Hall effect, and how. We show that unambiguous signatures of Laughlin-like states can be observed and characterized in synthetic ladders, thus being related to an unconventional fractional quantum Hall effect in the thin-torus limit. In particular we demonstrate the existence of a hierarchy of fractional insulating and conducting states, showing that the gapped states are characterized by density and magnetic order, whereas the gapless phases can support helical modes. Our analysis provides a guideline towards the observability and understanding of strongly correlated states of matter in synthetic ladders.

**Primary author:** ROSSINI, Davide (P)

**Co-authors:** Dr MAZZA, Leonardo (ENS, Paris); Mr CALVANESE STRINATI, Marcello (Scuola Normale Superiore); Dr DALMONTE, Marcello (ICTP, Trieste); Prof. FAZIO, Rosario (ICTP, Trieste); Dr BARBARINO, Simone (SISSA, Trieste)

**Presenter:** ROSSINI, Davide (P)

**Session Classification:** Talks on Specific Talks - III