



Contribution ID: 37

Type: **Poster contribution**

Search for B-meson decays to four baryons at Babar

Friday, April 21, 2017 5:00 PM (1 hour)

B mesons are the lightest mesons which can decay to various final states containing different baryons. Up to now, the discrepancy between the inclusive branching fraction of all B meson decay modes with at least a couple of baryons in the final state, measured by ARGUS to be $(6.8 \pm 0.6)\%$, and the sum of exclusive baryonic channels, averaged on neutral and positive B mesons at less than 1%, represents an open issue. Moreover, the measurement and comparison of exclusive branching fractions of baryonic B decays as well as studies on the dynamic of the decay, may allow better understanding of baryon production in B decays and, more generally, hadron fragmentation into baryons. We present here a search for the decay of a B meson in four baryons: $B \rightarrow p \bar{p} p \bar{p}$, not yet observed. The data set consists of about 470 million B-antiB pairs collected with the BABAR detector at the SLAC National Accelerator Laboratory.

Primary author: ZANI, Laura (PI)

Co-authors: ANULLI, Fabio (ROMA1); FORTI, Francesco (PI); Mr LUECK, Thomas (INFN sezione Pisa, University of Pisa)

Presenter: ZANI, Laura (PI)

Session Classification: Archivio Poster

Track Classification: Sessione Frontiera Intensità