



Contribution ID: 10

Type: **Oral contribution**

## **Study of the decays of the Higgs boson to boson pairs at Run-II with the ATLAS experiment**

*Wednesday, 19 April 2017 15:35 (15 minutes)*

The Higgs boson plays an important role in the Standard Model: it provides mass to the elementary particles through the electroweak spontaneous symmetry breaking (EWSB). Discovered in 2012 by the ATLAS and CMS collaborations, today the Higgs boson is a very powerful tool to test the validity of SM theory at the TeV scale. The Higgs boson can be produced at proton-proton colliders, such as LHC, in different ways and many decay channels are predicted by the SM. At ATLAS Run-2, this particle is widely studied mainly through its decays in vectors boson pairs ( $H \rightarrow ZZ$ ,  $H \rightarrow WW$  and  $H \rightarrow \gamma\gamma$ ). In this context, the latest results, obtained using the dataset collected by the ATLAS experiment during 2015 and 2016 at a center of mass energy of 13 TeV, are presented. Experimental techniques and estimate of the main backgrounds will be discussed in detail.

**Primary author:** SESSA, Marco (ROMA3)

**Presenter:** SESSA, Marco (ROMA3)

**Session Classification:** Sessione Frontiera Energia

**Track Classification:** Sessione Frontiera Energia