

Contribution ID: 48

Type: Oral contribution

Search for narrow resonances in dilepton mass spectra in p-p collisions at sqrt(s) = 13 TeV and combination with 8 TeV data

Thursday, 20 April 2017 18:10 (15 minutes)

any well established models extending beyond Standard Model (e.g. Grand Unified Theory, Sequential Standard Model or models proposing extra spatial dimension(s)) predict the existence of new heavy neutral bosons that would decay in two leptons. A search for new narrow resonances, generically referred as Z', in the dielectron and dimuon decay channels has been performed using data collected by the CMS experiment in 2016 from proton - proton collisions at a center of mass energy of sqrt(s) = 13 TeV (corresponding to an integrated luminosity of 13 fb-1) and combining 2015 13 TeV data (corresponding a luminosity of 2.9 fb-1) with a previous analysed set of data obtained at sqrt(s) = 8 TeV (corresponding a luminosity of 20 fb-1). In the absence of a significant deviation from the standard model predictions, 95% confidence level limits are set on the ratio of the production cross section times branching fraction for high-mass resonances to that for the Z boson. For several models, lower limits on the resonance mass are derived.

Primary author: ERRICO, Filippo (BA)

Presenter: ERRICO, Filippo (BA)

Session Classification: Sessione Frontiera Energia

Track Classification: Sessione Frontiera Energia