

## A Lambshift Polarimeter for Hydrogen Molecules

*Monday, 10 September 2018 18:00 (20 minutes)*

Usually, Lambshift polarimeters are used to measure the nuclear spin polarisation of hydrogen atomic beams or slow proton beams of about 1keV. As recent experiments aim to produce polarised hydrogen molecules (H<sub>2</sub> HD, D<sub>2</sub>), either by recombining atoms (IKP @FZ-Jülich) or with a new molecular-beam-source (BINP), a nuclear spin polarimeter for molecules is needed. As it will be shown in this talk it is also possible to measure the nuclear spin polarisation of the hydrogen molecules with a Lambshift polarimeter.

**Primary author:** Mr HUXOLD, Lukas (HHU Duesseldorf)

**Co-authors:** Prof. TOPORKOV, Dmitriy (Budker institute of nuclear Physics, Novosibirsk State University); AWWAD, Hani (HHU Düsseldorf); Dr GRIGORYEV, Kirill (Institut für Kernphysik, Forschungszentrum Jülich); Prof. BÜSCHER, Markus (Forschungszentrum Jülich / PGI-6); Dr ENGELS, Ralf (Institut für Kernphysik, Forschungszentrum Jülich); SHES-TAKOV, Yuri (Budker institute of nuclear Physics, Novosibirsk State University)

**Presenter:** Mr HUXOLD, Lukas (HHU Duesseldorf)

**Session Classification:** Polarized Ion and Lepton Sources and Targets

**Track Classification:** Polarized Ion and Lepton Sources and Targets