

Double helicity asymmetries for π^\pm production in $p + p$ collisions at $\sqrt{s}=510$ GeV at PHENIX Mid-rapidity

Wednesday, 12 September 2018 17:00 (20 minutes)

One of the main goals of the RHIC spin program is the determination of the gluon helicity contribution to the proton spin. This can be accessed by measuring double helicity asymmetries (A_{LL}) of pion production at mid-rapidity in longitudinally polarized proton collisions. The ordering of the asymmetries with the charge of the final state pions can in addition directly infer the sign of the gluon spin contribution.

Charged pions are reconstructed in the central PHENIX tracking system. The asymmetries are evaluated between the collisions of bunches with the same and opposite helicity after correcting for differences in luminosity and for beam polarizations.

To extend our understanding of the gluon polarization to a lower gluon momentum fraction (x), high statistics data was collected at a higher $\sqrt{s}=510$ GeV in 2012-2013.

We present the physics motivation, the analysis procedure, current status of the π^\pm A_{LL} measurements and comparisons to π^0 as well as π^\pm at $\sqrt{s}=200$ GeV.

Primary author: Mr MOON, Taebong (Yonsei University/RIKEN)

Presenter: Mr MOON, Taebong (Yonsei University/RIKEN)

Session Classification: Nucleon helicity structure

Track Classification: Nucleon Helicity Structure