



Contribution ID: 288

Type: poster

## Multi-MeV, stable electron bunch using supersonic air jet target

*Monday, 25 September 2017 19:30 (1 hour)*

Relativistic interaction of short-pulse, high intensity lasers with underdense plasmas results in acceleration of electrons in relativistic regime. These relativistic electron beams have the ability to serve as a source of spatially coherent collimated, point-like and femtosecond X-ray radiation. Generation of a quasi-monoenergetic bunch of electrons is important for generation of stable betatron radiation and X-ray pulses from inverse Compton scattering.

The purpose this work was to optimize conditions necessary for building this type of electron and X-ray source in a small laboratory equipped with a common femtosecond few-terawatt laser. Various gas jet targets, such as helium, a mixture of helium and argon, helium with an admixture of synthetic air, and dry air were tested for this purpose using different backing pressures and the obtained results are compared. Additionally, a razor blade was used to create steep density gradient of the neutral gas in order to achieve shock injection and thus to increase the number of trapped electrons. Simulations and experimental results are presented.

**Primary author:** Dr CHAULAGAIN, Uddhab (ELI beamlines)

**Co-authors:** NEJDL, Jaroslav (Institute of Plasma Physics ASCR); BOHACEK, Karel (2Czech Technical University in Prague, Faculty of Nuclear Sciences and Physical Engineering); TA PHUOC, Kim (Laboratoire d'Optique Appliquée, ENSTA); KOZLOVA, Michaela (Institute of Plasma Physics ASCR); KRUS, Miroslav (Institute of Plasma Physics ASCR); HORNY, Vojtěch (Institute of Plasma Physics ASCR)

**Presenter:** Dr CHAULAGAIN, Uddhab (ELI beamlines)

**Session Classification:** Wine and Poster Session 1(WG1-WG2-WG3-WG8)

**Track Classification:** WG1 - Electron Beams from Plasmas