



Contribution ID: 226

Type: talk

THz-Driven Accelerator Components

Monday, September 25, 2017 6:00 PM (20 minutes)

Plasmon- or phonon-polaritons excited at plane or structured interfaces or in in sub-wavelength resonators by strong THz pulses are interesting candidates for miniaturized accelerator components. Often these structures show either electric or magnetic near-field enhancement alleviating the need for a strong driving THz source. Today laser-driven THz sources can produce single-cycle pulses with field strengths between 10 to 100 MV/m (electric) and around several Tesla (magnetic); they can be easily synchronized to a gun laser or any other laser in the accelerator chain and the THz free-space wavelength or the plasmon wavelength are well matched to a typical transverse electron bunch size on the order of 100 microns.

Here, we will discuss deflecting structures for electron streaking diagnostics, accelerating structures, and also miniaturized undulator structures.

Primary author: Prof. FEURER, Thomas (University of Bern)

Co-authors: Ms HAYATI, Mozghan (University of Bern); Dr TARKESHIAN, Roxana (Universität Bern); Dr OLLMANN, Zoltan (University of Bern)

Presenter: Prof. FEURER, Thomas (University of Bern)

Session Classification: WG5_Parallel

Track Classification: WG5 - High-Gradient Plasma Structures/Advanced Beam Diagnostics