



# ALICE ACTIVITIES IN TRIESTE



<https://webint.ts.infn.it/ricerca/exp/alice/alice-its/welcome.html>



## A LARGE ION COLLIDER EXPERIMENT

The ALICE Collaboration has built a dedicated detector to exploit the unique physics potential of nucleus-nucleus collisions at LHC energies. Its aim is to study the physics of strongly interacting matter at the highest energy densities reached so far in the laboratory. In such condition, an extreme phase of matter – called the **quark-gluon plasma** – is formed.

Our universe is thought to have been in such a primordial state for the first few millionths of a second after the Big Bang. The properties of such a phase are key issues for Quantum Chromodynamics, the understanding of confinement-deconfinement and chiral phase transitions.

For this purpose, we are carrying out a comprehensive study of the hadrons, electrons, muons and photons produced in the collisions of heavy nuclei.

ALICE is also studying proton-proton and proton-nucleus collisions both as a comparison with nucleus-nucleus collisions and in their own right.



**Inner Tracking System (ITS):**




- Primary vertex
- Tracking
- Particle identification via  $dE/dx$

**Time Projection Chamber (TPC):**

- Global tracking
- Particle identification via  $dE/dx$

**Time Of Flight (TOF):**


- Particle identification via velocity measurement



## ITS UPGRADE

Aimed to replace ALICE ITS during the LHC Long Shutdown 2 in 2019/20


- Increase readout speed → Readout Pb-Pb collisions at 100 kHz
- Improve impact parameter resolution
  - Reduce pixel size:  $0.50 \times 425 \mu\text{m}^2 \rightarrow 0.30 \times 30 \mu\text{m}^2$
  - Reduce distance from the IP: 39 mm → 23 mm
- Reduce inner layers X0:  $\sim 1.14\% \rightarrow \sim 0.3\%$
- Improve tracking efficiency at low  $p_T$
- All 7 layers with binary pixels



**12.5 Gigapixels**  
**10 m<sup>2</sup> sensitive area**

ITS requirements	Inner Barrel	Outer Barrel
Spatial resolution	5 $\mu\text{m}$	10 $\mu\text{m}$
Detection efficiency	> 99 %	
Fake hit rate	< $10^{-6}$ pixel <sup>-1</sup> event <sup>-1</sup>	
Power density	< 300 mW cm <sup>-2</sup>	< 100 mW cm <sup>-2</sup>
TID radiation hardness	2700 krad	100 krad
NIEL radiation hardness	$1.7 \times 10^{13}$ 1 MeV n <sub>eq</sub> cm <sup>-2</sup>	$10^{12}$ 1 MeV n <sub>eq</sub> cm <sup>-2</sup>

## TECHNOLOGY




TowerJazz 180 nm CMOS imaging sensor process


- Deep p-well shielding n-well allowing in-pixel PMOS
- More complex in-pixel circuitry
- High-resistivity (> 1 k $\Omega$  cm) p-type epitaxial layer (18 to 30  $\mu\text{m}$ ) on p-type substrate
- Substrate bias → Increase of depletion volume
- Larger charge collected by seed pixel
- Lower input capacitance → better S/N ratio
- Short collection time
- Better non-ionising radiation tolerance

## BEAM TEST RESULTS

### SPATIAL RESOLUTION



### DETECTION EFFICIENCY




## FOR INFORMATION:

- Prof. Paolo Camerini (paolo.camerini@ts.infn.it)
- Prof. Giacomo-Vito Margagliotti (giacomo.margagliotti@ts.infn.it)

- Prof. Rinaldo Rui (rinaldo.rui@ts.infn.it)
- Dott. Stefano Piano (stefano.piano@ts.infn.it)

## THE LARGE HADRON COLLIDER





The Large Hadron Collider (LHC) is a synchrotron built 100 meters underground at the border of Switzerland and France able to accelerate pp, p-Pb and Pb-Pb. It has a circumference of 27 km and is instrumented with 8-Tesla superconducting magnets, providing the bending power to orbit 13 TeV protons in opposite directions.

## ANALYSIS ACTIVITIES


### HYPERNUCLEI PRODUCTION

$^3\text{H}$  is the lightest known hypernucleus and is formed by (p,n, $\Lambda$ ).  
Mass = 2.991 GeV/c<sup>2</sup>  $B_\Lambda = 0.13 \pm 0.05$  MeV Lifetime ~ 263 ps

- Signal extracted in 3 pt bins for 2 centrality classes:
  - The measured yield is in agreement with equilibrium thermal model expectations
  - The measured lifetime is smaller w.r.t. the free  $\Lambda$  one



### LIGHT NUCLEI ELLIPTIC FLOW




Angular distribution of reconstructed charged particles can be expanded into a Fourier series w.r.t. symmetry plane  $\Psi_n$ :

$$E \frac{d^3N}{dp^3} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left( 1 + \sum_{n=1}^{\infty} 2v_n \cos[n(\phi - \Psi_n)] \right)$$
$$v_n = \langle \cos[n(\phi - \Psi_n)] \rangle$$


### Blast-Wave vs. Coalescence:

- deuteron  $v_2$  follows the mass ordering
- a Blast-Wave (red curve) parameterization obtained from lower mass species can describe the deuteron  $v_2$  reasonably well
- a simple coalescence model (magenta band) is not able to reproduce the measured  $v_2$




### PRODUCTION OF RESONANCES

- Hadronic resonances decay under the strong interaction with lifetimes of the same order of magnitude as that of the fireball created in Pb-Pb collisions.
- After chemical freeze-out resonances decay and can undergo re-scattering and regeneration depending on:
  - lifetime of the hadronic phase
  - lifetime of resonances
  - scattering cross-section of the decay products
- Key measurements:
  - Resonance yields and ratios to long-lived particles vs. centrality
  - Spectra down to low transverse momentum ( $p_T$ )
  - Resonances with different lifetimes




### MEASUREMENT OF D-MESONS PRODUCTION

- Heavy flavours as a probe of the Quark-Gluon Plasma:
  - Heavy flavours (i.e. charm and beauty quarks) are produced in the initial stages of the collision:
    - Heavy-flavours production time:  $t_{\text{prod}} < \hbar/m_{c(b)} \sim 0.1(0.4)$  fm/c
    - Quark-Gluon Plasma formation time at LHC [1]:  $t_{\text{form}} \sim 0.3$  fm/c
- Heavy flavours experience the whole system evolution interacting with the medium constituents



### FEMTOSCOPY




Femtoscopia is based on the measurement of the two-particle correlation function:

$$C(k^*) = \frac{N_{\text{same}}(k^*)}{N_{\text{mixed}}(k^*)} \quad k^* = \frac{1}{2} |p_1 - p_2|$$
$$p_1 + p_2 = 0$$

Sensitive to the size of particle emission region and their final state interactions:

- Exploring interactions relatively which are unknown with femtoscopia!

## COMPUTING



- The ALICE experiment has originally been designed as a relatively low-rate experiment: this will not be the case anymore for the Run 3 that is scheduled to start in 2019: expected rate ~100 kHz
- A part of ALICE upgrade includes major improvements in the computing side of the experiment:
  - we are involved in the development of a virtual infrastructure based on Cloud Computing for the parallel data analysis of the ALICE experiment

CPU	10000 xHS06 - 10 TFlops	84 WIN's 790 cores
DISK	300 TBn + 400 TBn	2.7 TB RAM
TAPE	80/150 TBn (DSM)	Quantum Scalar 12000 (LTO4/LTO5)
NET	Storage 8 Gb/s Fibre Channel	Ethernet 1 - 10 Gb/s

<http://webint.ts.infn.it/ricerca/exp/alice.html>  
<http://df.units.it/?q=it/node/3167>