

A. Incicchitti
INFN sez. Roma
Assisi, October 2009

A stylized logo for DAMA (Dark Matter And MAterial) featuring a large, abstract shape composed of various colored segments (red, blue, purple, brown) and a central purple beam-like element. The word "Dama" is written in a cursive font across the bottom of the logo.

Particle Dark Matter and DAMA/LIBRA






Accelerators

Cosmology and Astrophysics

Indirect search

Direct search

Complementary information



Different information can be only matched by logic grids in model dependent scenarios
A model independent approach can be pursued by direct detection
to investigate DM particle component in the Galactic halo


Relic DM particles from primordial Universe

Heavy candidates:

- In thermal equilibrium in the early stage of Universe
- Non relativistic at decoupling time:
 $\langle \sigma_{\text{ann}} \cdot v \rangle \sim 10^{-26} / \Omega_{\text{WIMP}} h^2 \text{ cm}^3 \text{ s}^{-1} \rightarrow \sigma_{\text{ordinary matter}} \sim \sigma_{\text{weak}}$
- Expected flux: $\Phi \sim 10^7 \cdot (\text{GeV}/m_{\text{W}}) \text{ cm}^{-2} \text{ s}^{-1}$ ($0.2 < \rho_{\text{halo}} < 1.7 \text{ GeV cm}^{-3}$)
- Form a dissipationless gas trapped in the gravitational field of the Galaxy ($v \sim 10^{-3}c$)
- Neutral, massive, stable (or with half life \sim age of Universe) and weakly interacting

Light candidates:

axion, sterile neutrino, axion-like particles cold or warm DM (no positive results from direct searches for relic axions with resonant cavity)




+ multi-component halo?

even a suitable particle not yet foreseen by theories

Some direct detection processes:

- Scatterings on nuclei

→ detection of nuclear recoil energy



- Inelastic Dark Matter: $W + N \rightarrow W^* + N$

→ W has Two mass states χ_+ , χ_- with δ mass splitting

→ Kinematical constraint for the inelastic scattering of χ_- on a nucleus

$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$


- Excitation of bound electrons in scatterings on nuclei

→ detection of recoil nuclei + e.m. radiation

- Interaction only on atomic electrons

→ detection of e.m. radiation


... even WIMPs



e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the electromagnetic component of their counting rate

- Conversion of particle into e.m. radiation


→ detection of γ , X-rays, e^-



- Interaction of light DMp (LDM) on e^- or nucleus with production of a lighter particle

→ detection of electron/nucleus recoil energy


e.g. sterile ν



• ... and more

Investigating the presence of a DM particle component in the galactic halo by the model independent annual modulation signature

Drukier, Freese, Spergel PRD86
Freese et al. PRD88



- $v_{\text{sun}} \sim 232 \text{ km/s}$ (Sun velocity in the halo)
- $v_{\text{orb}} = 30 \text{ km/s}$ (Earth velocity around the Sun)
- $\gamma = \pi/3$
- $\omega = 2\pi/T$ $T = 1 \text{ year}$
- $t_0 = 2^{\text{nd}} \text{ June}$ (when v_{\oplus} is maximum)

$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos\gamma \cos[\omega(t-t_0)]$$


$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

Expected rate in given energy bin changes because of the Earth's motion around the Sun moving in the Galaxy


Requirements:

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2nd June)
- 5) For single hit in a multi-detector set-up
- 6) With modulated amplitude in the region of maximal sensitivity < 7% (for usually adopted halo distributions, but it can be larger in case of some possible scenarios)

To mimic this signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements



DAMA: an observatory for rare processes @LNGS



DAMA/NaI : ≈ 100 kg NaI(Tl)

Performances: N.Cim.A112(1999)545-575, EPJC18(2000)283,
Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

Results on rare processes:

- Possible Pauli exclusion principle violation PLB408(1997)439
- CNC processes PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) PLB460(1999)235
- Search for solar axions PLB515(2001)6
- Exotic Matter search EPJdirect C14(2002)1
- Search for superdense nuclear matter EPJA23(2005)7
- Search for heavy clusters decays EPJA24(2005)51

Results on DM particles:

- PSD PLB389(1996)757
- Investigation on diurnal effect N.Cim.A112(1999)1541
- Exotic Dark Matter search PRL83(1999)4918
- Annual Modulation Signature

PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23,
EPJC18(2000)283, PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503,
Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127, IJMPA21(2006)1445,
EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506,
MPLA23(2008)2125, other works in progress ...



*data taking completed on July 2002,
last data release 2003: total exposure
(7 annual cycles) 0.29 ton x yr*

model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

DAMA/LIBRA ~250 kg NaI(Tl) (Large sodium Iodide Bulk for RARE processes)



As a result of a second generation R&D for more radiopure NaI(Tl)
by exploiting new chemical/physical radiopurification techniques
(all operations involving crystals and PMTs - including photos - in HP Nitrogen atmosphere)



improving installation
and environment



PMT
+HV
divider

Cu etching with
super- and ultra-
pure HCl solutions,
dried and sealed in
HP N₂



storing new crystals



etching staff at work
in clean room



**The new DAMA/LIBRA set-up ~250 kg NaI(Tl)
(Large sodium Iodide Bulk for RARE processes)**




installing DAMA/LIBRA detectors



assembling a DAMA/ LIBRA detector


filling the inner Cu box with further shield



**closing the Cu box
housing the detectors**



detectors during installation; in the central and right up detectors the new shaped Cu shield surrounding light guides (acting also as optical windows) and PMTs was not yet applied




view at end of detectors' installation in the Cu box

The calibration system



Some on residual contaminants in new NaI(Tl) detectors




α/e pulse shape discrimination has practically 100% effectiveness in the MeV range

The measured α yield in the new DAMA/LIBRA detectors ranges from 7 to some tens $\alpha/\text{kg}/\text{keV}$

Second generation R&D for new DAMA/LIBRA crystals: new selected powders, physical/chemical radiopurification, new selection of overall materials, new protocol for growing and handling

^{232}Th residual contamination From time-amplitude method. If ^{232}Th chain at equilibrium: it ranges from 0.5 ppt to 7.5 ppt



^{238}U residual contamination First estimate: considering the measured α and ^{232}Th activity, if ^{238}U chain at equilibrium \Rightarrow ^{238}U contents in new detectors typically range from 0.7 to 10 ppt



^{238}U chain splitted into 5 subchains: $^{238}\text{U} \rightarrow ^{234}\text{U} \rightarrow ^{230}\text{Th} \rightarrow ^{226}\text{Ra} \rightarrow ^{210}\text{Pb} \rightarrow ^{206}\text{Pb}$

Thus, in this case: (2.1 ± 0.1) ppt of ^{232}Th ; (0.35 ± 0.06) ppt for ^{238}U and: (15.8 ± 1.6) $\mu\text{Bq}/\text{kg}$ for $^{234}\text{U} + ^{230}\text{Th}$; (21.7 ± 1.1) $\mu\text{Bq}/\text{kg}$ for ^{226}Ra ; (24.2 ± 1.6) $\mu\text{Bq}/\text{kg}$ for ^{210}Pb .

natK residual contamination
The analysis has given for the natK content in the crystals values not exceeding about 20 ppb




^{129}I and ^{210}Pb
 $^{129}\text{I}/\text{natI} \approx 1.7 \times 10^{-13}$ for all the new detectors
 ^{210}Pb in the new detectors: $(5 - 30)$ $\mu\text{Bq}/\text{kg}$.

No sizeable surface pollution by Radon daughters, thanks to the new handling protocols

... more on NIMA592(2008)297





DAMA/LIBRA: calibrations at low energy

Studied by using various external gamma sources (^{241}Am , ^{133}Ba) and internal X-rays or gamma's (^{40}K , ^{125}I , ^{129}I)



The curves superimposed to the experimental data have been obtained by simulations

- **Internal ^{40}K :** 3.2 keV due to X-rays/Auger electrons (tagged by 1461 keV γ in an adjacent detector).
- **Internal ^{125}I :** 67.3 keV peak (EC from K shell + 35.5 keV γ) and composite peak at 40.4 keV (EC from L,M,... shells + 35.5 keV γ).
- **External ^{241}Am source:** 59.5 keV γ peak and 30.4 keV composite peak.
- **External ^{133}Ba source:** 81.0 keV γ peak.
- **Internal ^{129}I :** 39.6 keV structure (39.6 keV γ + β spectrum).






$$\frac{\sigma_{LE}}{E} = \frac{(0.448 \pm 0.035)}{\sqrt{E(\text{keV})}} + (9.1 \pm 5.1) \cdot 10^{-3}$$

Routine calibrations with ^{241}Am

DAMA/LIBRA: calibrations at high energy

The data are taken on the full energy scale up to the MeV region by means QADC's

Studied by using external sources of gamma rays (e.g. ^{137}Cs , ^{60}Co and ^{133}Ba) and gamma rays of 1461 keV due to ^{40}K decays in an adjacent detector, tagged by the 3.2 keV X-rays





$$\frac{\sigma_{HE}}{E} = \frac{(1.12 \pm 0.06)}{\sqrt{E(\text{keV})}} + (17 \pm 23) \cdot 10^{-4}$$


The signals (unlike low energy events) for high energy events are taken only from one PMT

Noise rejection near the energy threshold

Typical pulse profiles of PMT noise and of scintillation event with the same area, just above the energy threshold of 2 keV



The different time characteristics of PMT noise (decay time of order of tens of ns) and of scintillation event (decay time about 240 ns) can be investigated building several variables



From the Waveform Analyser
2048 ns time window:

$$X_1 = \frac{\text{Area (from 100 ns to 600 ns)}}{\text{Area (from 0 ns to 600 ns)}}$$

$$X_2 = \frac{\text{Area (from 0 ns to 50 ns)}}{\text{Area (from 0 ns to 600 ns)}}$$

- The separation between noise and scintillation pulses is very good.
- Very clean samples of scintillation events selected by stringent acceptance windows.
- The related efficiencies evaluated by calibrations with ^{241}Am sources of suitable activity in the same experimental conditions and energy range as the production data (efficiency measurements performed each ~ 10 days; typically 10^4 - 10^5 events per keV collected)

This is the only procedure applied to the analysed data

Infos about DAMA/LIBRA data taking

EPJC 56 (2008) 333

- DAMA/LIBRA test runs: from March 2003 to September 2003
- DAMA/LIBRA normal operation: from September 2003 to August 2004
- High energy runs for TDs: September 2004
to allow internal α 's identification
(approximative exposure $\approx 5000 \text{ kg} \times \text{d}$)
- DAMA/LIBRA normal operation: from October 2004

Data released here:

- four annual cycles: $0.53 \text{ ton} \times \text{yr}$
- calibrations: acquired $\approx 44 \text{ M}$ events from sources
- acceptance window eff: acquired $\approx 2 \text{ M}$ events/keV

Period		Exposure (kg×day)	$\alpha - \beta^2$
DAMA/LIBRA-1	Sept. 9, 2003 - July 21, 2004	51405	0.562
DAMA/LIBRA-2	July 21, 2004 - Oct. 28, 2005	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 - July 18, 2006	39445	0.591
DAMA/LIBRA-4	July 19, 2006 - July 17, 2007	49377	0.541
Total		192824 $\approx 0.53 \text{ ton} \times \text{yr}$	0.537

DAMA/NaI (7 years) + DAMA/LIBRA (4 years)

total exposure: $300555 \text{ kg} \times \text{day} = 0.82 \text{ ton} \times \text{yr}$



Two remarks:

- One PMT problems after 6 months. Detector out of trigger since Sep. 2003 (since Sep. 2008 again in operation)
- Residual cosmogenic ^{125}I presence in the first year in some detectors (this motivates the Sept. 2003 as starting time)

**DAMA/LIBRA is
continuously running**

Model independent annual modulation result

DAMA/NaI (7 years) + DAMA/LIBRA (4 years) Total exposure: 300555 kg×day = 0.82 ton×yr
EPJC 56(2008)333



$A \cos[\omega(t-t_0)]$

continuous lines: $t_0 = 152.5$ d, $T = 1.00$ y

$A = (0.0129 \pm 0.0016)$ cpd/kg/keV

$\chi^2/\text{dof} = 54.3/66$ 8.2σ C.L.

Absence of modulation? No

$\chi^2/\text{dof} = 116.4/67 \Rightarrow P(A=0) = 1.8 \cdot 10^{-4}$

from the fit with all the parameters free:


$A = (0.0131 \pm 0.0016)$ cpd/kg/keV

$t_0 = (144 \pm 8)$ d

$T = (0.998 \pm 0.003)$ y

Experimental single-hit residuals rate vs time and energy in 2-6 keV over 11 annual cycles

experimental residual rate of the multiple hit events (DAMA/LIBRA 1-4) in the 2-6 keV energy interval:
 $A = -(0.0004 \pm 0.0006)$ cpd/kg/keV



Multiple hits events = Dark Matter particle "switched off"


No systematics or side reaction able to account for the measured modulation amplitude and to satisfy all the peculiarities of the signature

This result offers an additional strong support for the presence of DM particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background

model independent evidence of a particle Dark Matter component in the galactic halo at 8.2σ C.L.

Can a hypothetical background modulation account for the observed effect?

• No Modulation above 6 keV



Mod. Ampl. (6-10 keV): (0.0016 ± 0.0031) , $-(0.0010 \pm 0.0034)$, $-(0.0001 \pm 0.0031)$ and $-(0.0006 \pm 0.0029)$ cpd/kg/keV for DAMA/LIBRA-1, DAMA/LIBRA-2, DAMA/LIBRA-3, DAMA/LIBRA-4;
 → they can be considered statistically consistent with zero

+

In the same energy region where the effect is observed: no modulation of the multiple-hits events (see next slide)

• No modulation in the whole spectrum:

studying integral rate at higher energy, R90

- R_{90} percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA-1,2,3,4 running periods


→ cumulative gaussian behaviour with $\sigma \approx 1\%$, fully accounted by statistical considerations

- Fitting the behaviour with time, adding a term modulated according period and phase expected for Dark Matter particles:

Period	Mod. Ampl.
DAMA/LIBRA-1	$-(0.05 \pm 0.19)$ cpd/kg
DAMA/LIBRA-2	$-(0.12 \pm 0.19)$ cpd/kg
DAMA/LIBRA-3	$-(0.13 \pm 0.18)$ cpd/kg
DAMA/LIBRA-4	(0.15 ± 0.17) cpd/kg

consistent with zero

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region → $R_{90} \sim$ tens cpd/kg → $\sim 100 \sigma$ far away



**No modulation in the background:
 these results account for all sources of bckg (+ see later)**


Multiple-hits events in the region of the signal - DAMA/LIBRA 1-4

- Each detector has its own TDs read-out
→ pulse profiles of multiple-hits events (multiplicity > 1) acquired (exposure: 0.53 ton×yr).
- The same hardware and software procedures as the ones followed for single-hit events

signals by Dark Matter particles do not belong to multiple-hits events, that is:

multiple-hits events = Dark Matter particles events "switched off"

Evidence of annual modulation with proper features as required by the DM annual modulation signature is present in the *single-hit* residuals, while it is absent in the *multiple-hits* residual rate.



This result offers an additional strong support for the presence of Dark Matter particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background


Energy distribution of the modulation amplitudes, S_m , for the total exposure

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

DAMA/NaI (7 years) + DAMA/LIBRA (4 years)

total exposure: 300555 kg×day = 0.82 ton×yr

here $T=2\pi/\omega=1$ yr and $t_0=152.5$ day



A clear modulation is present in the (2-6) keV energy interval, while S_m values compatible with zero are present just above

In fact, the S_m values in the (6-20) keV energy interval have random fluctuations around zero with χ^2 equal to 24.4 for 28 degrees of freedom

Statistical distributions of the modulation amplitudes (S_m)

a) S_m values for each detector, each annual cycle and each considered energy bin (here 0.25 keV)


b) $\langle S_m \rangle$ = mean values over the detectors and the annual cycles for each energy bin; σ = errors associated to each S_m

DAMA/LIBRA (4 years)


total exposure: 0.53 ton \times yr

Each panel refers to each detector separately; 64 entries = 16 energy bins in 2-6 keV energy interval \times 4 DAMA/LIBRA annual cycles

2-6 keV



Standard deviations of the variable
 $(S_m - \langle S_m \rangle) / \sigma$
 for the DAMA/LIBRA detectors



$0.80 < r.m.s. < 1.16$

Individual S_m values follow a normal distribution since $(S_m - \langle S_m \rangle) / \sigma$ is distributed as a Gaussian with a unitary standard deviation (r.m.s.)

→ S_m statistically well distributed in all the detectors and annual cycles

Statistical analyses about modulation amplitudes (S_m)


$$\chi = (S_m - \langle S_m \rangle) / \sigma,$$

$$\chi^2 = \sum \chi^2$$

$\chi^2/d.o.f.$ values of S_m distributions for each DAMA/LIBRA detector in the (2–6) keV energy interval for the four annual cycles.

DAMA/LIBRA (4 years)

total exposure: 0.53 ton \times yr



The $\chi^2/d.o.f.$ values range from 0.7 to 1.28 (64 d.o.f. = 16 energy bins \times 4 annual cycles)

\Rightarrow at 95% C.L. the observed annual modulation effect is well distributed in all the detectors.

- The mean value of the twenty-four points is 1.072, slightly larger than 1. Although this can be still ascribed to statistical fluctuations, let us ascribe it to a possible systematics.
- In this case, one would have an additional error of $\leq 5 \times 10^{-4}$ cpd/kg/keV, if quadratically combined, or $\leq 7 \times 10^{-5}$ cpd/kg/keV, if linearly combined, to the modulation amplitude measured in the (2 – 6) keV energy interval.
- This possible additional error ($\leq 4.7\%$ or $\leq 0.7\%$, respectively, of the DAMA/LIBRA modulation amplitude) can be considered as an upper limit of possible systematic effects

Is there a sinusoidal contribution in the signal?



Phase \neq 152.5 day?

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)] = S_0 + Y_m \cos[\omega(t - t^*)]$$

For Dark Matter signals:

- $|Z_m| \ll |S_m| \approx |Y_m|$
- $\omega = 2\pi/T$
- $t^* \approx t_0 = 152.5d$
- $T = 1 \text{ year}$


Slight differences from 2nd June are expected in case of contributions from non thermalized DM components (as e.g. the SagDEG stream)



E (keV)	S_m (cpd/kg/keV)	Z_m (cpd/kg/keV)	Y_m (cpd/kg/keV)	t^* (day)
2-6	0.0122 ± 0.0016	-0.0019 ± 0.0017	0.0123 ± 0.0016	144.0 ± 7.5
6-14	0.0005 ± 0.0010	0.0011 ± 0.0012	0.0012 ± 0.0011	--

The analysis at energies above 6 keV, the analysis of the multiple-hits events and the statistical considerations about S_m already exclude any sizeable presence of systematical effects

Additional investigations



The analysis at energies above 6 keV, the analysis of the multiple-hits events and the statistical considerations about S_m already exclude any sizeable presence of systematical effects.

Additional investigations on the stability parameters

Modulation amplitudes obtained by fitting the time behaviours of main running parameters, acquired with the production data, when including a DM-like modulation

Running conditions stable
at a level better than 1%

	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
Temperature	$-(0.0001 \pm 0.0061) \text{ }^\circ\text{C}$	$(0.0026 \pm 0.0086) \text{ }^\circ\text{C}$	$(0.001 \pm 0.015) \text{ }^\circ\text{C}$	$(0.0004 \pm 0.0047) \text{ }^\circ\text{C}$
Flux N_2	$(0.13 \pm 0.22) \text{ l/h}$	$(0.10 \pm 0.25) \text{ l/h}$	$-(0.07 \pm 0.18) \text{ l/h}$	$-(0.05 \pm 0.24) \text{ l/h}$
Pressure	$(0.015 \pm 0.030) \text{ mbar}$	$-(0.013 \pm 0.025) \text{ mbar}$	$(0.022 \pm 0.027) \text{ mbar}$	$(0.0018 \pm 0.0074) \text{ mbar}$
Radon	$-(0.029 \pm 0.029) \text{ Bq/m}^3$	$-(0.030 \pm 0.027) \text{ Bq/m}^3$	$(0.015 \pm 0.029) \text{ Bq/m}^3$	$-(0.052 \pm 0.039) \text{ Bq/m}^3$
Hardware rate above single photoelectron	$-(0.20 \pm 0.18) \times 10^{-2} \text{ Hz}$	$(0.09 \pm 0.17) \times 10^{-2} \text{ Hz}$	$-(0.03 \pm 0.20) \times 10^{-2} \text{ Hz}$	$(0.15 \pm 0.15) \times 10^{-2} \text{ Hz}$


**All the measured amplitudes well compatible with zero
+none can account for the observed effect**

(to mimic such signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also simultaneously satisfy all the 6 requirements)


Summary of the results obtained in the additional investigations of possible systematics or side reactions

EPJC 56(2008)333

<i>Source</i>	<i>Main comment</i>	<i>Cautious upper limit (90%C.L.)</i>
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5 \times 10^{-6}$ cpd/kg/keV
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
NOISE	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
ENERGY SCALE	Routine + intrinsic calibrations	$<1-2 \times 10^{-4}$ cpd/kg/keV
EFFICIENCIES	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
SIDE REACTIONS	Muon flux variation measured by MACRO	$<3 \times 10^{-5}$ cpd/kg/keV




+ even if larger they cannot satisfy all the requirements of annual modulation signature



Thus, they can not mimic the observed annual modulation effect

The positive and model independent result by DAMA/NaI + DAMA/LIBRA




- Presence of modulation for 11 annual cycles at $\sim 8.2\sigma$ C.L. with the proper distinctive features of the DM signature; all the features satisfied by the data over 11 independent experiments of 1 year each one
- Absence of known sources of possible systematics and side processes able to quantitatively account for the observed modulation amplitude and to contemporaneously satisfy all the peculiarities of the signature

No other experiment whose result can be directly compared in model independent way is available so far




Corollary quests for candidates




- Just to offer some naive feeling on the complexity of the argument:
experimental S_m values vs expected behaviours

for some DM candidates in few of the many possible astrophysical, nuclear and particle physics scenarios and parameters values




Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



WIMP DM candidate (as in [4])
considering elastic scattering on
nuclei

SI dominant coupling
 $v_0 = 170$ km/s




About the same C.L.

...scaling from NaI


channeling contribution as
in EPJC53(2008)205
considered for curve *b*

Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm ³)	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)
<i>a</i>	A5 (NFW)	0.2	A	15 GeV	3.1×10^{-4}
<i>b</i>	A5 (NFW)	0.2	A	15 GeV	1.3×10^{-5}
<i>c</i>	A5 (NFW)	0.2	B	60 GeV	5.5×10^{-6}
<i>d</i>	B3 (Evans power law)	0.17	B	100 GeV	6.5×10^{-6}
<i>e</i>	B3 (Evans power law)	0.17	A	120 GeV	1.3×10^{-5}

Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



WIMP DM candidate (as in [4])
 Elastic scattering on nuclei
 SI & SD mixed coupling
 $v_0 = 170$ km/s




About the same C.L.

...scaling from NaI



$\theta = 2.435$

Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm ³)	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)	$\xi\sigma_{SD}$ (pb)
<i>f</i>	A5 (NFW)	0.2	A	15 GeV	10^{-7}	2.6
<i>g</i>	A5 (NFW)	0.2	A	15 GeV	1.4×10^{-4}	1.4
<i>h</i>	A5 (NFW)	0.2	B	60 GeV	10^{-7}	1.4
<i>i</i>	A5 (NFW)	0.2	B	60 GeV	8.7×10^{-6}	8.7×10^{-2}
<i>j</i>	B3 (Evans power law)	0.17	A	100 GeV	10^{-7}	1.7
<i>k</i>	B3 (Evans power law)	0.17	A	100 GeV	1.1×10^{-5}	0.11



Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



LDM candidate
 (as in arXiv:0802.4336):
 inelastic interaction
 with electron or nucleus targets



Light bosonic candidate
 (as in IJMPA21(2006)1445):
 axion-like particles totally
 absorbed by target material



About the same C.L.

(NFW) halo model as in [4, 34], local density = 0.17 GeV/cm³, local velocity = 170 km/s

Curve label	DM particle	Interaction	Set as in [4]	m_H	Δ	Cross section (pb)
<i>l</i>	LDM	coherent on nuclei	A	30 MeV	18 MeV	$\xi\sigma_m^{coh} = 1.8 \times 10^{-6}$
<i>m</i>	LDM	coherent on nuclei	A	100 MeV	55 MeV	$\xi\sigma_m^{coh} = 2.8 \times 10^{-6}$
<i>n</i>	LDM	incoherent on nuclei	A	30 MeV	3 MeV	$\xi\sigma_m^{inc} = 2.2 \times 10^{-2}$
<i>o</i>	LDM	incoherent on nuclei	A	100 MeV	55 MeV	$\xi\sigma_m^{inc} = 4.6 \times 10^{-2}$
<i>p</i>	LDM	coherent on nuclei	A	28 MeV	28 MeV	$\xi\sigma_m^{coh} = 1.6 \times 10^{-6}$
<i>q</i>	LDM	incoherent on nuclei	A	88 MeV	88 MeV	$\xi\sigma_m^{inc} = 4.1 \times 10^{-2}$
<i>r</i>	LDM	on electrons	-	60 keV	60 keV	$\xi\sigma_m^e = 0.3 \times 10^{-6}$

curve *r*: also pseudoscalar axion-like candidates (e.g. majoron)
 $m_a=3.2$ keV $g_{aee}=3.9 \cdot 10^{-11}$

Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates (in several of the many astrophysical, nuclear and particle physics scenarios); other ones are open

Neutralino as LSP in SUSY theories

Various kinds of WIMP candidates with several different kind of interactions
Pure SI, pure SD, mixed + Migdal effect + channeling, ... (from low to high mass)

a heavy ν of the 4-th family

Pseudoscalar, scalar or mixed light bosons with axion-like interactions

WIMP with preferred inelastic scattering

Mirror Dark Matter

Light Dark Matter

Dark Matter (including some scenarios for WIMP) electron-interacting

Sterile neutrino


Self interacting Dark Matter

Elementary Black holes such as the Daemons

heavy exotic candidates, as "4th family atoms", ...

... and more

Kaluza Klein particles



Possible model dependent positive hints from indirect searches not in conflict with DAMA results
(but interpretation, evidence itself, derived mass and cross sections depend e.g. on bckg modeling, on DM spatial velocity distribution in the galactic halo, etc.)

Available results from direct searches using different target materials and approaches do not give any robust conflict

where we are ...

- DAMA/LIBRA over 4 annual cycles (0.53 ton×yr) confirms the results of DAMA/NaI (0.29 ton×yr)
- The cumulative confidence level for the model independent evidence for presence of DM particle in the galactic halo is 8.2σ (total exposure 0.82 ton × yr)




- First upgrading of the experimental set-up in Sept. 2008



- Opening of the shield of DAMA/LIBRA set-up in HP N₂ atmosphere
- Replacement of some PMTs in HP N₂ atmosphere



- Dismounting of the Tektronix TDs and mounting of the new Acqiris TDs and of the new DAQ system with optical read-out



Since Oct. 2008 again in data taking

...and where we are going

- *Continuing the data taking*
- *Updating of corollary analyses in some of the many possible scenarios for DM candidates, interactions, halo models, nuclear/atomic properties, etc. is in progress*
- *Next upgrading: replacement of all the PMTs with higher Q.E. ones*
- *Production of new Q.E. PMTs in progress. Goal: to study if it is possible to lower the energy threshold of the detectors*
- *Analyses/data taking to investigate other rare processes in progress/foreseen*

A possible highly radiopure NaI(Tl) multi-purpose set-up DAMA/1 ton (proposed by DAMA in 1996) is at present at R&D phase