



Contribution ID: 100

Type: Oral

Status and prospects of charged lepton flavor violation searches with the MEG-II experiment

Friday, 7 June 2019 15:08 (30 minutes)

The MEG experiment took data at the Paul Scherrer Institute in the years 2009-2013 and published the most stringent limit on the charged lepton flavor violating decay $\mu \rightarrow e\gamma$: $BR(\mu \rightarrow e\gamma) < 4.2 \times 10^{-13}$ @90% C.L.

The MEG detector has been upgraded in order to reach a sensitivity of 5×10^{-14} , which corresponds to an improvement of one order of magnitude.

The basic idea of MEG-II is to achieve the highest possible sensitivity by making the maximum use (7×10^7 muons/s) of the available muon intensity at PSI with an improved detector, since MEG ran at a reduced intensity (3×10^7 muons/s) in order to keep the background at a manageable level.

The key features of the MEG-II are the increase of the rate capability of all detectors to enable running at the intensity frontier, and to increase the resolutions while maintaining the same detector concept.

A new mass, single volume, high granularity tracker, together with a thinner muon stopping target, leads to better spatial, angular and energy positron resolution.

A new highly segmented timing counter improves positron timing capabilities. The detector acceptance for positrons is increased by more than a factor 2 by diminishing the material between these two detectors. The liquid Xenon calorimeter has new smaller photosensors (VUV-sensitive SiPM) that replace current phototubes and improve in particular photon energy resolution. The results of the 2018 pre-engineering run, the first with all the sub-detectors, and the current schedule will be presented.

Collaboration name

MEGII

Presenter: RENGA, Francesco (ROMA1)

Session Classification: Flavor and Precision Physics

Track Classification: Flavor and Precision Physics