

Hadronic uncertainties in $b \rightarrow sll$ exclusive decays

Marco Ciuchini



- The paradigm: $B \rightarrow K^* \mu\mu$ at low q^2
- Factorizable amplitudes → form factors
- Non-factorizable contributions → the charm loop
 - theoretical estimates & phenomenological approaches



Angular analysis of $B \rightarrow K^* \mu\mu$

$$\frac{d^{(4)}\Gamma}{dq^2 d(\cos\theta_l)d(\cos\theta_k)d\phi} = \frac{9}{32\pi} \left(I_1^s \sin^2\theta_k + I_1^c \cos^2\theta_k + (I_2^s \sin^2\theta_k + I_2^c \cos^2\theta_k) \cos 2\theta_l \right. \\ \left. + I_3 \sin^2\theta_k \sin^2\theta_l \cos 2\phi + I_4 \sin 2\theta_k \sin 2\theta_l \cos\phi \right. \\ \left. + I_5 \sin 2\theta_k \sin\theta_l \cos\phi + (I_6^s \sin^2\theta_k + I_6^c \cos^2\theta_K) \cos\theta_l \right. \\ \left. + I_7 \sin 2\theta_k \sin\theta_l \sin\phi + I_8 \sin 2\theta_k \sin 2\theta_l \sin\phi \right. \\ \left. + I_9 \sin^2\theta_k \sin^2\theta_l \sin 2\phi \right)$$

angular
analysis

$$S_i = \left(I_i^{(s,c)} + \bar{I}_i^{(s,c)} \right) / \Gamma' \\ (2\Gamma' \equiv d\Gamma/dq^2 + d\bar{\Gamma}/dq^2)$$

8 CP-AVERAGED OBSERVABLES

$$F_L, A_{FB}, S_{3,4,5,7,8,9}$$

In the helicity amplitude formalism ($m_\ell \sim 0$),
we need to compute few helicity amplitudes:

$$H_{V,A}^\lambda \quad \lambda = 0, \pm$$

$$I_1^c = -I_2^c = \frac{F}{2} (|H_V^0|^2 + |H_A^0|^2),$$

$$I_6^s = F \text{Re}[H_V^-(H_A^-)^* - H_V^+(H_A^+)^*],$$

$$I_1^s = 3I_2^s = \frac{3}{8}F (|H_V^+|^2 + |H_V^-|^2 + |H_A^+|^2 + |H_A^-|^2), \quad I_6^c = 0,$$

$$I_7 = \frac{F}{2} \text{Im} [(H_A^+ + H_A^-)(H_V^0)^* + (H_V^+ + H_V^-)(H_A^0)^*],$$

$$I_3 = -\frac{F}{2} \text{Re} [H_V^+(H_V^-)^* + H_A^+(H_A^-)^*], \quad I_8 = \frac{F}{4} \text{Im} [(H_V^- - H_V^+)(H_V^0)^* + (H_A^- - H_A^+)(H_A^0)^*],$$

$$I_4 = \frac{F}{4} \text{Re} [(H_V^+ + H_V^-)(H_V^0)^* + (H_A^+ + H_A^-)(H_A^0)^*], \quad I_9 = \frac{F}{4} \text{Im} [H_V^+(H_V^-)^* + H_A^+(H_A^-)^*].$$


$$H_V^\lambda = \frac{4iG_F m_B}{\sqrt{2}} \frac{\alpha_e}{4\pi} \lambda_t \left\{ C_9^{\text{eff}} \tilde{V}_{L\lambda} + \frac{m_B^2}{q^2} \left[\frac{2m_b}{m_B} C_7^{\text{eff}} \tilde{T}_{L\lambda} - 16\pi^2 h_\lambda \right] \right\},$$

$$H_A^\lambda = \frac{4iG_F m_B}{\sqrt{2}} \frac{\alpha_e}{4\pi} \lambda_t C_{10} \tilde{V}_{L\lambda}. \quad \lambda = 0, \pm$$

NNLO Wilson coefficients from the $\Delta B=1$, $\Delta S=1$ effective Hamiltonian:

$$\mathcal{H}_{\text{eff}}^{\Delta B=1} = \mathcal{H}_{\text{eff}}^{sl+\gamma} + \mathcal{H}_{\text{eff}}^{\text{had}}$$

$$\mathcal{H}_{\text{eff}}^{sl+\gamma} = -\frac{4G_F}{\sqrt{2}} \lambda_t (C_7 Q_{7\gamma} + C_9 Q_{9V} + C_{10} Q_{10A})$$



$$Q_{7\gamma} = \frac{e}{16\pi^2} m_b \bar{s}_L \sigma_{\mu\nu} F^{\mu\nu} b_R,$$

$$Q_{9V} = \frac{\alpha_e}{4\pi} (\bar{s}_L \gamma_\mu b_L) (\bar{\ell} \gamma^\mu \ell),$$

$$Q_{10A} = \frac{\alpha_e}{4\pi} (\bar{s}_L \gamma_\mu b_L) (\bar{\ell} \gamma^\mu \gamma^5 \ell).$$

Hadronic matrix elements
of quark currents:
FORM FACTORS

Form factors

Six (seven) form factors need to be computed: $\tilde{V}_{L\lambda}, \tilde{T}_{L\lambda}, (\tilde{S}_L)$
or, in the transversity basis, $V, A_{(0),1,2}, T_{0,1,2}$

Two main approaches:

Heavy quark symmetry

Isgur, Wise; J. Charles et al., hep-ph/9812358;
Grinstein, Pirjol, hep-ph/0201298

- 7 FF's \rightarrow 2 soft functions in the infinite mass limit
- useful to define optimized observables (FF-independent for $m_b \rightarrow \infty$)
- symmetry breaking corrections still require a dynamical approach

Non-perturbative QCD approaches

- 2+1 lattice QCD calculations at low recoil - uncertainty $\sim 5\text{-}10\%$

Horgan, Liu, Meinel, Wingate, arXiv:1310.3722

- light-cone sum rules at large recoil - uncertainty $\sim 10\text{-}15\%$

Bharucha, Straub, Zwicky, arXiv:1503.05534


Results given in terms of the coefficients of a z-expansion

$$F^{(i)}(q^2) = \sum_k \alpha_k^{(i)} \frac{(z(q^2) - z(0))^k}{1 - q^2/m_{R,i}^2}$$



Correlations induced by the HQ symmetry accounted for by the provided correlation matrix

In the low q^2 region, one has to rely on LCSR results, yet the extrapolation to high q^2 matches quite well lattice QCD

Bharucha, Straub, Zwicky, arXiv:1503.05534v3



Charm-loop effects



Charm loop in the effective theory


$$\mathcal{H}_{\text{eff}}^{\text{had}} = \frac{4G_F}{\sqrt{2}} \left\{ \lambda_u [C_1 (Q_1^u - Q_1^c) + C_2 (Q_2^u - Q_2^c)] - \lambda_t \left[C_1 Q_1^c + C_2 Q_2^c + \sum_{i=3}^6 C_i Q_i + C_8 Q_{8g} \right] \right\}$$

top loops in the SM give rise to penguin operators

- non-perturbative matrix elements of local operators
- α_s suppressed matching conditions, small Wilson coefficients

charm (and up) loops appear as Wick contractions in the MEs

- dominated by the insertion of $Q_{1,2}$, namely $O(1)$ Wilson coefficients
- easily produce intermediate real states, i.e. rescattering, non-local contributions, strong phases, etc.



Non-leptonic $b \rightarrow s$ decays: “charming” penguins

Colangelo, Nardulli, Paver, Riazuddin, Z.Phys. C45 (1990) 575
MC, Franco, Martinelli, Pierini, Silvestrini, hep-ph/9703353

Charm penguin is doubly Cabibbo-enhanced in $b \rightarrow s \bar{u} u$ transitions

- Threatened factorization of non-leptonic B decays in the infinite mass limit → tamed
Beneke, Buchalla, Neubert, Sachrajda, arXiv:0902.4446
see also Bauer, Pirjol, Rothstein, Stewart, hep-ph/0502094
- Questioned the extraction of the CKM angle γ from $B \rightarrow K\pi$ decays (but helped to account for the $K\pi$ and $\pi\pi$ BRs and CP asymmetries)
- Challenged NP sensitivity of various non-leptonic B decays
→ still to be tamed
MC, Franco, Martinelli, Pierini, Silvestrini, hep-ph/0208048
Beneke, Buchalla, Neubert, Sachrajda, hep-ph/0104110
Fleischer, Matias, hep-ph/9906274, ...

BUT 4-quark operators also contribute to the ME. In the helicity amplitude formalism, they appear in

Jäger, Camalich, arXiv:1212.2263;
Melikhov, Nikitin, Simula, hep-ph/9807464


$$H_V^\lambda = \frac{4iG_F m_B}{\sqrt{2}} \frac{e^2}{16\pi^2} \lambda_t \left\{ C_9^{\text{eff}} \tilde{V}_{L\lambda} + \frac{m_B^2}{q^2} \left[\frac{2m_b}{m_B} C_7^{\text{eff}} \tilde{T}_{L\lambda} - 16\pi^2 h_\lambda \right] \right\}$$

$$h_\lambda(q^2) = \frac{\epsilon_\mu^*(\lambda)}{m_B^2} \int d^4x e^{iqx} \langle \bar{K}^* | T\{ j_{\text{em}}^\mu(x) \mathcal{H}_{\text{eff}}^{\text{had}}(0) \} | \bar{B} \rangle$$

- At small values of $q^2 = m_{\ell\ell}^2$, the H^{had} matrix element factorizes in the infinite mass limit

Beneke, Feldmann, Seidel,
hep-ph/0106067

- Yet the “charming penguin” issues are present:
 - how large is the genuine power-suppressed contribution?
 - how much does it increase approaching the resonant region where factorization badly fails?



Taming the charm-loop monster...




An estimate in 2 steps:

Khodjamirian, Mannel, Pivovarov, Wang, arXiv:1006.4945

1. at $q^2 \ll 4m_c^2$ the charm loop is dominated by light-cone dynamics.

One can write the ME $\left[\mathcal{H}_\mu^{(B \rightarrow K^{(*)})}(p, q) \right]_{non\ fact} = 2C_1 \langle K^{(*)}(p) | \tilde{\mathcal{O}}_\mu(q) | B(p+q) \rangle,$

where $\tilde{\mathcal{O}}_\mu(q) = \int d\omega I_{\mu\rho\alpha\beta}(q, \omega) \bar{s}_L \gamma^\rho \delta[\omega - \frac{(in + \mathcal{D})}{2}] \tilde{G}_{\alpha\beta} b_L$ is a non-local operator representing the first subleading term of an expansion in $\Lambda^2/(4m_c^2 - q^2)$ (single soft gluon approximation), whose ME is computed using light-cone sum rules



step 1 → estimate of the hadronic contribution at small $q^2 < \text{few GeV}^2$
but large uncertainties (100%? more?)
no hard gluons, no phases, no scale and scheme dependence, ...

2. extend the previous result to larger q^2 using a dispersion relation, modeling the spectral function (2 physical $\Psi^{(i)}$ + effective poles)



step 2

$$\Delta C_{9,i}^{(c\bar{c})}(q^2) = \frac{r_{1,i} \left(1 - \frac{\bar{q}^2}{q^2}\right) + \Delta C_{9,i}^{(c\bar{c})}(\bar{q}^2) \frac{\bar{q}^2}{q^2}}{1 + r_{2,i} \frac{\bar{q}^2 - q^2}{m_{J/\psi}^2}}$$

	$r_{1,i}$	$r_{2,i}$
	$0.10^{+0.02}_{-0.00}$	$1.13^{+0.00}_{-0.01}$
	$0.09^{+0.01}_{-0.00}$	$1.12^{+0.00}_{-0.01}$
	$0.06^{+0.04}_{-0.10}$	$1.05^{+0.05}_{-0.04}$

but model dependence, no pert. gluons and phases: uncertainty ?

Khodjamirian, Mannel, Pivovarov, Wang, arXiv:1006.4945



2010 → today

Step 1: no new non-perturbative calculation. However an hierarchy among contributions in the helicity basis has been found

$$h_+ \sim \mathcal{O} \left(\frac{\Lambda}{m_b} \right) h_-$$

Jäger, Camalich, arXiv:1212.2263

Step 2: recent attempts to gain more control over the q^2 dependence improving the dispersion relation approach

1. empirical model using resonance data over the full dimuon spectrum

Blake, Egede, Owen, Pomery, Petridis, arXiv:1709.03921


2. replace the dispersion relation with a z-expansion of h_λ , constraining the coefficients using analiticity and

Bobeth, Chrzaszcz, van Dyk, Virto, arXiv:1707.07305

1. resonant $B \rightarrow \Psi^{(n)} K^*$ data (masses and amplitudes)
2. LCSR + QCDF theoretical results at small/negative q^2

empirical model

Blake, Egede, Owen, Pomery, Petridis, arXiv:1709.03921
see also LHCb collaboration, arXiv:1612.06764



The hadronic contribution is modeled as the sum of 1^- resonances represented by relativistic Breit-Wigner functions


$$\Delta C_{9,\lambda}^{\text{had}}(q^2) = \sum_j \eta_j^\lambda e^{i\theta_j^\lambda} A_j^{\text{res}}(q^2)$$

$$A_j^{\text{res}}(q^2) = \frac{m_{\text{res},j} \Gamma_{\text{res},j}}{(m_{\text{res},j}^2 - q^2) - im_{\text{res},j} \Gamma_j(q^2)}$$

Open issues:

Why should it work far from the resonances? What about double counting? How large is the model uncertainty?

Illustrate nicely the importance of strong phases



c loop from analyticity

Bobeth, Chrzaszcz, van Dyk, Virto, arXiv:1707.07305

Features:


- get rid of DD branch cut modeling by mapping it at the boundary of the expansion region
- exploits the $\psi^{(n)}$ resonance data to constrain the expansion

Open issues:

- strong phases related to the DD_s cut in p^2 are taken from LCSR and QCDF calculations. Are they reliable?

k	0	1	2
$\text{Re}[\alpha_k^{(\perp)}]$	-0.06 ± 0.21	-6.77 ± 0.27	18.96 ± 0.59
$\text{Re}[\alpha_k^{(\parallel)}]$	-0.35 ± 0.62	-3.13 ± 0.41	12.20 ± 1.34
$\text{Re}[\alpha_k^{(0)}]$	0.05 ± 1.52	17.26 ± 1.64	—
$\text{Im}[\alpha_k^{(\perp)}]$	-0.21 ± 2.25	1.17 ± 3.58	-0.08 ± 2.24
$\text{Im}[\alpha_k^{(\parallel)}]$	-0.04 ± 3.67	-2.14 ± 2.46	6.03 ± 2.50
$\text{Im}[\alpha_k^{(0)}]$	-0.05 ± 4.99	4.29 ± 3.14	—

- z expansion: no sign of convergence for the typical values $|z| \sim 0.2\text{-}0.4$
NB: z expansion of FF at much smaller values



Parametrizing the charm loop

Jäger, Camalich, arXiv:1212.2263


MC, Fedele, Franco, Mishima, Paul, Silvestrini, Valli, arXiv:1512.07157
+ preliminary update

$$\begin{aligned} H_V^- &= -iN \left\{ (C_9^{\text{eff}} + h_-^1) V_{L-} + \frac{m_B^2}{q^2} \left[\frac{2m_b}{m_B} (C_7^{\text{eff}} + h_-^0) T_{L-} - 16\pi^2 h_-^2 q^4 \right] \right\} \\ H_V^0 &= -iN \left\{ (C_9^{\text{eff}} + h_-^1) \tilde{V}_{L0} + \frac{m_B^2}{q^2} \left[\frac{2m_b}{m_B} (C_7^{\text{eff}} + h_-^0) \tilde{T}_{L0} - 16\pi^2 (\tilde{h}_0^0 + \tilde{h}_0^1 q^2) \right] \right\} \\ H_V^+ &= -iN \left\{ (C_9^{\text{eff}} + h_-^1) V_{L+} + \frac{m_B^2}{q^2} \left[\frac{2m_b}{m_B} (C_7^{\text{eff}} + h_-^0) T_{L+} - 16\pi^2 (h_+^0 + h_+^1 q^2 + h_+^2 q^4) \right] \right\} \end{aligned}$$

$\Delta C_7^{(cc)} = h_-^0$ and $\Delta C_9^{(cc)} = h_-^1$ shift the corresponding Wilson coefficients (as NP contributions do), while the other parameters have no short-distance counterparts


Fitting hadronic parameters

- Compute all amplitudes using QCD factorization and form factors from LQCD (Bailey et al. '15) and LCSR (Bharucha, Straub & Zwicky '15)
- add hadronic parameters and
 - use LCSR calculation from KMPW at low q^2 (0 and 1 GeV 2) only (PDD)
or
 - extrapolate LCSR calculation to larger q^2 using KMPW (PMD)
- fit all available experimental data using the **HEPfit** code
- compare different models using $IC = -2\overline{\log L} + 4\sigma_{\log L}^2$




SM fit of the h_λ 's

B \rightarrow K $^*\ell\ell$ data accounted for by the hadronic contributions



MC, Coutinho, Fedele, Franco, Paul,
Silvestrini, Valli, in preparation



SM fit of the h_λ 's

B \rightarrow K $^*\ell\ell$ data accounted for by
the hadronic contributions


$|\Delta C_7|$ fixed by the KMPW value
at $q^2 = 0$

No clear evidence for other
non-vanishing hadronic parameters
but interesting correlation


$|\Delta C_9| - h_-^{(2)}$

Future experimental uncertainties
will be able to pin down the $h^{(2)}$'s

MC, Coutinho, Fedele, Franco, Paul,
Silvestrini, Valli, in preparation



preliminary



PDD

SM fit projection

Central values fixed to the present fit global mode

Experimental errors reduced by a factor of 6 (202?)

Many coefficient can be measured, even in the presence of NP (barring ΔC_7 and ΔC_9)

MC, Coutinho, Fedele, Franco, Paul, Silvestrini, Valli, in preparation


NP in $C_{9,\mu}$ and $C_{9,e}$

MC, Coutinho, Fedele, Franco, Paul, Silvestrini, Valli, arXiv:1704.05447 + work in progress


LFUV hints are not affected by hadronic parameters...

$$C_{9,\pm} = \frac{1}{2} (C_{9,\mu}^{\text{NP}} \pm C_{9,e}^{\text{NP}})$$

PMD



PDD





NP in C_{10}^{μ} and C_{10}^e

MC, Coutinho, Fedele, Franco, Paul, Silvestrini, Valli, arXiv:1704.05447 + work in progress

... BUT viable NP scenarios ARE!

$$C_{10,\pm} = \frac{1}{2} (C_{10,\mu}^{\text{NP}} \pm C_{10,e}^{\text{NP}})$$



Summary & Conclusions

The case of $B \rightarrow K^* \ell \bar{\ell}$ was discussed, but similar issues are present in other exclusive $b \rightarrow s$ semileptonic decays in the low q^2 region, albeit to different extents

Two dominant sources of theoretical uncertainties:
form factors and charm loop effect

Form factors uncertainty is in the 10-15% ballpark, but optimized observables have reduced sensitivity

- (-) only LCSR results for FFs available in the low q^2 region
 - (+) full correlation matrix available
- (+) matching between extrapolated LCSR and LQCD results

Charm loop: dangerous or harmless?



A clear-cut non-perturbative calculation is not available yet


Combinations of QCDF, LCSR, analiticity and unitarity point to a moderate effect with a flat q^2 dependence in the region of interest. Yet their ability to fully describe c-loop rescattering is questionable

Future data could be able to pin down hadronic contributions with no short-distance counterparts (all but ΔC_7 and ΔC_9)


LFUV signals are not affected, but their interpretation may be



Backup

Belle data [S. Wehle @ Belle Col.]



$IC = 103$





$$\frac{d^{(4)}\Gamma}{dq^2 d(\cos \theta_l) d(\cos \theta_k) d\phi} = \frac{9}{32\pi} \left(I_1^s \sin^2 \theta_k + I_1^c \cos^2 \theta_k + (I_2^s \sin^2 \theta_k + I_2^c \cos^2 \theta_k) \cos 2\theta_l \right. \\ \left. + I_3 \sin^2 \theta_k \sin^2 \theta_l \cos 2\phi + I_4 \sin 2\theta_k \sin 2\theta_l \cos \phi \right. \\ \left. + I_5 \sin 2\theta_k \sin \theta_l \cos \phi + (I_6^s \sin^2 \theta_k + I_6^c \cos^2 \theta_K) \cos \theta_l \right. \\ \left. + I_7 \sin 2\theta_k \sin \theta_l \sin \phi + I_8 \sin 2\theta_k \sin 2\theta_l \sin \phi \right. \\ \left. + I_9 \sin^2 \theta_k \sin^2 \theta_l \sin 2\phi \right)$$

angular analysis

$$S_i = \left(I_i^{(s,c)} + \bar{I}_i^{(s,c)} \right) / \Gamma' \\ (2\Gamma' \equiv d\Gamma/dq^2 + d\bar{\Gamma}/dq^2)$$

8 CP-AVERAGED OBSERVABLES



$$F_L, A_{FB}, S_{3,4,5,7,8,9}$$

In the helicity amplitude formalism: ($m_\ell \sim 0$)



$$\begin{aligned} I_1^c &= -I_2^c = \frac{F}{2} (|H_V^0|^2 + |H_A^0|^2), & I_6^s &= F \text{Re}[H_V^- (H_A^-)^* - H_V^+ (H_A^+)^*], \\ I_1^s &= 3I_2^s = \frac{3}{8}F (|H_V^+|^2 + |H_V^-|^2 + |H_A^+|^2 + |H_A^-|^2), & I_6^c &= 0, \\ I_3 &= -\frac{F}{2} \text{Re}[H_V^+ (H_V^-)^* + H_A^+ (H_A^-)^*], & I_7 &= \frac{F}{2} \text{Im}[(H_A^+ + H_A^-)(H_V^0)^* + (H_V^+ + H_V^-)(H_A^0)^*], \\ I_4 &= \frac{F}{4} \text{Re}[(H_V^+ + H_V^-)(H_V^0)^* + (H_A^+ + H_A^-)(H_A^0)^*], & I_8 &= \frac{F}{4} \text{Im}[(H_V^- - H_V^+)(H_V^0)^* + (H_A^- - H_A^+)(H_A^0)^*], \\ I_5 &= \frac{F}{4} \text{Re}[(H_V^- - H_V^+)(H_A^0)^* + (H_A^- - H_A^+)(H_V^0)^*], & I_9 &= \frac{F}{4} \text{Im}[H_V^+ (H_V^-)^* + H_A^+ (H_A^-)^*]. \end{aligned}$$

We need to compute few helicity amplitudes: $H_{V,A}^\lambda$ $\lambda = 0, \pm$



NP IN C_9^μ AND C_9^e



NP IN C_9^μ AND C_{10}^μ



NP IN $C_7, C_{9,10}^{e,\mu}$




FUTURE PROJECTIONS



- Choose a theory setup (SM or NP; rising or non-rising charm loop)
- Generate experimental results from current best fit point in the given setup
- Assume future exp errors scaling LHCb statistical errors to 50/fb (roughly /6) and including BelleII estimates
- Fit parameters from generated data

See also Hurth, Mahmoudi, Martinez Santos & Neshatpour '17;
Albrecht, Bernlochner, Kenzie, Reichert, Straub, Tully '17



SM PROJECTION





NP PROJECTION





NP PROJECTION





NP IN C_9^μ AND C_9^e



NP IN C_9^μ AND C_9^e



NP IN C_9^μ AND C_9^e



NP IN C_7 , C_{10}^μ AND C_{10}^e

