

CP Violation in B decays

Keri Vos

Universität Siegen



Theor. Physik 1



DFG FOR 1873

Motivation

- Studies of CP violation are an important part of the flavour program
 - Determining precisely SM inputs (CKM parameters)
 - Search for new physics through sensitivity for new CP violating phases
- Non-leptonic B decays are key players
 - Large data sets from B-factories and LHCb-run I, many observables
 - Already impressive experimental uncertainties
- Foresee unprecedented precision for LHCb upgrade and Belle II
 - Challenges theorists to keep up


→ see also talk by Gilberto T-X
→ see also talk by Stefan de Boer

Focus on recent progress and very selected topics

$B_q - \bar{B}_q$ mixing observables

Leading contribution in the SM

see Buras, Buchalla, Lautenbacher [1995]



Mass eigenstates H and L arise from diagonalization of $\Delta F = 2$ Hamiltonian


Mass difference $\Delta M_q \equiv M_H^q - M_L^q \sim 2|M_{12}^q| > 0$

- Governed by short-distance contributions
- **New Physics** can have a significant impact see also: di Lucio, Kirk, Lenz [2018]

$B_q - \bar{B}_q$ mixing observables

Leading contribution in the SM

see Buras, Buchalla, Lautenbacher [1995]



Mass eigenstates H and L arise from diagonalization of $\Delta F = 2$ Hamiltonian

Mass difference $\Delta M_q \equiv M_H^q - M_L^q \sim 2|M_{12}^q| > 0$

- Governed by short-distance contributions
- **New Physics** can have a significant impact see also: di Lucio, Kirk, Lenz [2018]



Width difference $\Delta\Gamma_q \equiv \Gamma_L^q - \Gamma_H^q \sim 2\Gamma_{12}^q \cos\phi_q$

- $\Delta\Gamma_s$ sizeable Dunietz, Fleischer, Nierste [2001]; Lenz et al. [2012]; Badin, Gabbiani, Petrov [2007]
- Dominated by tree decays, rather insensitive to New Physics

$B_q - \bar{B}_q$ mixing observables

Leading contribution in the SM

see Buras, Buchalla, Lautenbacher [1995]



Mass eigenstates H and L arise from diagonalization of $\Delta F = 2$ Hamiltonian

Mass difference $\Delta M_q \equiv M_H^q - M_L^q \sim 2|M_{12}^q| > 0$

- Governed by short-distance contributions
- **New Physics** can have a significant impact see also: di Lucio, Kirk, Lenz [2018]

Width difference $\Delta\Gamma_q \equiv \Gamma_L^q - \Gamma_H^q \sim 2\Gamma_{12}^q \cos\phi_q$

- $\Delta\Gamma_s$ sizeable Dunietz, Fleischer, Nierste [2001]; Lenz et al. [2012]; Badin, Gabbiani, Petrov [2007]
- Dominated by tree decays, rather insensitive to New Physics


CP-violating mixing phase $\phi_q \equiv \arg(-M_{12}^q/\Gamma_{12}^q)$

CP violation in $B_q-\bar{B}_q$ mixing

Flavour-specific semi-leptonic decays probe CP violation in **mixing**

$$a_{\text{sl}}^q = \frac{\Gamma(\bar{B}_q(t) \rightarrow f) - \Gamma(B_q(t) \rightarrow \bar{f})}{\Gamma(\bar{B}_q(t) \rightarrow f) + \Gamma(B_q(t) \rightarrow \bar{f})} = \left(\frac{\Delta\Gamma_q}{\Delta M_q} \right) \tan \phi_q$$

Current Status:



Artuso, Borissov, Lenz [2015]

Inclusive SM prediction using HQE

$$a_{\text{sl}}^d|_{\text{SM}} = (-4.7 \pm 0.6) \times 10^{-4}$$

$$a_{\text{sl}}^s|_{\text{SM}} = (2.22 \pm 0.27) \times 10^{-5}$$

see: Jubb, Kirk, Lenz, Tetlalmatzi-Xolocotzi [2017]

- Assumes quark-hadron duality
- Requires lattice calculations of higher-dimensional matrix elements
- Sensitivity to CP violating New Physics

CP violation in $B_s^0-\bar{B}_s^0$ mixing

Fleischer, KKV, Phys.Lett. B770 (2017) 319

New physics would also show up in **exclusive determinations** of ϕ_s

$$a_{sl}^s = \left(\frac{\Delta\Gamma_s}{\Delta M_s} \right) \times \tan(\phi_s)$$

CP violation in $B_s^0-\bar{B}_s^0$ mixing

Fleischer, KKV, Phys.Lett. B770 (2017) 319

New physics would also show up in **exclusive determinations** of ϕ_s

$$a_{\text{sl}}^s = [(0.46 \pm 0.04) \times 10^{-2}] \times \tan(\langle \phi_s \rangle + \Delta\Psi)$$

- a_{sl}^s already suppressed by measurements of ΔM_s and Γ_s

CP violation in $B_s^0 - \bar{B}_s^0$ mixing

Fleischer, KKV, Phys.Lett. B770 (2017) 319

New physics would also show up in **exclusive determinations** of ϕ_s

$$a_{\text{sl}}^s = [(0.46 \pm 0.04) \times 10^{-2}] \times \tan(\langle \phi_s \rangle + \Delta\Psi)$$

- a_{sl}^s already suppressed by measurements of ΔM_s and Γ_s

Implications of exclusive ϕ_s determinations

- Available determinations are all **consistent** with the SM
- Significantly constrains possible new physics effects
- $\langle \phi_s \rangle$ average of ϕ_s^f with $f = J/\psi\phi, D_s^- D_s^+, J/\psi\pi^+\pi^-, \dots$
- Phase $\Delta\Psi$ determined from **experimental data**

$$\Delta\Psi = \arg \left[\sum_f \eta_f w_f e^{i(\phi_s^f - \langle \phi_s \rangle)} \right] \quad w_f = \Gamma(B_s^0 \rightarrow f) \sqrt{\frac{1 - \mathcal{A}_{\text{CP}}^{\text{dir}}(B_s \rightarrow f)}{1 + \mathcal{A}_{\text{CP}}^{\text{dir}}(B_s \rightarrow f)}} \quad \text{measurable weight function}$$

CP violation in $B_s^0-\bar{B}_s^0$ mixing


Fleischer, KKV, Phys.Lett. B770 (2017) 319

New physics would also show up in **exclusive determinations** of ϕ_s

$$a_{\text{sl}}^s = [(0.46 \pm 0.04) \times 10^{-2}] \times \tan(\langle\phi_s\rangle + \Delta\Psi)$$

- a_{sl}^s already suppressed by measurements of ΔM_s and Γ_s

Exclusive Prediction:



- Limits the room for new physics
- Interesting to confront with more precise measurements
- Opens also new windows to search for CPV in D_s^\pm decays

Limited by $B_s \rightarrow D_s^- D_s^+$ (small band: upgrade scenario)

CP violation in non-leptonic B decays

Non-leptonic B decays

see Buras, Buchalla, Lautenbacher [1995]

Effective Hamiltonian

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} \sum_{j=u,c} V_{jq}^* V_{jb} \left(\sum_{i=1,2} C_i(\mu) \mathcal{O}_i^{jq}(\mu) + \sum_{i=3}^{10} C_i(\mu) \mathcal{O}_i^q \right)$$

- $C_i(\mu)$ real short-distance coefficient, $\langle \mathcal{O}_i \rangle$ long-distance physics

Current-current operators

$$\mathcal{O}_1^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) j_\beta \bar{j}_\beta \gamma_\mu (1 - \gamma_5) b_\alpha$$

$$\mathcal{O}_2^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) j_\alpha \bar{j}_\beta \gamma_\mu (1 - \gamma_5) b_\beta$$

QCD penguin operators ($q' = u, d, s, c, b$)

$$\mathcal{O}_3^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\alpha \sum_{q'} \bar{q}'_\beta \gamma_\mu (1 - \gamma_5) q'_\beta$$

$$\mathcal{O}_4^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\beta \sum_{q'} \bar{q}'_\beta \gamma_\mu (1 - \gamma_5) q'_\alpha$$

$$\mathcal{O}_5^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\alpha \sum_{q'} \bar{q}'_\beta \gamma_\mu (1 + \gamma_5) q'_\beta$$

$$\mathcal{O}_6^q = \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\beta \sum_{q'} \bar{q}'_\beta \gamma_\mu (1 + \gamma_5) q'_\alpha$$

EW penguin operators

$$\mathcal{O}_7^q = \frac{3}{2} \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\alpha \sum_{q'} Q_{q'} \bar{q}'_\beta \gamma_\mu (1 + \gamma_5) q'_\beta$$

$$\mathcal{O}_8^q = \frac{3}{2} \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\beta \sum_{q'} Q_{q'} \bar{q}'_\beta \gamma_\mu (1 + \gamma_5) q'_\alpha$$

$$\mathcal{O}_9^q = \frac{3}{2} \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\alpha \sum_{q'} Q_{q'} \bar{q}'_\beta \gamma_\mu (1 - \gamma_5) q'_\beta$$

$$\mathcal{O}_{10}^q = \frac{3}{2} \bar{q}_\alpha \gamma^\mu (1 - \gamma_5) b_\beta \sum_{q'} Q_{q'} \bar{q}'_\beta \gamma_\mu (1 - \gamma_5) q'_\alpha$$

Non-leptonic B decays

see Buras, Buchalla, Lautenbacher [1995]

Effective Hamiltonian

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} \sum_{j=u,c} V_{jq}^* V_{jb} \left(\sum_{i=1,2} C_i(\mu) \mathcal{O}_i^{jq}(\mu) + \sum_{i=3}^{10} C_i(\mu) \mathcal{O}_i^q \right)$$

- $C_i(\mu)$ real short-distance coefficient, $\langle \mathcal{O}_i \rangle$ long-distance physics

General non-leptonic B decay (CKM unitarity implies: at most two independent CKM amplitudes)

$$A(B \rightarrow f) = e^{i\varphi_1} |A_1| e^{i\theta_1} + e^{i\varphi_2} |A_2| e^{i\theta_2}$$
$$|A_i| e^{i\delta_i} = \sum_k C_k(\mu) \times \langle f | \mathcal{O}_k^i(\mu) | B \rangle$$

Perturbatively calculable
Hadronic matrix element

Hadronic matrix elements theoretically challenging

Continuum methods to determine Hadronic Matrix Elements

- pQCD

Li, Yu [1995]; Li, Yang [1999]; Keum, Li, Sanda [2000]

- QCD Factorization

[Beneke, Buchalla, Neubert, Sachrajda, Bell, Huber, Feldmann, Li, Jaeger, Zupan, ...]

- Strong phases generated at $\mathcal{O}(\alpha)$
- Completion of penguin parameters at NNLO in progress
- Power corrections challenging to control

Beneke, Bell, Huber [in progress]

- Soft Collinear Effective Theory (SCET)

Bauer, Pirjol, Stewart [2001]; Bauer, Grinstein, Pirjol, Stewart [2003]; ...

- Important tool to establish QCDF at higher orders

Flavour symmetries [Gronau, Rosner, London, Buras, Fleischer, Zupan, Pirjol, Jaeger, Mannel, Jung, ...]

- Allow determination of CKM phases
- Permit valuable insights into non-perturbative effects

CP violation in B decays in the SM

Charged B mesons

$$\begin{aligned} \mathcal{A}_{\text{CP}}^{\text{dir}} &\equiv \frac{|A(B \rightarrow f)|^2 - |A(\bar{B} \rightarrow f)|^2}{|A(B \rightarrow f)|^2 + |A(\bar{B} \rightarrow f)|^2} \\ &= \frac{2|\mathcal{A}_1||\mathcal{A}_2| \sin(\Delta\theta) \sin \Delta\varphi}{|\mathcal{A}_1|^2 + |\mathcal{A}_2|^2 + 2|\mathcal{A}_1||\mathcal{A}_2| \cos(\Delta\theta) \cos \Delta\varphi} \end{aligned}$$

Direct CP asymmetry

- Interference between two different decay amplitudes
- Non-trivial CP-conserving strong phase difference $\Delta\theta$
- Non-trivial CP-violating weak phase difference $\Delta\varphi$ (extraction of CKM angle γ)

CP violation in B decays in the SM

Neutral B_d and B_s mesons

$$\mathcal{A}_{\text{CP}}(t) \equiv \frac{\Gamma(B_q^0(t) \rightarrow f) - \Gamma(\bar{B}_q^0(t) \rightarrow \bar{f})}{\Gamma(B_q^0(t) \rightarrow f) + \Gamma(\bar{B}_q^0(t) \rightarrow \bar{f})} = \frac{\mathcal{A}_{\text{CP}}^{\text{dir}} \cos(\Delta M_q t) + \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta M_q t)}{\cosh(\Delta\Gamma_q t/2) + \mathcal{A}^{\Delta\Gamma_q} \sinh(\Delta\Gamma_q t/2)}$$


$$\mathcal{A}_{\text{CP}}^{\text{dir}} \equiv \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, \quad \mathcal{A}^{\Delta\Gamma} \equiv \frac{-2\text{Re}\lambda_f}{1 + |\lambda_f|^2}, \quad \lambda_f = \frac{q}{p} \frac{\bar{\mathcal{A}}_f}{\mathcal{A}_f}$$

$$\mathcal{A}_{\text{CP}}^{\text{mix}} \equiv \frac{-2\text{Im}\lambda_f}{1 + |\lambda_f|^2} = \frac{2|\lambda_f|}{1 + |\lambda_f|^2} \sin \phi_q$$

Mixing-induced CP asymmetry

- Arises from interference between mixing and decay
- Offers an important additional observable
- Can also be sizeable if only one amplitude dominates

Determination of γ from $B \rightarrow DK$




→ see talk by Matthew Kenzie

Determination of γ from $B \rightarrow DK$


$$\gamma = \arg \left(-\frac{V_{ud} V_{ub}^*}{V_{cd} V_{cb}^*} \right)$$

Gronau, Wyler [1991]; Gronau, London [1991]; Atwood, Dunietz, Soni [1997]

Giri, Grossman, Soffer, Zupan [2003]



$$\propto V_{cb} V_{us(d)}^*$$



$$\propto V_{ub} V_{cs(d)}^*$$

- Important parameter: **key input of the CKM**
- Theoretically extremely clean (no penguin contributions)
 - Electroweak box corrections tiny Brod, Zupan [2013]; Brod [2014]
- Incredible precision of 1° expected at LHCb upgrade
- New physics contributions in $C_{1,2}$ may cause **sizeable shifts** in γ

Brod, Lenz, Wiebusch, Tetlalmatzi-Xolocotzi [2014]

γ determination from $B_s \rightarrow D_s^\pm K^\mp, \dots$

Aleksan, Dunietz, Kayser [1990]; de Bruyn, Fleischer, Kneijens, Merk, Schiller, Tuning [2012]; Fleischer [2003]

Another theoretically clean probe

Time-dependent analysis of $B_s \rightarrow D_s^\pm K^\mp, \dots$ probes $\phi_s + \gamma$

- Most precise measurement of γ from B_s system LHCb JHEP 03 [2018] 059

$$\gamma = (128_{-22}^{+17})^\circ \quad (\text{using } \phi_s \text{ from } b \rightarrow \bar{c}cs)$$

- Great potential for the LHCb upgrade
- Possible to perform a **joint analysis to determine γ and ϕ_s**

Fleischer, Nucl. Phys. B61 (2003) 459

Similarly $B_d \rightarrow D_s^\pm \pi^\mp, \dots$ decays probe $\phi_d + \gamma$

see talk by Greig Cowan

Mixing angles ϕ_s and ϕ_d

Effective mixing angles ϕ_d and ϕ_s

CP asymmetries determine the “effective” mixing angle

$$\sin \phi_q^{\text{eff}} = \frac{\mathcal{A}_{\text{CP}}^{\text{mix}}(B_q^0 \rightarrow f)}{\sqrt{1 - \mathcal{A}_{\text{CP}}^{\text{dir}}(B_q^0 \rightarrow f)^2}} = \sin \left(\phi_q^{\text{SM}} + \Delta\phi_q + \phi_q^{\text{NP}} \right)$$

- New era of precision physics: reach of $\mathcal{O}(0.5^\circ)$ foreseen
- Subleading terms are doubly Cabibbo suppressed
- Controlling hadronic effects crucial
- Penguin shift $\Delta\phi_q$ decay is mode specific

Non-perturbative effects

Frings, Nierste, Wiebush [2015]

- $B \rightarrow J/\psi M$ factorizes in $N_c \rightarrow \infty$, but large corrections
- Flavour symmetries provide valuable insights into hadronic parameters

Fleischer [1999]

Controlling penguin effects in $B_d \rightarrow J/\psi K_S$

Fleischer [1999]; Ciuchini, Pierini, Silvestrini [2005, 2011]

Faller, Fleischer, Jung, Mannel [2008]; Jung [2012]

de Bruyn, Fleischer [2015]

$$\phi_d^{\text{SM}} \equiv 2\beta = 2\arg\left(-\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*}\right)$$

Penguin suppressed golden mode:

$$\mathcal{A}(B_d^0 \rightarrow J/\psi K_S) = \left(1 - \frac{\lambda^2}{2}\right) C' \left[1 + \epsilon a' e^{i\theta'} e^{i\gamma}\right], \quad \epsilon = \frac{\lambda^2}{1 + \lambda^2} \sim 0.05$$


Penguin enhanced control mode:

$$\mathcal{A}(B_s^0 \rightarrow J/\psi K_S) = -\lambda C \left[1 - a e^{i\theta} e^{i\gamma}\right]$$


- Extract penguin parameters (a, θ) using γ as input
- Decays are related via U -spin (s -quark \leftrightarrow d -quark)
- Only sensitive to **non-factorizable** U -spin breaking de Bruyn, Fleischer [2015]

Controlling penguin effects in $B_d \rightarrow J/\psi K_S$

de Bruyn, Fleischer, JHEP 1503 [2015] 145



Current data



Benchmark scenario (LHCb upgrade)

- Current data gives $\Delta\phi_d^{J/\psi K_S} = (-0.71^{+0.56}_{-0.65})^\circ$ Some theoretical assumptions
- Benchmark scenario matches experimental precision in upgrade era

Penguin effects can be controlled!

Controlling penguin effects in $B_s \rightarrow J/\psi\phi$

Fleischer [1999]; Ciuchini, Pierini, Silvestrini [2005, 2011]

Faller, Fleischer, Jung, Mannel [2008]; Jung [2013]

de Bruyn, Fleischer [2015]; Fleischer [2007]; Jung, Schacht [2014]

$$\phi_s^{\text{SM}} \equiv 2\beta_s = 2\arg\left(-\frac{V_{ts}V_{tb}^*}{V_{cs}V_{cb}^*}\right)$$

See Talk by Greig Cowan

Penguin suppressed golden mode:

$B_s^0 \rightarrow J/\psi\phi$ (requires polarization measurements)

Penguin enhanced control mode:

$B_d^0 \rightarrow J/\psi\rho^0$ (but also $B_s \rightarrow J/\psi\bar{K}^{*0}$)

- Implement U -spin symmetry and use γ as input
- CP asymmetries measurements are key inputs
- Already implemented by LHCb LHCb, JHEP 1511 (2015) 082
- Penguin effects under control \rightarrow additional tests of QCD possible

Similar strategy allows extraction of ϕ_s from $B_s \rightarrow D_s\bar{D}_s$

CP violation in $B_s^0 \rightarrow K^- K^+$

Flavor symmetries in $B_s^0 \rightarrow K^- K^+$ and $B_d \rightarrow \pi^- \pi^+$

Fleischer [1999, 2007]; Fleischer, Kneijens [2011]

- $B_s^0 \rightarrow K^- K^+$ dominated by QCD Penguin topologies
- Related to $B_d^0 \rightarrow \pi^- \pi^+$ via *U-spin symmetry*
- Extract γ and ϕ_s from direct and mixing-induced CP asymmetries

Fleischer [1999,2007]; Fleischer, Kneijens [2011]; Cuichini, Franco, Mishima, Silvestrini [2012]; LHCb [2015]

$$\gamma = (63.5_{-6.7}^{+7.2})^\circ \quad \phi_s = -(6.9_{-8.0}^{+9.2})^\circ$$

- Allows comparison between pure tree and penguin determinations
- Quickly limited by dominant *U-spin* breaking corrections

Flavor symmetries in $B_s^0 \rightarrow K^- K^+$ and $B_d \rightarrow \pi^- \pi^+$

Fleischer [1999, 2007]; Fleischer, Kneijens [2011]

- $B_s^0 \rightarrow K^- K^+$ dominated by QCD Penguin topologies
- Related to $B_d^0 \rightarrow \pi^- \pi^+$ via ***U-spin symmetry***
- Extract γ and ϕ_s from direct and mixing-induced CP asymmetries

Fleischer [1999,2007]; Fleischer, Kneijens [2011]; Cuichini, Franco, Mishima, Silvestrini [2012]; LHCb [2015]

$$\gamma = (63.5_{-6.7}^{+7.2})^\circ \quad \phi_s = -(6.9_{-8.0}^{+9.2})^\circ$$

Controlling $SU(3)$ breaking effects

- γ and ϕ_d input parameters; extract ϕ_s
- Split U -spin corrections: factorizable and **non-factorizable** effects
- Semileptonic ratios provide additional input

$$R_K \equiv \frac{\Gamma(B_s \rightarrow K^- K^+)}{|d\Gamma(B_s \rightarrow K^- \ell^+ \nu_\ell)/dq^2|_{q^2=m_K^2}}$$

Controlling $SU(3)$ breaking effects

Gronau, Rosner [1995]; Fleischer, Jaarsma, and KKV[2016]

Non-factorizable U -spin breaking probed by

$$\zeta_{\text{NF}}^a \equiv \left| \frac{a_{\text{NF}}}{a'_{\text{NF}}} \right| = \left| \frac{a_{\text{NF}}^T}{a_{\text{NF}}^{T'}} \right| \left| \frac{1+r_P}{1+r'_P} \right| \left| \frac{1+x}{1+x'} \right|$$

$$r_P \equiv P^{(ut)} / T \sim \mathcal{O}(\lambda)$$

$$x \equiv \frac{E+PA^{(ut)}}{T+P^{(ut)}} \sim \mathcal{O}(\lambda)$$

- Very favourable and robust structure
- Use data-driven methods to quantify U -spin breaking corrections

Controlling $SU(3)$ breaking effects

Gronau, Rosner [1995]; Fleischer, Jaarsma, and KKV[2016]

Non-factorizable U -spin breaking probed by

$$\zeta_{\text{NF}}^a \equiv \left| \frac{a_{\text{NF}}}{a'_{\text{NF}}} \right| = \left| \frac{a_{\text{NF}}^T}{a_{\text{NF}}^{T'}} \right| \left| \frac{1+r_P}{1+r'_P} \right| \left| \frac{1+x}{1+x'} \right|$$

$$r_P \equiv P^{(ut)} / T \sim \mathcal{O}(\lambda)$$

$$x \equiv \frac{E+PA^{(ut)}}{T+P^{(ut)}} \sim \mathcal{O}(\lambda)$$


- Very favourable and robust structure
- Use data-driven methods to quantify U -spin breaking corrections

Hadronic uncertainties

- QCDF probes the tree-level contributions a_{NF}^T Beneke, Huber, Li [2010]
- More insights from future measurements of CP asymmetries
 - Pure penguin (P) $B_d^0 \rightarrow K^0 \bar{K}^0, B_s^0 \rightarrow K^0 \bar{K}^0$
 - Pure exchange (E) and penguin annihilation (PA) topologies $B_d^0 \rightarrow K^+ K^-, B_s^0 \rightarrow \pi^+ \pi^-$

Illustration of the future error on $\Delta\phi_{KK}$

Fleischer, Jaarsma, and KKV[2016]



Matching the experimental precision of 0.5° requires

- 5% precision on differential rate of $B_s \rightarrow K^- \ell^+ \nu_\ell$ **not yet measured**
- 5% precision $SU(3)$ -breaking corrections **achievable**

The $B \rightarrow \pi K$ Puzzle

The $B \rightarrow \pi K$ puzzle

Buras, Fleischer, Recksiegel, Schwab [2004]; Fleischer, Recksiegel, Schwab [2007]; Fleischer, Jaeger, Pirjol, Zupan [2008]
Neubert, Rosner [1998]; Beneke, Neubert [2003]; Beaudry, Datta, London, Rashed, Roux [2018]

$B \rightarrow \pi K$ decays have been in the spotlight for decades

- Puzzling correlation between CP asymmetries found
- Large discrepancy between experiment and QCDF
- Electroweak penguins (EWP) contribute at the same level as Trees $\rightarrow V_{ub}$ suppressed
- **EWP sector** offers an interesting avenue for NP to enter via

$$qe^{i\phi} e^{i\omega} \equiv - \left(\frac{\hat{P}_{EW} + \hat{P}_{EW}^C}{\hat{T} + \hat{C}} \right)$$

The $B \rightarrow \pi K$ puzzle

Buras, Fleischer, Recksiegel, Schwab [2004]; Fleischer, Recksiegel, Schwab [2007]; Fleischer, Jaeger, Pirjol, Zupan [2008]
Neubert, Rosner [1998]; Beneke, Neubert [2003]; Beaudry, Datta, London, Rashed, Roux [2018]

$B \rightarrow \pi K$ decays have been in the spotlight for decades

- Puzzling correlation between CP asymmetries found
- Large discrepancy between experiment and QCDF
- Electroweak penguins (EWP) contribute at the same level as Trees $\rightarrow V_{ub}$ suppressed
- **EWP sector** offers an interesting avenue for NP to enter via

$$qe^{i\phi} e^{i\omega} \equiv - \left(\frac{\hat{P}_{EW} + \hat{P}_{EW}^C}{\hat{T} + \hat{C}} \right)$$

Electroweak penguin parameters

- $\phi(\omega)$ CP-violating (conserving) phases, ω model-independently small
- New CP violating physics might enter with large phase ϕ

Neubert, Rosner [1998]

CP asymmetries in $B \rightarrow \pi K$

Gronau[2005]; Gronau, Rosner [2006]

$$\begin{aligned} \Delta_{\text{SR}} = & \mathcal{A}_{\text{CP}}^{\text{dir}}(\pi^{\pm} K^{\mp}) + \mathcal{A}_{\text{CP}}^{\text{dir}}(\pi^{\pm} K^0) \frac{\text{Br}(\pi^{\pm} K^0)}{\text{Br}(\pi^{\pm} K^{\mp})} \frac{\tau_{B^0}}{\tau_{B^+}} \\ & - \mathcal{A}_{\text{CP}}^{\text{dir}}(\pi^0 K^{\pm}) \frac{2\text{Br}(\pi^0 K^{\pm})}{\text{Br}(\pi^{\pm} K^{\mp})} \frac{\tau_{B^0}}{\tau_{B^+}} - \mathcal{A}_{\text{CP}}^{\text{dir}}(\pi^0 K_S) \frac{2\text{Br}(\pi^0 K^0)}{\text{Br}(\pi^{\pm} K^{\mp})} = 0 + \mathcal{O}(\lambda^2) \end{aligned}$$

Sum rule provides a Standard Model test

- Satisfied experimentally \rightarrow still large uncertainties for $B_d^0 \rightarrow \pi^0 K^0$
- Predicts $\mathcal{A}_{\text{CP}}^{\text{dir}}(B_d^0 \rightarrow \pi^0 K^0) = -0.14 \pm 0.03$ (PDG: $A_{\text{CP}}^{\pi^0 K^0} = 0.00 \pm 0.13$)
- Intriguing opportunities for Belle II

Mixing-induced CP asymmetry in $B_d^0 \rightarrow \pi^0 K^0$ provides additional tests

Isospin Amplitude Triangles

Nir, Quin [1991]; Gronau, Hernandez, London, Rosner [1995]

Fleischer, Jaeger, Pirjol, Zupan [2008]; Fleischer, Jaarsma, KKV [2018]

$$\begin{aligned} & \sqrt{2}A(B^0 \rightarrow \pi^0 K^0) + A(B^0 \rightarrow \pi^- K^+) \\ &= \sqrt{2}A(B^+ \rightarrow \pi^0 K^+) + A(B^+ \rightarrow \pi^+ K^0) \\ &= -(\hat{T} + \hat{C})(e^{i\gamma} - qe^{i\phi}e^{i\omega}) \equiv 3A_{3/2} = 3|A_{3/2}|e^{i\phi_{3/2}}, \end{aligned}$$

- QCD penguin and colour-suppressed EWPs cancel
- Gives a clean correlation between the CP asymmetries in $B_d \rightarrow \pi^0 K_S$
- Minimal $SU(3)$ input

$$|\hat{T} + \hat{C}| = R_{T+C} |V_{us}/V_{ud}| \sqrt{2} |A(B^+ \rightarrow \pi^+ \pi^0)|$$


$$R_{T+C}|_{\text{fact}} = f_K/f_\pi = 1.2 \pm 0.2$$

Uncertainty accounts for non-factorizable $SU(3)$ breaking

Isospin Amplitude Triangles

Nir, Quin [1991]; Gronau, Hernandez, London, Rosner [1995]

Fleischer, Jaeger, Pirjol, Zupan [2008]; Fleischer, Jaarsma, KKV [2018]




Hints towards New Physics in the EWP sector?

Pinning down New Physics in EWP sector


see Poster by Ruben Jaarsma

Fleischer, Jaarsma, KKV [2018]; Fleischer, Jaarsma, Malami, KKV [2018]



Current data

Additional constraint from mixing-induced CP asymmetry



Benchmark scenario


Exciting prospects for Belle-II

CP violation in multibody decays

- Large part of the non-leptonic B decays
- Rich structure of CP violation
 - Especially for $B \rightarrow \pi\pi\pi$


Theoretically challenging:

- T-odd correlations Durieux, Grossman [2015]; Gronau, Rosner [2015]
- Using flavour symmetries Bhattacharya, Gronau, Imbeault, Rosner, London, Bediaga, Guerrer, de Miranda
- Applying CPT-invariance Nogueira, Bediaga, Cavalcante, Frederico, Lourenco [2015]; ...
- Using heavy meson chiral perturbation theory Cheng, Chua, Soni [2007]; Cheng, Chua, Zhang [2017]



QCD Factorization in three-body decays

Kraenkl, Mannel, Virto [2015]; Klein, Mannel, Virto, KKV [2017]



Factorization theorem at the phase space edge

$$\langle \pi^+ \pi^+ \pi^- | \mathcal{O}_i | B \rangle = T_i^I \otimes F^{B \rightarrow \pi^+} \otimes \Phi_{\pi^+ \pi^-} + T_i^I \otimes F^{B \rightarrow \pi^+ \pi^-} \otimes \Phi_{\pi^+}$$

- Improvement over quasi-two body interpretation
- Introduces **new non-perturbative strong phases**
 - Light-cone sum rules for $B \rightarrow hh$ form factors Khodjamirian, Cheng, Virto [2017]; Khodjamirian, Descotes-Genon, Virto, KKV [wip]
- Challenge: Reach the same level as two-body QCDF

Summary

Summary

- Extraction of γ from $B \rightarrow DK$ is theoretically clean
 - Impressive 1° precision in the upgrade era expected
 - Will play an increasingly important role as input parameter
- Penguin pollution in ϕ_s determinations under control
- Penguin dominated $B_s \rightarrow KK$ offers additional probe of ϕ_s
 - Requires analyses of $B_s^0 \rightarrow K^- \ell^+ \nu_\ell$
- $B \rightarrow \pi K$ decays remain puzzling \rightarrow good prospects
 - Improved CP asymmetries in $B_d \rightarrow \pi^0 K_S$ needed
 - Crucial to distinguish New Physics from QCD effects
- Three-body decays still offer many interesting avenues to explore
 - Study QCDF in $B^0 \rightarrow D^- \pi^+ \pi^0$

Summary


- Extraction of γ from $B \rightarrow DK$ is theoretically clean
 - Impressive 1° precision in the upgrade era expected
 - Will play an increasingly important role as input parameter
- Penguin pollution in ϕ_s determinations under control
- Penguin dominated $B_s \rightarrow KK$ offers additional probe of ϕ_s
 - Requires analyses of $B_s^0 \rightarrow K^- \ell^+ \nu_\ell$
- $B \rightarrow \pi K$ decays remain puzzling \rightarrow good prospects
 - Improved CP asymmetries in $B_d \rightarrow \pi^0 K_S$ needed
 - Crucial to distinguish New Physics from QCD effects
- Three-body decays still offer many interesting avenues to explore
 - Study QCDF in $B^0 \rightarrow D^- \pi^+ \pi^0$

Thank you for your attention

Back up



Constraints on new physics from the sum rule

- Limited sensitivity to q and ϕ for $q < 3$



Correlation between CP asymmetries in $B_d^0 \rightarrow \pi^0 K^0$

Fleischer, Jaeger, Pirjol, Zupan [2008]; Fleischer, Jaarsma, KKV [2018]



New element: constraint on angle $\phi_{\pm} = \arg(\bar{A}_{\pm} A_{\pm}^*)$

$$\phi_{\pm}|_{SM, \phi=0} = 2r \cos \delta \sin \gamma + \mathcal{O}(\lambda^2) = (8.7 \pm 3.5)^{\circ}$$

Pinning down New Physics in EWP sector

- Complement the isospin analysis with $S_{\text{CP}}^{\pi^0 K_S}$

$$\tan \phi_{00} = 2(r \cos \delta - r_c \cos \delta_c) \sin \gamma + 2r_c(\cos \delta_c - 2\tilde{a}_C/3)q \sin \phi + \mathcal{O}(\lambda^2)$$

- r, δ, r_c and δ_c hadronic parameters determined from $B \rightarrow \pi\pi$
- Only cosines of small phases, low sensitivity to variations
- Includes color-suppressed EWPs $\tilde{a}_C = a_C \cos(\Delta_C + \delta_c)$
- Effects included in a data-driven way

$$R \equiv \frac{\text{Br}(\pi^- K^+)}{\text{Br}(\pi^+ K^0)} = 0.89 \pm 0.04 = 1 - 2r \cos \delta \cos \gamma + 2r_c \tilde{a}_C q \cos \phi + \mathcal{O}(\lambda^2)$$

Controlling penguin effects in $B_s \rightarrow J/\psi\phi$

de Bruyn, Fleischer, JHEP 1503 [2015] 145

