



Contribution ID: 295

Type: not specified

Achievements and challenges in understanding nucleon-deuteron reactions

Comparison of theoretical predictions based on a nucleon-nucleon potential with data for elastic nucleon-deuteron (Nd) scattering and nucleon induced deuteron breakup reveals the importance of a three-nucleon force (3NF). Inclusion of semi-phenomenological 3NF models, such as Tucson-Melbourne or Urbana IX, into calculations in many cases improves the data description. However, some serious discrepancies remain even when a 3NF is included.

At low energies the prominent examples of discrepancies between theory and data were found for the vector analyzing power in elastic Nd scattering and for the neutron-deuteron (nd) breakup cross sections in neutron-neutron (nn) quasi-free-scattering (QFS) and symmetric-space-star (SST) geometries. Since both these configurations depend predominantly on the S-wave nucleon-nucleon (NN) force components, these cross section discrepancies have serious consequences for the nn $1S_0$ force component.

At energies above about 100 MeV current 3NF's only partially improve the description of data for cross section and spin observables in elastic Nd scattering and breakup. The complex angular and energy behavior of analyzing powers, spin correlation and spin transfer coefficients fails to be explained by standard nucleon-nucleon interactions alone or combined with current models of 3NF's.

One of the reasons for the above disagreements could be a lack of consistency between 2N and 3N phenomenological potentials used or/and omission of important terms in the applied 3NF.

The Chiral Effective Field Theory approach provides consistent two- and three-nucleon forces. The 3NF occurs for the first time at next-to-next-to

leading order (N²LO) of chiral expansion. The N³LO and N⁴LO NN forces when used in 3N calculations provide description of NN data of the same quality as standard, realistic NN potentials.

Application of improved, semilocal coordinate-space regularized chiral NN interactions up to N⁴LO order of chiral expansion combined with N²LO 3NF's supports conclusions obtained with standard forces. It can be expected that an application of consistent chiral NN and 3NF's up to N³LO will play an important role in understanding of elastic scattering and breakup reactions at higher energies.

Primary authors: Prof. WITALA, Henryk (Jagiellonian University); Prof. GOLAK, Jacek (Jagiellonian University Institute of Physics); Dr TOPOLNICKI, Kacper (Jagiellonian University Institute of Physics); Dr SKIBINSKI, Roman (Jagiellonian University)

Presenter: Prof. WITALA, Henryk (Jagiellonian University)