

Neutron capture reactions at the n_TOF facility at CERN

M. Barbagallo^{1,2}, on behalf of the n_TOF Collaboration²


1-Istituto Nazionale di Fisica Nucleare, sez. di Bari

2-CERN



GIANTS- IX Incontro dei Gruppi Italiani di Astrofisica Nucleare Teorica e Sperimentale
Palazzo Poggi, Bologna, 5-6 Ottobre 2017

- The Physics Case: s-process Nucleosynthesis
- The n_TOF facility at CERN (time-line, main features, “how-to”)
- Experimental program on neutron capture reactions



- ~~The Physics Case: s-process Nucleosynthesis~~
- The n_TOF facility at CERN (time-line, main features, “how-to”)
- Experimental program on neutron capture reactions








FIG. 6. Uncertainties of the stellar (n, γ) cross sections for s-process nucleosynthesis. These values refer to a thermal energy of $kT = 30$ keV, but may be considerably larger at lower and higher temperatures.










- Neutron spallation source
- Pulsed neutron beam produced by the high energy proton beam

- Two beam lines (**180m e 20m**)
- **Two** experimental areas (**EAR1** and **EAR2**), class A laboratories



n_TOF Time-line

F. Gunsing et al (n_TOF Coll.), Eur. Phys. J. Plus (2016) 131: 371


<p>May 1998</p>	<p>Feasibility CERN/LHC/98-02+Add</p>	<p>2000</p> <p>Commissioning</p> 	<p>2004-2007</p> <p>Upgrade</p> 	<p>May 2009</p> <p>Commissioning</p> 	<p>2010</p> <p>Upgrades: Borated-H₂O Second Line Class-A</p>	<p>2014-2018</p> <p>Phase III Capture: 14 Fission: 4 (n,cp): 4</p> <p>July 2014</p> <p>Commissioning of EAR-2</p>	
<p>1997 2017 ...</p>							
<p>1997</p>	<p>Concept by C.Rubbia CERN/ET/Int. Note 97-19</p>	<p>Aug 1998</p> <p>Proposal submitted</p>	<p>1999</p> <p>Construction started</p> 	<p>2001-2004</p> <p>Phase I Isotopes Capture: 25 Fission: 11</p> 	<p>2008</p> <p>New Target construction</p> 	<p>2009 - 2012</p> <p>Phase II Isotopes Capture: 14 Fission: 3 (n,cp): 2</p>	<p>2011</p> <p>EAR2 Design and Construction</p>  

Advantages of the Proton Synchrotron beam: high energy, high peak current (7e12 ppp/7 ns)

High instantaneous neutron flux
(10^5 n/cm²/pulse and 10^7 n/cm²/pulse).

Very convenient for measurements of:

- radioactive isotopes,
- low cross sections,
- isotopes available in small quantities



Other features of the neutron beams:



- High **resolution in energy** ($\Delta E/E = 10^{-4}$ in EAR1 and $\Delta E/E = 10^{-3}$ in EAR2)
- Wide **energy range** ($1 \text{ meV} < E_n < 1 \text{ GeV}$)
- Low **repetition** rate ($< 0.8 \text{ Hz}$) (no wrap-around)

C. Guerrero et al (n_TOF Coll.), Eur. Phys. J. A (2013) 49: 27

M. Barbagallo et al (n_TOF Coll.), Eur. Phys. J. A (2013) 49: 156

Two systems (and two different techniques) were used at n_TOF to measure neutron capture cross sections:

- Several C_6D_6 detectors \longrightarrow Total energy method
- A 4π BaF2 array (TAC) \longrightarrow Total absorption calorimetry




A cutting-edge technology for detection of γ rays is mandatory for high accuracy measurements

n_TOF at glance

■ Capture reactions (56)


■ Fission reactions (33)

■ (n,cp) reactions (8)




Nearly 50 (n, γ) reaction of interest for Nuclear Astrophysics studied at n_TOF so far:

- Branching points isotopes
- Abundancies in pre-solar grains
- Magic nuclei and end-points
- Seed-isotopes
- Cosmo-Cronometer




Nearly 50 (n, γ) reaction of interest for Nuclear Astrophysics studied at n_TOF so far:




- Branching points isotopes
- Abundancies in pre-solar grains
- Magic nuclei and end-points
- Seed-isotopes
- Cosmo-Cronometer




^{151}Sm ($t_{1/2}=93$ y), determines **abundance** of $^{152,154}\text{Gd}$




^{63}Ni ($t_{1/2}=100$ y) **first branching point** determines **abundance** of $^{63,65}\text{Cu}$



^{151}Sm ($t_{1/2}=93$ y), determines **abundance** of $^{152,154}\text{Gd}$



^{63}Ni ($t_{1/2}=100$ y) **first branching point** determines **abundance** of $^{63,65}\text{Cu}$




First, strong evidence of thermal pulsing in AGB stars.

Strong constraints on Cu, Ni and Zn production in massive stars and subsequent supernovae


U. Abbondanno et al., Phys Rev. Lett. 93, 161103 (2004)

C. Lederer et al., Phys. Rev. Lett. 110, 022501, (2013)

^{151}Sm ($t_{1/2}=93$ y), determines **abundance** of $^{152,154}\text{Gd}$



^{63}Ni ($t_{1/2}=100$ y) **first branching point** determines **abundance** of $^{63,65}\text{Cu}$




First, strong evidence of thermal pulsing in AGB stars.

Strong constraints on Cu, Ni and Zn production in massive stars and subsequent supernovae

U. Abbondanno et al., Phys Rev. Lett. 93, 161103 (2004)


C. Lederer et al., Phys. Rev. Lett. 110, 022501, (2013)

^{147}Pm ($t_{1/2}=2.6$ y) directly constrains the stellar reaction rate as used in the astrophysical models.



^{147}Pm ($t_{1/2}=2.6$ y) directly constrains the stellar reaction rate as used in the astrophysical models.

^{171}Tm ($t_{1/2}=1.9$ y) is branching point weakly dependent on stellar temperature \rightarrow constraint neutron density in low mass AGB stars.




The present (fraction of)

^{147}Pm ($t_{1/2}=2.6$ y) directly constrains the stellar reaction rate as used in the astrophysical models.

^{171}Tm ($t_{1/2}=1.9$ y) is branching point weakly dependent on stellar temperature \rightarrow constraint neutron density in low mass AGB stars.

^{204}Tl ($t_{1/2}=3.8$ y) capture cross-section affects the $^{205}\text{Pb}/^{205}\text{Tl}$ ratio and the s-only isotope ^{204}Pb .




The present (fraction of)

^{147}Pm ($t_{1/2}=2.6$ y) directly constrains the stellar reaction rate as used in the astrophysical models.

^{171}Tm ($t_{1/2}=1.9$ y) is branching point weakly dependent on stellar temperature \rightarrow constraint neutron density in low mass AGB stars.


^{204}Tl ($t_{1/2}=3.8$ y) capture cross-section affects the $^{205}\text{Pb}/^{205}\text{Tl}$ ratio and the s-only isotope ^{204}Pb .




Isotope production by neutron irradiation at @ ILL ($^{146}\text{Nd}(n,\gamma)^{147}\text{Nd}(\beta)$, $^{170}\text{Er}(n,\gamma)^{171}\text{Er}(\beta)$, $^{203}\text{Tl}(n,\gamma)^{204}\text{Tl}$)

Chemical separation and sample preparation @ PSI


^{171}Tm (1.9 y) - 3.6 mg sample



^{147}Pm (2.6 y) - 80 μg sample




^{204}Tl (3.8 y) - 11 mg sample (210 GBq)




A. Casanovas et al., in preparation

Isotope production by neutron irradiation at @ ILL ($^{146}\text{Nd}(n,\gamma)^{147}\text{Nd}(\beta)$, $^{170}\text{Er}(n,\gamma)^{171}\text{Er}(\beta)$, $^{203}\text{Tl}(n,\gamma)^{204}\text{Tl}$)
 Chemical separation and sample preparation @ PSI


^{171}Tm (1.9 y) - 3.6 mg sample



^{147}Pm (2.6 y) - 80 μg sample




^{204}Tl (3.8 y) - 11 mg sample (210 GBq)




A. Casanovas et al., in preparation

Isotope production by neutron irradiation at @ ILL ($^{146}\text{Nd}(n,\gamma)^{147}\text{Nd}(\beta)$, $^{170}\text{Er}(n,\gamma)^{171}\text{Er}(\beta)$, $^{203}\text{Tl}(n,\gamma)^{204}\text{Tl}$)
 Chemical separation and sample preparation @ PSI


^{171}Tm (1.9 y) - 3.6 mg sample



^{147}Pm (2.6 y) - 80 μg sample



^{204}Tl (3.8 y) - 11 mg sample (210 GBq)




A. Casanovas et al., in preparation

Preliminary results indicate **important differences** in the MACS relative to calculations and (very few) previous integral measurements.

The present (fraction of)





^{154}Gd and ^{156}Gd isotopes are of particular interest because their pure s-process origin allows to check the robustness of stellar models in galactic chemical evolution (GCE) models (^{13}C pocket).



The present (fraction of)

^{154}Gd and ^{156}Gd isotopes are of particular interest because their pure s-process origin allows to check the robustness of stellar models in galactic chemical evolution (GCE) models (^{13}C pocket).

Disagreement of more than 20% between observation and model calculation of s-process abundances



- S. Bisterzo, *et al.*, The Astrophysical Journal **787** (2014) 10
- C. Trippella, *et al.*, The Astrophysical Journal **787** (2014) 41
- S. Cristallo, *et al.*, The Astrophysical Journal **801** (2015) 53

To date a systematic study of all isotopes has never been carried out in the energy region from thermal to about 1 MeV. Accuracy so far limited by:

- The detector used in these experiments, which in some cases suffered from high neutron sensitivity
- The experimental determination of the neutron flux, which might have been biased in some previous measurements
- The lack of information on the cross section of impurities
- The enrichment of the samples and their quality in terms of canning and other material needed for the container.


To date a systematic study of all isotopes has never been carried out in the energy region from thermal to about 1 MeV. Accuracy so far limited by:

- The detector used in these experiments, which in some cases suffered from high neutron sensitivity
- The experimental determination of the neutron flux, which might have been biased in some previous measurements
- The lack of information on the cross section of impurities
- The enrichment of the samples and their quality in terms of canning and other material needed for the container.

Measurement at n_TOF-EAR1 of
 $^{152,154,155,156,157,158,160}\text{Gd}(n,\gamma)$

+

A priori and a posteriori sample
 characterization



use of an array of C_6D_6 detectors. The objective is to determine stellar cross sections with overall uncertainty below 5% for thermal energies up to about $kT = 100$ keV. In total 3.3×10^{18}

- There is need of **accurate new data** on neutron cross-section for **Nuclear Astrophysics**.
- The combination of **excellent resolution, unique brightness and low background** allows to collect at n_TOF high-accuracy data, in some cases for the first time ever.
- **Since 2001**, n_TOF@CERN has provided an important contribution on many capture cross section measurements for stellar nucleosynthesis. **Stay tuned for new results..**
- **A second experimental area (EAR2)** at 20 m flight path has recently made possible to perform very challenging measurements of interest for Nuclear Astrophysics, **on short-lived radionuclides, on sub-mg samples, and on low cross section reactions.**
- **A new spallation target will be installed before 2020, for the experimental program in Nuclear Astrophysics (and more) in the next decade.**