

Impact of electron-captures on nuclei near $N=50$ on core-collapse supernovae

Friday, 29 June 2018 10:30 (15 minutes)

Sensitivity studies of the late stages of stellar core collapse with respect to electron-capture rates indicate the importance of a region of nuclei near the $N=50$ shell closure, just above doubly magic ^{78}Ni . In the present work, it has been demonstrated that uncertainties in key characteristics of the evolution, such as the lepton fraction, electron fraction, entropy, stellar density, and in-fall velocity are about 50% due to uncertainties in the electron-capture rates on nuclei in this region, although thousands of nuclei are included in the simulations. The present electron-capture rate estimates used for the nuclei in this region of interest are primarily based on a simple approximation, and it is shown that the estimated rates are likely overestimated by an order of magnitude or more. More accurate microscopic theoretical models are required to obtain Gamow-Teller strength distributions, upon which electron-capture rates are based. The development of these models and the benchmarking of such calculations rely on data from charge-exchange experiments at intermediate energies. An experimental campaign to study Gamow-Teller strength distributions in nuclei at and near $N=50$, including ^{86}Kr and ^{88}Sr , with the $(t,^3\text{He})$ reaction at NSCL is underway and preliminary results, and their effects on future astrophysical work, will be presented.

Primary author: TITUS, Rachel (NSCL/MSU)

Co-authors: BROWN, B. Alex (NSCL/MSU); GAO, Bingshui (NSCL/MSU); SULLIVAN, Chris (NSCL/MSU); ZEGERS, Remco G. T. (NSCL/MSU)

Presenter: TITUS, Rachel (NSCL/MSU)

Session Classification: Stellar contribution: NS mergers