



ID contributo: 27

Tipo: non specificato

## On divisibility and quantum Markovianity for non-invertible dynamical maps

*venerdì 29 giugno 2018 09:30 (45 minuti)*

We analyze the relation between CP-divisibility and the lack of information back flow for an arbitrary not necessarily invertible dynamical map [1,2,3]. It is well known that CP-divisibility always implies lack of information back flow. Moreover, these two notions are equivalent for invertible maps. We show that for a map which is not invertible the lack of information back flow always implies the existence of completely positive (CP) propagator which, however, needs not be trace-preserving. Interestingly, for a class of image non-increasing dynamical maps [majority of examples studied in the literature do belong to this class] this propagator becomes trace-preserving as well and hence the lack of information back flow implies CP-divisibility. This result sheds new light into the structure of the time-local generators giving rise to CP-divisible evolutions. It is shown that if the map is not invertible then positivity of dissipation/decoherence rates is no longer necessary for CP-divisibility.

[1] D. Chruściński, A. Kossakowski, and A. Rivas, Phys. Rev. A 83, 052128 (2011).

[2] B. Bylicka, M. Johansson, and A. Acín, Phys. Rev. Lett. 118, 120501 (2017).

[3] D. Chruściński, A. Rivas, and E. Størmer, arXiv:1710.06771

**Autore principale:** Prof. CHRUSCINSKI, Dariusz (Nicolaus Copernicus University Torun)

**Relatore:** Prof. CHRUSCINSKI, Dariusz (Nicolaus Copernicus University Torun)