



Contribution ID: 86

Type: **Poster**

## **SuperCDMS IMPACT: an Ionization Yield Calibration Program**

*Tuesday, July 23, 2019 6:45 PM (15 minutes)*

The SuperCDMS collaboration has been developing cryogenic silicon and germanium detectors optimized for phonon signals from dark matter-nucleus collisions. The detectors are sensitive to dark matter masses between about 1 and 10  $\text{GeV}/c^2$ , which corresponds to sub-keV energy deposits from the nuclear recoil signal. The sensitivity of a SuperCDMS high voltage detector is achieved by applying a high voltage across the crystal. Under the electric field, the signal from electron-hole pairs generated from nuclear recoil events is amplified through the Neganov-Trofimov-Luke (NTL) effect. However, the yield of electron-hole pairs, which is critical to reconstructing the energy of the recoiling nucleus, is not well characterized in the sub-keV nuclear recoil energy region. I will describe a neutron scattering experiment called IMPACT (Ionization Measurement with Phonons At Cryogenic Temperatures), which is designed to measure the ionization yield in SuperCDMS style detectors.

### **Less than 5 years of experience since completion of Ph.D**

Y

### **Student (Ph.D., M.Sc. or B.Sc.)**

Y

**Primary author:** REN, Runze (Northwestern University)

**Presenter:** REN, Runze (Northwestern University)

**Session Classification:** Poster session

**Track Classification:** Low Temperature Detector Applications