

Contribution ID: 345

Type: Poster

X-ray spectroscopy of muonic atoms isolated in vacuum using transition edge sensors

Thursday, 25 July 2019 18:45 (15 minutes)

High-resolution X-ray spectroscopy of highly-charged muonic atoms/ions isolated in vacuum is an ideal probe to explore quantum electrodynamics (QED) effects. One of the major topic in fundamental atomic physics is to conduct these experiments in high-Z atom in which the bound particles experience extremely strong electric fields.

A negatively-charged muon can bind to a nucleus via the Coulomb field. This "muonic atom" is essentially hydrogen-like in its electronic structure. Since a muon is 200 times more massive than an electron, a muonic atom has a Bohr radius 200 times smaller than that of atomic hydrogen. This allows to test QED in strong field in a very different regime, since at such short distances the dominant QED contribution is the vacuum polarization, while it is the self-energy in highly-charged ions. After a negatively-charged muon is captured by the nucleus in a highly excited state, the muon peels off most (or all) of the electrons bound to the nucleus as Auger electrons, and thereby generating highly-charged muonic atoms in vacuum.

While a low-density target is required to avoid rapid refilling of electrons into the highly charged muonic atom from the surrounding atoms, it is experimentally difficult to efficiently stop muons in a low-density target. This is due to their large momentum distribution via traveling pion decay, resulting in insufficient x-ray yields with the conventional high-resolution x-ray spectroscopy technology based on diffraction from Bragg crystals, unless one uses a device like the PSI cyclotron trap.

We aim to realize the high-resolution muonic atom X-ray spectroscopy with low-density gas target with a combination of the world highest intensity pulsed negative muon beam at J-PARC MLF MUSE (Tokai, Japan) and an X-ray spectrometer based on a 240 pixel array of superconducting transition-edge-sensor (TES) microcalorimeters.

In April 2019 we will perform a feasibility test at J-PARC and report the outcome of these new result.

Less than 5 years of experience since completion of Ph.D

Ν

Student (Ph.D., M.Sc. or B.Sc.)

Ν

Primary author: Dr OKADA, Shinji (RIKEN)

Co-authors: REINTSEMA, Carl (NIST); SCHMIDT, Daniel (NIST); SWETZ, Daniel (NIST); Dr BENNETT, Douglas (NIST); O'NEIL, Galen (NIST); HILTON, Gene (NIST-Boulder); TATSUNO, Hideyuki; Dr NODA, Hi-rofumi (Osaka University); ULLOM, Joel (NIST/University of Colorado); GARD, Johnathon (NIST); FOWLER, Joseph (NIST); Dr NINOMIYA, Kazuhiko (Osaka University); Dr MORGAN, Kelsey (NIST); Prof. SHIMOMURA, Kouichiro (KEK); Dr DURKIN, Malcolm (NIST); Dr KATSURAGAWA, Miho (IPMU); Dr KAWAMURA, Nari-toshi (KEK); Prof. INDELICATO, Paul (CNRS); Dr CARADONNA, Pietro (IPMU); HAYAKAWA, Ryota (Tokyo

Metropolitan University); Dr WATANABE, Shin (JAXA); Dr TAKEDA, Shinichiro (IPMU); YAMADA, Shinya (Tokyo Metropolitan University); Dr KANDA, Sohtaro (RIKEN); Dr TAKESHITA, Soshi (KEK); ISOBE, Tadaaki (RIKEN); HASHIMOTO, Tadashi (Japan Atomic Energy Agency); Prof. TAKAHASHI, Tadayuki (IPMU); Dr OKU-MURA, Takuma (RIKEN); AZUMA, Toshiyuki (RIKEN); DORIESE, W. Bertrand (Randy) (NIST); Prof. MIYAKE, Yasuhiro (KEK); Dr UENO, Yasuhiro (RIKEN); Prof. KINO, Yasushi (Tohoku University); Dr ICHINOHE, Yuto (Rikkyo University)

Presenter: Dr OKADA, Shinji (RIKEN)

Session Classification: Poster session

Track Classification: Low Temperature Detector Applications