

Chern-Simons gravity coupled to a spectator field during inflation

Based on:

- *N. Bartolo, G. Orlando, arXiv:1706.04627*
- *Ongoing project*

Giorgio Orlando

Padua University, Department of Physics and Astronomy

ASI/COSMOS meeting - "Astroparticle and Fundamental Physics with the CMB", Ferrara, 26 June 2018

Chern-Simons gravity

$$\mathcal{L} = \sqrt{g} \left[\frac{1}{2} M_{Pl}^2 R - \frac{1}{2} g^{\mu\nu} \partial_\mu \chi \partial^\mu \chi - V(\chi) \right] + f(\chi) \epsilon^{\mu\nu\rho\sigma} R_{\mu\nu}{}^{\kappa\lambda} R_{\rho\sigma\kappa\lambda}.$$

Peculiarities of the Chern-Simons term:

- $\epsilon^{\mu\nu\rho\sigma} R_{\mu\nu}{}^{\kappa\lambda} R_{\rho\sigma\kappa\lambda}$ is a surface term, we need the coupling $f(\chi)$.
- **P broken**, T broken, CPT preserved.

→ Standard **GR restored** if at a certain time χ **decades**.

→ Search **remnant signatures** of Chern-Simons gravity **from inflation**.

(χ can be both the inflaton field ϕ or a **spectator field** in multi-field models)

- Key observable: **chirality of primordial GW** $\longrightarrow \Theta = \frac{P_\gamma^R - P_\gamma^L}{P_\gamma^R + P_\gamma^L}$.

$$\gamma_R = \frac{1}{\sqrt{2}}(\gamma_+ - i\gamma_\times),$$

$$\gamma_L = \frac{1}{\sqrt{2}}(\gamma_+ + i\gamma_\times).$$

Linear analysis


- $\tau \equiv$ conformal time

Quadratic action of GW

$$S|_{\gamma\gamma} = \sum_{s=L,R} \int d\tau \frac{d^3k}{(2\pi)^3} A_{\gamma,s}^2 \left[|\gamma'_s(\tau, k)|^2 - k^2 |\gamma_s(\tau, k)|^2 \right],$$

$$A_{\gamma,s}^2 = \frac{M_{Pl}^2}{2} a^2 \left(1 - \lambda_s \frac{k_{phys}}{M_{CS}} \right), \quad M_{CS} = M_{Pl}^2 / 8\dot{f}(\chi).$$

- M_{CS} : **Chern-Simons mass**, characteristic energy scale.
- $\lambda_R = +1, \lambda_L = -1 \implies$ When $k_{phys} > M_{CS}$, γ_R shows **instabilities**.



- We consider **physical scales** that at the beginning of inflation were **inside Hubble radius**, but **outside the Chern-Simons scale**.

$\longrightarrow H/M_{CS} \ll 1.$

Equations of motion for GW

- Field redefinition: $\mu_s = A_{\gamma,s} \gamma_s$.

$$\mu_{R/L}'' + k^2 \left(1 - \frac{2}{k^2 \tau^2} \pm \mathcal{V} \right) \mu_{R/L} = 0.$$

$$\mathcal{V} = \frac{1}{k\tau} \frac{H}{M_{CS}} \mathcal{A},$$

$$\mathcal{A} = \frac{1}{(1 \mp k_p/M_{CS})^2} \left\{ \left[1 - \xi + \frac{1}{2}\omega - \frac{1}{2H} \frac{1}{\tau} \xi \right] \left(1 \mp \frac{k_p}{M_{CS}} \right) + \mp \frac{k_p}{2M_{CS}} \left[\frac{1}{2} + \xi + \frac{1}{2}\xi^2 \right] \right\}.$$

$$k_p = k/a, \quad \xi = \frac{\dot{M}_{CS}}{M_{CS}H}, \quad \omega = \frac{\ddot{M}_{CS}}{M_{CS}H^2}.$$

• $H/M_{CS} \ll 1$ (previous slide) \longrightarrow \mathcal{V} vanishes vs standard gravity.

Example: $M_{CS} = \text{constant case}$

- E.o.m.'s become

$$\mu''_{R/L} + k^2 \left(1 - \frac{2}{k^2 \tau^2} \pm \frac{1}{k\tau} \frac{H}{M_{CS}} \right) \mu_{R/L} = 0 .$$

- It is Whittaker equation, we can find analytic solutions.

Super-horizon power-spectra (M. Satoh, 2010)

$$P_{\gamma}^L = \frac{P_{\gamma}}{2} e^{-\frac{\pi}{4} H/M_{CS}} ,$$

$$P_{\gamma}^R = \frac{P_{\gamma}}{2} e^{+\frac{\pi}{4} H/M_{CS}} .$$


$$\Theta = \frac{P_{\gamma}^R - P_{\gamma}^L}{P_{\gamma}^R + P_{\gamma}^L} = \frac{\pi}{2} \frac{H}{M_{CS}} .$$

- $H/M_{CS} \ll 1 \rightarrow \Theta \ll 1 .$

$$P_{\gamma} = \frac{4}{k^3} \frac{H^2}{M_{Pl}^2}$$

Forecasts from CMB data

- TB and EB correlators are sensitive to GW chirality.



M. Gerbino et. al. (2016)

- Impossible to observe small chirality given the current experimental constraint on $r = P_\gamma/P_S$, $r < 0.07$ (95%CL) $\rightarrow \sigma_\Theta \geq 0.3$.

Higher order statistics

- At linear level we get no measurable imprints.
- Need for a non-linear analysis.
- In JCAP07(2017)034 “Parity breaking signatures from a Chern-Simons coupling during inflation: the case of non-Gaussian gravitational waves” N. Bartolo, G. Orlando, **we treated the bispectrum statistics of the model.**

In-In formalism

$$\langle \delta_a(\vec{k}_1) \delta_b(\vec{k}_2) \delta_c(\vec{k}_3) \rangle(t) = -i \int_{t_0}^t dt' \langle 0 | \left[\delta_a^I(\vec{k}_1, t) \delta_b^I(\vec{k}_2, t) \delta_c^I(\vec{k}_3, t) , H_{int}^I(t') \right] | 0 \rangle .$$

- This formalism is **based on quantum interaction picture**. It allows to compute/estimate non-Gaussian cosmological correlators starting from free fields (i.e. using linear solutions).

Main results

- Amplitudes of $\langle \gamma\gamma\gamma \rangle$ and $\langle \gamma\delta\chi\delta\chi \rangle$ are suppressed by H/M_{CS} .
- **Amplitude of $\langle \gamma\gamma\delta\chi \rangle$ bispectrum** shows a different behaviour:

$$\begin{aligned} \langle \gamma_R(\vec{k}_1)\gamma_R(\vec{k}_2)\delta\chi(\vec{k}_3) \rangle &= (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{\pi}{64} \left(\sum_{i \neq j} P_\gamma(k_i) P_\gamma(k_j) \right) \times \\ &\times \frac{\dot{\chi}}{H} \left(H^2 \frac{\partial^2 f(\chi)}{\partial^2 \chi} \right) \frac{(k_1 + k_2)k_1 k_2}{\sum_i k_i^3} \cos\theta (1 - \cos\theta)^2, \end{aligned}$$

$$\langle \gamma_L(\vec{k}_1)\gamma_L(\vec{k}_2)\delta\chi(\vec{k}_3) \rangle = -\langle \gamma_R(\vec{k}_1)\gamma_R(\vec{k}_2)\delta\chi(\vec{k}_3) \rangle, \quad \langle \gamma_L(\vec{k}_1)\gamma_R(\vec{k}_2)\delta\chi(\vec{k}_3) \rangle = 0.$$

- The amplitude of this bispectrum is **proportional to the second order derivative** of $f(\chi) \rightarrow$ No suppression.
- Paper in preparation, *N. Bartolo, G. Orlando, M. Shiraishi*: **forecast** on the possibility to constrain the theory **from CMB bispectra**.


Squeezed shape function

- The shape function gives the dependence of the bispectrum on the three momenta k_i .


$$F(k_1^\gamma, k_2^\gamma, k_3^{\delta\chi}) = \left(\sum_{i \neq j} \frac{1}{k_i^3 k_j^3} \right) \frac{(k_1 + k_2) k_1 k_2}{\sum_i k_i^3} \cos \theta (1 - \cos \theta)^2,$$

$$\cos \theta = \frac{k_3^2 - k_2^2 - k_1^2}{2k_1 k_2}.$$

$$x_2 = k_2/k_1, \quad x_3 = k_3/k_1.$$



- The **shape function is maximum in the squeezed limit** when the momentum of $\delta\chi$ is much smaller than the momenta of the two γ 's ($k_3 \ll k_1 \simeq k_2$).



Squeezed modulation (1)

- The **squeezed limit** of a generic 3-points function gives the correlation of 2 short modes with 1 long mode. This **induces a modulation on the short modes**.

Squeezed modulation of the tensor power spectra

$$P_{\gamma}^{R/L}(q)|_{\delta\chi} = P_{\gamma}^{R/L}(q)|_{(0)} + \delta\chi(Q) \frac{\langle \gamma_{R/L}(q_1) \gamma_{R/L}(q_2) \delta\chi(Q) \rangle'}{P_{\delta\chi}(Q)},$$

$$Q \rightarrow 0, \quad q_1 \simeq q_2 = q.$$

- **Physical interpretation**: after horizon crossing the scalar long mode freezes at a certain amplitude, inducing a **local modification of the spatial curvature**. The short modes evolve in this modified background space and **inherits a specific modulation in their power spectrum statistics**.
- R and L squeezed bispectra differ for a minus sign.
→ **Contribution to the GW chirality**

Squeezed modulation (2)

Cumulative effect of soft modes

$$\mathcal{M}(q, \vec{x})_{R/L} = \int_{Q_{min} < |\vec{Q}| < Q_L} d^3 Q e^{i\vec{Q} \cdot \vec{x}} \delta\chi(Q) \frac{\langle \gamma_{R/L}(q) \gamma_{R/L}(q) \delta\chi(Q) \rangle'}{P_{\delta\chi}(Q)}.$$

Local amplification of chirality

$$P_{\gamma}^{R/L}(q; \vec{x}) = \frac{P_T(q)}{2} [1 \pm \mathcal{A}(\vec{x})], \quad \Theta = \mathcal{A}(\vec{x}).$$

- $\mathcal{A}(\vec{x})$ follows a Gaussian distribution with mean 0 and variance

$$\sigma^2 = 32\pi^2 \epsilon_{\chi} \left(\frac{H^2 f''(\chi_0)}{M_{Pl}} \right)^2 \int_{Q_{min}}^{Q_L} dQ Q^2 P_{\delta\chi}(Q).$$

$$\epsilon_{\chi} = \dot{\chi}_0^2 / 2H^2 M_{Pl}^2.$$

- χ has to be a spectator field. Squeezed bispectra are re-absorbed in single clock-inflation, consequence of gauge freedom. (see, e.g., *Creminelli et. al., 2013; Pajer et. al., 2013*).

Summary and Conclusion

- Chern-Simons gravity coupled to a scalar field χ can be tested during inflationary epoch.
- At linear level the model can not be constrained through CMB experiments. Chirality produced is low if we avoid instabilities.
- Non-linearities produce a model dependent local squeezed amplification of chirality. χ has to be different from the inflaton field.
- In progression: toy model building, parameter space.
- What about interferometers?
 - ▶ Coplanar interferometers are not sensitive to chirality of GW. We need a more complicated geometry (*Smith and Caldwell, 2016*).
 - ▶ Inflationary GW are red-tilted. We need a small scale amplification mechanism.
 - ▶ Idea: study the effect of chiral GW production while χ is decaying (Next step).

Thank You!

Back-up slide, example of a toy model

- Assume $\rho_\chi \ll \rho_\phi$, also take $\epsilon_H = -\frac{\dot{H}}{H^2} \simeq \epsilon_\phi$. ($\epsilon_\chi \ll \epsilon_\phi$), $\eta_\chi \simeq 0$.
- Both ϕ and χ have vacuum fluctuations on a quasi De-Sitter space.
 $P_\chi, P_\phi \simeq H^2/2k^3$.

$$\frac{\zeta_\chi}{\zeta_\phi} = \frac{\delta\rho_\chi}{\delta\rho_\phi} \simeq \frac{V'(\chi)\delta\chi}{U'(\phi)\delta\phi} = \sqrt{\frac{\epsilon_\chi}{\epsilon_\phi}} \ll 1 \longrightarrow \zeta \simeq \zeta_\phi.$$

- Postulate $f(\chi)$ as:

$$f(\chi) = \lambda \left(\frac{\chi}{M_{Pl}} \right)^n,$$

$$\longrightarrow \sigma^2 = 32\pi^2 \epsilon_\chi (H^2 f''(\chi_0))^2 \left(\frac{H}{M_{Pl}} \right)^2 \times \ln \left(\frac{Q_L}{Q_{min}} \right).$$

- Take $\left(\frac{H}{M_{Pl}} \right) = 10^{-5}$, $\epsilon_\chi = 10^{-6}$, $\ln \left(\frac{Q_L}{Q_{min}} \right) = 10$, $H^2 f''(\chi_0) = 10^6$:
 $\longrightarrow \sigma^2 = 1.$

- $\frac{H}{M_{CS}} \ll 1 \longrightarrow \left(\frac{\chi_0}{M_{Pl}} \right) \ll 10^{-4}$, small field potential for χ .