PHOTON 2019 - International Conference on the Structure and the Interactions of the Photon. Satellite Workshop: Photon Physics and Simulation at Hadron Colliders.

PHOTON 2019 - International Conference on the Structure and the Interactions of the Photon

3-7 June 2019

Satellite Workshop:

INFN - LNF, Frascati

Photon Physics and Simulation at Hadron Colliders 6-7 June 2019

Contribution ID: 30 Type: Talk

Production of W^+W^- and $t\bar{t}$ pairs in photon-photon mechanism in proton-proton scattering

Tuesday, 4 June 2019 11:45 (20 minutes)

Photon-induced processes in proton-proton interactions have become recently very topical. The large energy at the LHC, when combined with relatively large luminosity at run II, allows to start the exploration of such processes.

We discuss production of W^+W^- pairs and $t\bar{t}$ quark-antiquark pairs in proton-proton collisions induced by two-photon fusion including, for a first time, transverse momenta of incoming photons. The unintegrated inelastic fluxes (related to proton dissociation) of photons are calculated based on modern parametrizations of deep inelastic structure functions in a broad range of x and Q^2 .

We focus on processes with single and double proton dissociation. Highly excited remnant systems hadronise producing particles that can be vetoed in the calorimeter. We calculate associated effective gap survival factors. The gap survival factors depend on the process, mass of the remnant system and collision energy. The rapidity gap survival factor due to remnant fragmentation for double

dissociative (DD) collisions is smaller than that for single dissociative (SD) process. We observe approximate factorisation: $S_{R,DD} \approx S_{R,SD}^2$ when imposing rapidity veto. For the W^+W^- final state, the remnant fragmentation leads to a taming of the cross section when the rapidity gap requirement is imposed. Also for $t\bar{t}$ quark-antiquark pairs such a condition reverses the hierarchy observed for the case when such condition is taken into account.

Our results imply that for the production of such heavy objects as t quark and \bar{t} antiquark the virtuality of the photons attached to the dissociative system are very large ($Q^2 < 10^4 \text{ GeV}^2$). A similar effect was observed for the W^+W^- system.

- 1. M. Luszczak, L. Forthomme, W. Schafer, A. Szczurek, arXiv:1810.12432; JHEP (2019) in print.
- L. Forthomme, M. Luszczak, W. Schafer, A. Szczurek, Phys.Lett. B789 (2019) 300-307.
- M. Luszczak, W. Schafer, A. Szczurek, JHEP 1805 (2018) 064.
- M. Luszczak, W. Schafer and A. Szczurek, Phys. Rev. D93 (2016) 7, 074018.
- 5. M. Luszczak, A. Szczurek, Ch. Royon, JHEP 1502 (2015) 098.

Summary

Primary authors: SZCZUREK, Antoni (Institute of Nuclear Physics, Krakow and Rzeszow University, Rzeszow); Dr ŁUSZCZAK, Marta (Faculty of Mathematics and Natural Sciences, University of Rzeszow, Poland); Dr

LAURENT, Forthomme (Helsinki Institute of Physics); SCHAEFER, Wolfgang (Institute of Nuclear Physics PAN)

Presenters: SZCZUREK, Antoni (Institute of Nuclear Physics, Krakow and Rzeszow University, Rzeszow); Dr ŁUSZCZAK, Marta (Faculty of Mathematics and Natural Sciences, University of Rzeszow, Poland); Dr LAURENT, Forthomme (Helsinki Institute of Physics); SCHAEFER, Wolfgang (Institute of Nuclear Physics PAN)

Session Classification: Gamma-Gamma Collisions

Track Classification: Gamma-Gamma Collisions