



Contribution ID: 87

Type: Poster

Nuclei in the limits of the drip line: a study of the regions $N=20$ and $N=34$

By using a non-relativistic independent particle approach, we have investigated the mechanism promoting $N = 34$ as a new magic number [1]. We carried out Hartree-Fock plus Bardeen-Cooper Schrieffer and Quasi-particle Random Phase Approximation calculations by consistently using the same finite-range interaction in all the steps of our approach [2]. We used four Gogny-like interactions, with and without tensor terms. We found that the shell closure for $N = 34$ neutrons appear in isotones with proton number smaller than 26. The smaller is the proton number, the more evident is the shell closure at $N=34$. For this reason, we studied in detail the nucleus ^{48}Si , that appears to be the nucleus with the lowest value of Z for $N = 34$ before the drip line [3]. By using the same techniques, we have investigated also the mechanism producing the extinction of the $N = 20$ neutron shell closure in the nucleus ^{32}Mg [4]. We studied the contribution of the tensor interaction in the emergence of new magic numbers or in the disappearance of the traditionally ones.

REFERENCES

- [1] D. Steppenbeck et al., Nature 502, 207 (2013).
- [2] V. De Donno et al., Phys. Rev. C95, 054329 (2017).
- [3] J. Erler et al., Nature 486, 509 (2012).
- [4] A. Chaudhuri et al., Phys. Rev. C88, 054317 (2013).

Primary author: Dr ANGUIANO, Marta (Universidad de Granada)

Co-authors: Dr CO', Giampaolo (Università del Salento); Dr LALLENA, Antonio M. (Universidad de Granada)

Presenter: Dr ANGUIANO, Marta (Universidad de Granada)

Session Classification: POSTER SESSION