



Contribution ID: 190

Type: Poster

Reactions with ^9Li at HIE-ISOLDE

The increase of energy of the ISOLDE radioactive beams made possible through the HIE-ISOLDE project has opened new possibilities for reactions studies with exotic beams. A particular challenge is presented by the light nuclei where unbound final states play an important role and the role of the continuum is by now known to be important. Theoretical work on the reaction mechanisms involved aiming for a detailed description of the reaction processes is ongoing and evolves as new experimental data appear.

In particular we are interested in the region of neutron-rich Li-isotopes. ^{11}Li is a famous example of a so-called “nuclear halo”, where loosely bound neutrons extends to large distances. However, to understand and aid the theoretical description we require better experimental information on both ^{11}Li itself, but also ^{10}Li and ^9Li .

I shall report on results from two experiments performed in the Scattering Experiments Chamber (SEC), both using ^9Li as reaccelerated radioactive beam. The first experiment ran November 2016 with a beam energy of 6.8 MeV/u on a deuterated polyethylene target. The second experiment took place in the beginning of November 2018 with a beam energy of 8 MeV/u on a deuterated polyethylene target. Multiple reaction channels are open at these energies, but the major one of interest are the elastic and one-neutron transfer reactions to the isotopes ^8Li and ^{10}Li . Particularly interesting are the final channels leading to ^{10}Li and where knowledge on the detailed structure is still lacking in spite of many earlier experiments at widely different beam energies.

The presentation will give an overview of the experimental set-up used and report on the results that have been extracted so far. A comparison will also be given between the current results and the $^9\text{Li}+2\text{H}$ experiments carried out earlier at REX-ISOLDE at beam energies of 2.3 MeV/u and 2.8 MeV/u.

Primary authors: Mr JENSEN, Jesper (Aarhus University (DK)); Prof. RIISAGER, K (Department of Astronomy and Physics, University of Aarhus)

Co-authors: Prof. TENGBLAD, Olof (IEM - CSIC); SOTTY, Ch. (“Horia Hulubei” National Institute of Physics and Nuclear Engineering, RO-077125 Bucharest, Romania); Mr PEREA, Angel (IEM-CSIC); Dr MARTEL, Ismael (Universidad de Huelva); DÍAZ OVEJAS, Javier (CSIC); CEDERKALL, Joakim (Lund University); NILSSON, Thomas (Chalmers University of Technology); Dr NACHER, Enrique (IEM-CSIC); Dr MUNCH, Michael (Department of Astronomy and physics, University of Aarhus); LICA, R. (CERN, CH-1211 Geneva 23, Switzerland); JONSON, Bjorn (Fundamental Physics, Chalmers University of Technology); FRAILE, L. M. (Grupo de Física Nuclear and UPARCOS, Universidad Complutense de Madrid, CEI Moncloa, E-28040 Madrid, Spain); FIGUERA, Pierpaolo (LNS); BERGE, M. J. G. (Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain); FYNBO, Hans O. U. (Department of Physics and Astronomy, University of Aarhus, Denmark); VIÑALS, Silvia (CSIC)

Presenter: Mr JENSEN, Jesper (Aarhus University (DK))

Session Classification: POSTER SESSION