

# Experimental studies of neutron-rich nuclei around $N = 126$ at KEK isotope separation system

Y.X. Watanabe  
KEK Wako Nuclear Science Center (WNSC)




## Contents

1. Introduction: astrophysical motivation
2. Nuclear production around  $N = 126$  by MNT reactions
3. KISS facility and recent experimental results
4. R&D and future plan of KISS
5. Summary


# r-process nucleosynthesis and $N = 126$ neutron-rich nuclei

Nuclear properties of neutron closed shell  $N = 126$  nuclei  
→ r-process in astrophysical nucleosynthesis




H. Grawe et al., Rept. Prog. Phys. 70, 1525 – 1582 (2007).

Lifetime and mass of waiting point nuclei  
→ Astrophysical environments of r-process




# Uncertainties of r-process abundance pattern: half-life

Theoretical nuclear models play crucial roles in the simulation of the r-process nucleosynthesis.



**Isotopic abundance pattern variance  
from uncertain  $\beta$ -decay half-lives  
( $0.1 \sim 10 \times T_{1/2}$ )**



KISS project   
KIS Isotope Separation System


Systematic nuclear spectroscopy (lifetime, mass,  $\beta$ - $\gamma$  spectroscopy, laser spectroscopy) around  $N = 126$   
→ Astrophysical environments of r-process

M.R. Mumpower et al., Prog. Part. Nucl. Phys. 86, 86 (2016).

# Nuclear production around $N = 126$ by MNT reaction



## Macroscopic approach

(Langevin-type dynamical equation of motion)




V. Zagrebaev and W. Greiner, Phys. Rev. Lett. 101, 122701 (2008).

## $^{136}\text{Xe} + ^{208}\text{Pb}$ (5.8 MeV/A) at ANL



## Semi-classical approach

(Single-particle transfer probability)



## GRAZING calculation

A. Winther, Nucl. Phys. A 572, 191 – 235 (1994);  
594, 203 – 245 (1995).


J.S. Barrett et al., Phys. Rev. C 91, 064615 (2015).

# Experimental study for MNT reactions of $^{136}\text{Xe} + ^{198}\text{Pt}$

$^{136}\text{Xe}$  (8 MeV/nucleon) +  $^{198}\text{Pt}$  (1.3 mg/cm<sup>2</sup>)



Projectile-like fragments (PLFs) were detected by large acceptance magnetic spectrometer VAMOS++ at GANIL, and target-like fragment (TLF) distributions were deduced.

Y.X. Watanabe et al., Phys. Rev. Lett. 115, 172503 (2015).




# TKEL dependence of isotopic distribution


TKEL: Total Kinetic Energy Loss




# KEK Isotope Separation System (KISS)




# KISS experimental setup



MSPGC (Multi-Segmented Proportional Gas Counter)




Four HPGe (High-Purity Germanium) clover detectors





8  
Collaboration with IBS (Korea)

# Experimental results




# $\beta$ - $\gamma$ spectroscopy of $^{198}\text{Os}$


Y. Hirayama et al., Phys. Rev. C 98, 014321 (2018).



# Systematics of osmium isotopes




H. Koura, JAEA-Review 2010-056, pp.83–84 (2010).



# Laser spectroscopy of nuclei around $N = 126$

## Laser spectroscopy → Hyperfine structure

- $\mu, Q \rightarrow$  Wave-function
  - Isotope shift  $\rightarrow$  Charge radius  $\rightarrow$  Nuclear deformation






## In-gas-cell laser ionization spectroscopy at KISS for study of nuclear structure

D. Verney et al., Eur. Phys. J. A 30, 489 (2006).

# HFS measurement for ground and isomeric states of $^{199}\text{Pt}$


*Y. Hirayama et al., Phys. Rev. C 96, 014307 (2017).*



# HFS measurement for ground and isomeric states of $^{199}\text{Pt}$

*Y. Hirayama et al., Phys. Rev. C 96, 014307 (2017).*

Nuclide	$I^\pi$	$\mu (\mu_N)$	$\delta\langle r^2 \rangle^{194,A} (\text{fm}^2)$	$ \langle \beta_2^2 \rangle ^{1/2}$
$^{199\text{m}}\text{Pt}$	(13/2)+	-0.57(5)	0.166(30)	0.110(12)
$^{199\text{g}}\text{Pt}$	5/2-	+0.75(8)	0.268(34)	0.144(10)




# High-precision In-gas-jet laser spectroscopy


More precise study of wave-function and deformation

← High-precision laser spectroscopy

Narrow-band LD laser + Dye amplifier + Intense YAG laser



In-gas-jet laser spectroscopy  
 $^{194}\text{Pt}$  ( $I^\pi = 0^+$ )  
Width : 0.60(1) GHz (FWHM)



Ar gas cell: 80 kPa,  $P_{\text{B.G.}} \sim 50$  Pa


# 3D tracking gas counter for low-background rate of 0.01 cps

To go further to lifetime measurements of more neutron-rich nuclei, lower background rate of the gas counter is necessary ( $\sim 0.01$  cps)

Proportional gas counter: Ar + CH<sub>4</sub>(10%), 0.1 MPa


2D tracking :  $\Delta\Omega = 80\%$ , background rate 0.1 cps

2-layered 16-segmented proportional gas counters




Anode wire: Carbon wire ( $\phi 10$  mm,  $3 \text{ k}\Omega/\text{cm}$ )  
→ Longitudinal hit-positions of  $\beta$ -rays can be identified.  
→ Better separation from the B.G.



One proto-type counter




2 mm in FWHM



# Mass measurements at KISS



P. Schury et al., Phys. Rev. C 95, 011305(R) (2017).



# Summary

- Systematic nuclear spectroscopy (lifetime, mass,  $\beta$ - $\gamma$  spectroscopy, laser spectroscopy) around neutron magic number 126 are important for identification of astrophysical environments of r-process.
- MNT reactions are promising for production of neutron-rich nuclei around  $N = 126$ .
- KEK Isotope Separation System (KISS)
  - MNT reactions of  $^{136}\text{Xe} + ^{198}\text{Pt}$
  - Gas cell + Laser ionization + ISOL
    - Efficient collection and separation of MNT reaction products
  - Lifetime measurements &  $\beta$ - $\gamma$  spectroscopy  
 $^{199-201}\text{Pt}$ ,  $^{196-200}\text{Ir}$ ,  $^{195-198}\text{Os}$  ( $^{136}\text{Xe} + ^{198}\text{Pt}$ ),  $^{185-187}\text{Ta}$  ( $^{136}\text{Xe} + \text{nat.W}$ )
  - Laser spectroscopy  
 $^{199}\text{Pt}$ ,  $^{196-198}\text{Ir}$
  - High-precision in-gas-jet laser spectroscopy was prepared
  - 3D tracking gas counter is under development
  - Mass measurements with MRTOF-MS is planned

KISS is open for external user programs

Pre-proposals will be discussed in the SSRI-PNS collaboration meeting (September)

[SSRI-PNS\\_contact@kek.jp](mailto:SSRI-PNS_contact@kek.jp)

*THANK YOU FOR YOUR ATTENTION*

# Collaboration

## KISS project



**KEK**

**IBS**

**Seoul National University**

**Tsukuba University**

**CNS**

**RIKEN**

**K.U. Leuven**

Y. Hirayama, Y. Kakiguchi, H. Miyatake,  
M. Oyaizu, P.H. Schury, M. Wada, Y.X. Watanabe  
S.C. Jeong, J.Y. Moon, J.H. Park  
H.S. Choi  
M. Mukai, M. Ahmed  
N. Imai  
H. Ishiyama, S. Kimura, T. Sonoda  
P. Van Duppen , Yu. Kudryavtsev, M. Huyse

## MNT measurements at GANIL

**KEK**

**IBS**

**CNS**

**GANIL**

**Torino University**

**LNL**

**Padova University**

**Seoul National University**

**University of Tokyo**

**RIKEN**

**Osaka University**

Y. Hirayama, H.S. Jung, H. Miyatake, Y.X. Watanabe  
H. Ishiyama, S.C. Jeong  
N. Imai  
Y.H. Kim, M. Rejmund, C. Schmitt, A. Navin,  
G. de France, E. Clement  
G. Pollarolo  
L. Corradi, E. Fioretto  
D. Montanari  
S.H. Choi, J.S. Song  
M. Niikura  
D. Suzuki  
H. Nishibata, J. Takatsu