

Heterogeneity, visco-elasticity and the universality class of earthquake occurrence

Wednesday, 11 December 2019 15:45 (15 minutes)

The organization in time, space and energy of earthquakes exhibits scaling laws with exponents α , β , γ and p which are universal, i.e. they are independent of time and of the geographic region.

A possible explanation of this critical-like behavior is provided by the possibility to describe the evolution of a seismic fault

under friction within the general context of the depinning transition. In fact, minimal models for the seismic fault, such as the Burridge-Knopoff model, can be mapped to classical quenched Edwards-Wilkinson (qEW) interfaces. Nevertheless the exponents of the scaling laws

of the qEW interfaces are different from those exhibited by instrumental earthquakes.

In this talk I will present a more realistic description of the seismic fault which can be viewed as a qEW interface evolving over

a viscous-ductile substrate. I will show that this description produces scaling laws with exponents α , β , γ and p in very good agreement with experimental values. More precisely, the values of the exponents α , β and γ are quite independent of model parameters and this explains their universal character. Conversely the exponent p , controlling the temporal clustering of seismic sequences,

depends on the specific law implemented for the viscosity. The value $p = 1$ of instrumental data is recovered assuming a velocity-strengthening law for the viscous layer, which is a quite realistic description of real fault systems.

Primary authors: LIPPIELLO, Eugenio; Prof. ROSSO, Alberto; Dr LANDES, François; Dr PETRILLO, Giuseppe

Presenter: LIPPIELLO, Eugenio

Session Classification: Session 3