


# Scenario for a world wide $e^+e^-$ collider


---

**LC09** Perugia, Italy, 21-24 September, 2009

F. Richard LAL/Orsay



F. Richard



# Outline

---

- Introduction
- Challenges
- Present views
- What can we expect to observe
- Alternate scenarios
- Expectations from first data at LHC
- Summary and conclusions

# Introduction

---



- There is a consensus that to get adequate accuracies, e.g. on the Higgs sector, we need a leptonic collider beyond LHC
- By far the most advanced project is the TeV SC collider called ILC which is developed by an worldwide collaboration
- It aims at a TDR for end of 2012
- CLIC for a multi-TeV collider is in an R&D phase and intends to prove its feasibility (CDR) by 2010
- These machines are very challenging projects in comparison to LEP2/SLC
- A muon collider is studied at FNAL
- Even more futuristic R&D is actively performed with Laser-plasma and Beam-plasma acceleration

# The ILC project

---

- ILC has a large 5% 'prototype' under construction, the [DESY XFEL](#), and intense R&D on critical aspects in Asia, Europe and NA
- Italy (+other Europeans like FR, Ge, SP, UK, CERN) is actively contributing to ILC through XFEL and studies in DAFNE (e-cloud, kickers)
- GDE, under B. Barish, is an international organization set by [ICFA](#), recognized by the 3 regions which is about to produce an 'almost' ready for construction project to be proposed to governments in 2012
- ILC works with a large community ~1000 physicists and engineers preparing detectors and refurbishing solid physics arguments in favor of such a project

# CLIC and ILC layouts



# Some parameters

$$L \sim \eta \frac{P_{\text{electrical}}}{E_{\text{CM}}} \sqrt{\frac{\delta E}{\varepsilon_{n,y}}} H_D$$

Type	LEP200	SLC100	ILC500	CLIC500
Vertical size nm	4000	700	5.7	2.3
Total P MW	65	50	216	129.4
Wall plug transf % $\eta$			9.4	7.4
Luminosity $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$	5	0.2	1500	1400
Interval between bunches ns	>>>	>>>	176	0.5
Polarisation %	No	80	>80	>80
Gradient MV/m	8	17	31.5	100

- ILC and CLIC intend to start at 500 GeV
- ILC is upgradable, with present technology, at 1 TeV
- CLIC could reach 3 TeV but with  $\sim$ constant luminosity (same  $\delta$ )



# Physics at ILC

---

- ❑ Physics arguments in favor of ILC are solid
- ❑ Precise measurements with a light Higgs as predicted within SUSY and the SM interpretation of LEP/SLC/TeVatron precision measurements (PM)
- ❑ Energy 0.5-1 TeV optimal to cover Higgs physics and presumably lightest SUSY particles
- ❑ ILC accuracy needed for the Higgs and SUSY sectors (as illustrated below)
- ❑ SC technology is well suited for this energy range but, while not strictly limited to 1 TeV cannot, with present SC materials, go well beyond

# $ee \rightarrow Z^* \rightarrow HZ$

- The recoil mass technique with  $Z \rightarrow \mu^+ \mu^-$  gives a very clean signal at  $\sqrt{s} = M_H + 110$  GeV
- Works even if H decays into invisible or complex modes
- ZZH coupling constant determined to 1%
- In the SM case most BR ratios known 10 times more precisely than at LHC




**Full Simulation**



# Dark matter & SUSY

- With LHC+LC it is possible to reach sufficient accuracy on the predicted dark matter to match cosmological observations
- Do they coincide ?



# Challenges & Hopes

---

- Why is this simple and convincing vision facing several challenges ?
- In the US, using its own costing approach (x3-4 6 B\$),DOE is judging the project too expensive and is no more proposing to build it in the US it is however contributing substantially ~35 M\$ (FNAL, JLAB, Cornell) and achieving good progress
- In Europe CERN is heavily committed on LHC but there is an initiative to allow a '[Scientific and geographical enlargement of CERN](#)' which would facilitate an organized participation to an international project (not necessarily at CERN). CLIC-ILC collaboration helps.
- Japan is highly committed on JPARC and super-Belle but after the Nobel shows increased motivation (lobbying at parliament, contacts with industry) to house ILC

# Breaking news

<http://www.physorg.com/news172317407.html>

---



- **CERN boss wants to bid for linear collider**
- September 16th, 2009 CERN's director general Rolf-Dieter Heuer will push for the linear collider, the next big experiment in particle physics after the Large Hadron Collider (LHC), to be built at the Geneva lab. Heuer made his call to situate the linear collider at CERN in an exclusive video **interview with *Physicsworld***, which is being relaunched today, Wednesday 16 September.

# Physics ?

---

- ❑ Physics arguments have also been challenged
- ❑ So far there is no compelling evidence for SUSY
- ❑ There is no unique prediction for SUSY masses (could even escape LHC)
- ❑ There are competing scenarios without necessarily a light Higgs and even Higgsless

# Present view


---

- ❑ Get ready technically to propose a construction of ILC end 2012
- ❑ See what comes out from LHC (and Tevatron)
- ❑ See if there are convincing indications from these results to trigger a decision on ILC
- ❑ If not wait and prepare for alternate scenarios: CLIC (or  $\mu$  colliders?)
- ❑ In this view, advocated by CERN DG, ILC and CLIC projects have decided to share efforts
- ❑ Also true for Detectors where CERN has signed the 3 proposed ILC concepts



# Detectors

- They need also to be ready end 2012 and well integrated to the machine(push pull issue)
- 3 Letters of Intent (1000 P+I) have been examined for validation by peer review (IDAG) during summer
- **SiD** **ILD** have been validated based on PFLOW ideas
- Important to pursue R&D on the multi-readout technology proposed by the **Fourth** concept
- Challenging detectors quite different from LEP
- CERN has joined this effort and intends to use the same detector concepts for CLIC
- Does it work? Seems OK for WW/ZZ separation but watch for duty cycle effects

# JETS



- ❑ High granularity+high density (SiW)
- ❑  $\mu$ electronics integrated inside calorimeters
- ❑ Possible with new technology+power pulsing
- ❑ Requires R&D



ILC : 100 $\mu$ W/ch

Physics Proto. 18ch 10\*10mm 5mW/ch

ATLAS LAr FEB 128ch 400\*500mm 1 W/ch

# Alternate Physics Scenarios

---


- ❑ What are the alternate physics scenarios, and how can they influence our choices?
- ❑ Most of these scenarios have a hard time to pass PM
- ❑ It seems however possible to accommodate a heavy Higgs and even an absence of Higgs
- ❑ Examples: strongly coupled field theory (TC) dual to extra dimensions (RS), 4<sup>th</sup> generation, BESS etc...
- ❑ These models provide S,T extra contributions and therefore alter the light Higgs prediction ('conspiracy')
- ❑ What could a LC observe in such scenarios ?



# RS in a nutshell

- The Randall Sundrum model provides an interpretation for Planck/EW and fermion masses hierarchies with **no new scale**
- S,T constraints requires extended groups and hence not only KK states but also  $Z'$  and 'custodians'
- KK bosons couple preferentially to  $b$  and even more to  $t$ , most likely  $t_R$
- AFB $b$  at LEP1 could be interpreted within RS by  $Z$ - $Z'$  mixing in RS
- AFB $t$  indication at Tevatron could be interpreted as Gkk exchange

Planck




Higgs

$SU(2)_L \times SU(2)_R$


$\begin{pmatrix} t \\ b \end{pmatrix}_L$

$t_R$



# Example on ZZH

- ❑ Reduced ZZH coupling
- ❑ Not a problem at ILC (but severe for LEP2)
- ❑ Could be the only signal if KK are heavy in which case ILC could be sensitive to this new physics well beyond LHC (up to  $\sim 10$  TeV)




G. Moreau C. Bouchart  
LPT-ORSAY-09-70

Giudice et al.  
**JHEP 0706:045,2007.**

# Top physics



- Plays a very peculiar role in most of these models
- In RS tR couples preferentially to  $Z'$  through  $Z$ - $Z'$  mixing
- Large effect on  $ALR_t$  expected from the AFBb ( $M_{Z'}$  up to  $\sim 10$  TeV)

[A. Djouadi et al.](#)  
*Nucl.Phys.B773:43-64,2007*



# Top at ILC

- ❑ LC 1 pb, LHC 1nb but with larger uncertainties
- ❑ Very good s/b at ILC and energy conservation allows to reconstruct modes with a neutrino
- ❑  $M_t$  and  $\Gamma_t$  with 50 MeV error, 0.4% on cross section
- ❑ Polarisation allows to separate tR and tL (extra dimensions)





# Higgsless scenarios

---

- Hyp: SM Higgs excluded by TeVatron+LHC
- Truly Higgsless ?
- There is the distinct possibility that a light Higgs was missed even at LEP2 if it cascades to 2 light CP odd Higgs NMSSM
- Would be covered by ILC irrespective of its decay modes (if narrow resonance) and with reduced xsection
- Without Higgs (RS, TC, BESS) several deviations expected in  $ee \rightarrow WW, ZWW, WW\bar{\nu}\nu$  better observed at CLIC
- Very demanding in luminosity at LHC which would delay any decision to build a LC
- Instead one could observe  $\sim$ TeV resonances which could be accessible earlier

# BESS model


- In extra dimensions (and 'deconstructed' versions) they require additional Z to control unitarity violation in WLWL
- ILC could see a signal but it will require CLIC to see the whole picture
- In some cases very large luminosity is needed both at LC and LHC



# How soon can LHC give answers?

- Will start with reduced lumi & energy ( $\leq 5$  TeV/beam) and with few 100 pb-1 not enough to discover the Higgs boson (shut down end 2010 to reach 7 TeV)
- CMSSM could be explored with 200pb-1 at 5+5 TeV
- AFBt not easy to confirm since at LHC  $q\bar{q} \ll gg$
- Heavy quarks  $< 500$  GeV either from 4<sup>th</sup> generation or from RS are accessible
- BESS needs  $> 1$  fb-1 ?

## LHC REACH VS CMSSM



mueller, Cavanaugh, De Roeck, Ellis, Flacher, Heinemeyer, Isidori, Olive, Paradisi, Ronga, Weiglein

# Summary on the HEP strategy

---

- ❑ Connect CLIC and ILC efforts to avoid duplication and potentially damaging competition
- ❑ Prepare for major challenges: technical (industrialisation 16000 SC cavities), financial (~6 B\$), political with a worldwide machine (LHC different, ~ITER ?) OCDE, ESFRI
- ❑ ILC and CLIC projects intend to address these problems
- ❑ Present uncertainties justify an open scenario
- ❑ However ILC is ready to go while it will take longer to complete the CLIC project



# In conclusion


---

- The HEP community has developed a consistent and worldwide strategy to construct an  $e^+e^-$  LC
- A viable project, ILC, can be presented to the governments end of 2012
- A final decision (ILC/CLIC) will depend on the physics results from LHC (or Tevatron)


# Why so precise ?

## Deviations from SM

(By S. Yamashita)



**SUSY**  
**(2 Higgs Doublet Model)**




**Extra dimension**  
**(Higgs-radion mixing)**


# An example of 'conspiracy'

parameter set	$m_{u_4}$	$m_{d_4}$	$m_H$	$\Delta S_{\text{tot}}$	$\Delta T_{\text{tot}}$
(a)	310	260	115	0.15	0.19
(b)	320	260	200	0.19	0.20
(c)	330	260	300	0.21	0.22
(d)	400	350	115	0.15	0.19
(e)	400	340	200	0.19	0.20
(f)	400	325	300	0.21	0.25

- G. Kribs et al  
<http://arxiv.org/abs/0706.3718v>
- Heavy Higgs allowed



# Combination of 0j and 2j, H to WW to II





# CLIC 3 TeV main parameters



Center-of-mass energy	CLIC conserv.	CLIC Nominal
Total (Peak 1%) luminosity	1.5(0.73)10 <sup>34</sup>	5.9(2.0)·10 <sup>34</sup>
Repetition rate (Hz)	50	
Loaded accel. gradient MV/m	100	
Main linac RF frequency GHz	12 (NC)	
Bunch charge10 <sup>9</sup>	3.72	
Bunch separation ns	0.5	
Beam pulse duration (ns)	156	
Beam power/linac (MWatts)	14	
Hor./vert. norm. emitt (10 <sup>-6</sup> /10 <sup>-9</sup> )	3 / 40	2.4 / 25
Hor/Vert FF focusing (mm)	10/0.4	8/0.1
Hor./vert. IP beam size (nm)	83 / 2.0	40 / 1.0
Soft Hadronic event at IP	0.57	2.7
Coherent pairs/crossing at IP	5 10 <sup>7</sup>	3.8 10 <sup>8</sup>
BDS length (km)	2.75	
Total site length (km)	48.3	
Wall plug to beam transfer eff.	6.8%	
Total power consumption (MW)	415	



# LC 500 GeV Main parameters



Center-of-mass energy	ILC	CLIC Conserv.	CLIC Nominal
Total (Peak 1%) luminosity	2.0(1.5)·10 <sup>34</sup>	0.9(0.6)·10 <sup>34</sup>	2.3(1.4)·10 <sup>34</sup>
Repetition rate (Hz)	5	50	
Loaded accel. gradient MV/m	33.5	80	
Main linac RF frequency GHz	1.3 (SC)	12 (NC)	
Bunch charge 10 <sup>9</sup>	20	6.8	
Bunch separation ns	176	0.5	
Beam pulse duration (ns)	1000	177	
Beam power/linac (MWatts)	10.2	4.9	
Hor./vert. norm. emitt (10 <sup>-6</sup> /10 <sup>-9</sup> )	10/40	3 / 40	2.4 / 25
Hor/Vert FF focusing (mm)	20/0.4	10/0.4	8/0.1
Hor./vert. IP beam size (nm)	640/5.7	248 / 5.7	202/ 2.3
Soft Hadronic event at IP	0.12	0.07	0.19
Coherent pairs/crossing at IP	10?	10	100
BDS length (km)	2.23 (1 TeV)	1.87	
Total site length (km)	31	13.0	
Wall plug to beam transfer eff.	9.4%	7.5%	
Total power consumption MW	216	129.4	