

# Conformal invariance of TMD rapidity evolution

Giovanni Antonio Chirilli

University of Regensburg

Resummation Evolution Factorization 2019


University of Pavia, Italy  
28 November, 2019

based on [Phys.Rev. D100 \(2019\) no.5, 051504](#)

- Motivation.
- BK equation and  $SL(2, C)$  Möbius group.
- Rapidity factorization for particle production in hadron-hadron collisions.
- Conformal invariance of TMD Operators.
- Conformal TMD Rapidity evolution equation in Sudakov region.
- Conclusions and outlook.

# Particle production in hadron-hadron collisions

e.g. production of Higgs particle



- typical TMD region:  $s \sim q^2 = m_X^2 \gg q_\perp^2 \sim 1\text{Gev}$
- Sudakov region:  $s \sim q^2 = m_X^2 \gg q_\perp^2 \gg 1\text{Gev}$
- small-x region :  $s \gg q^2 \sim q_\perp^2 \gg m_N^2$

$$\frac{d\sigma}{d\eta d^2q_\perp} = \sum_f \int d^2b_\perp e^{i(q,b)_\perp} \mathcal{D}_{f/A}(x_A, b_\perp, \eta) \mathcal{D}_{f/B}(x_B, b_\perp, \eta) \sigma(ff \rightarrow H) \\ + \text{power corrections} + \text{Y-terms}$$

- rapidity:  $\eta = \frac{1}{2} \ln \frac{q^+}{q^-}$ .
- $\mathcal{D}_{f/A}$ : TMD density of parton  $f$  in hadron  $A$ .
- $\sigma(ff \rightarrow H)$ : cross section of production of particle  $H$  of invariant mass  $m_H^2 = Q^2$  from two partons scattering.
- Power corrections:  $\frac{q_\perp^2}{Q^2}$ .
- Y-terms: for  $q_\perp^2 \sim q^2$  allow transition to collinear factorization formula.

- TMD evolution equations are analyzed by different methods at moderate  $x_B$ , CSS and SCET, and at small- $x_B$  resulting in different evolution equations.
- At the future Electron Ion Collider, TMD will be probed from low to high  $x_B$ . It is then necessary to develop a formalism which is valid in both limits.
- In the region of moderate  $x_B$  TMD analysis is performed with a combination of UV and rapidity cutoff which results in two evolution equations, in  $\mu^2$  and  $\zeta$  (related to rapidity). See Collins' book
  - Such evolution equations are known at two and three-loop, but their relation to the conformal properties of TMD is not known.

- **I. Balitsky and A. Tarasov (2016):** Evolution equation for gluon TMD valid for all  $x_B = \beta_B$  and all  $k_\perp (\geq 1\text{GeV})$ .

$$\begin{aligned}
 \frac{d}{d \ln \sigma} \tilde{\mathcal{F}}_i^a(\beta_B, x_\perp) \mathcal{F}_j^a(\beta_B, y_\perp) &= -\alpha_s \text{Tr} \left\{ \int d^2 k_\perp (x_\perp | \left\{ U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} (U k_k + p_k U) \frac{\sigma \beta_{BS} g_{\mu i} - 2k_\mu^\perp k_i}{\sigma \beta_{BS} + k_\perp^2} \right. \right. \\
 &- 2k_\mu^\perp g_{ik} U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} U - 2g_{\mu k} U^\dagger \frac{p_i}{\sigma \beta_{BS} + p_\perp^2} U + \left. \left. \frac{2k_\mu^\perp}{k_\perp^2} g_{ik} \right\} \tilde{\mathcal{F}}^k \left( \beta_B + \frac{k_\perp^2}{\sigma S} \right) | k_\perp \right) \\
 &\times (k_\perp | \mathcal{F}^l \left( \beta_B + \frac{k_\perp^2}{\sigma S} \right) \left\{ \frac{\sigma \beta_{BS} \delta_j^\mu - 2k_\perp^\mu k_j}{\sigma \beta_{BS} + k_\perp^2} (k_l U^\dagger + U^\dagger p_l) \frac{1}{\sigma \beta_{BS} + p_\perp^2} U \right. \\
 &\quad \left. - 2k_\perp^\mu g_{jl} U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} U - 2\delta_l^\mu U^\dagger \frac{p_j}{\sigma \beta_{BS} + p_\perp^2} U + 2g_{jl} \frac{k_l^\mu}{k_\perp^2} \right\} | y_\perp ) \\
 &+ 2\tilde{\mathcal{F}}_i(\beta_B, x_\perp) (y_\perp | \frac{p^m}{p_\perp^2} \mathcal{F}_k(\beta_B) (i \overleftarrow{\partial}_l + U_l) (2\delta_m^k \delta_j^l - g_{jm} g^{kl}) U^\dagger \frac{1}{\sigma \beta_{BS} - p_\perp^2 + i\epsilon} U \\
 &\quad + \mathcal{F}_j(\beta_B) \frac{\sigma \beta_{BS}}{p_\perp^2 (\sigma \beta_{BS} - p_\perp^2 + i\epsilon)} | y_\perp ) \\
 &+ 2(x_\perp | - U^\dagger \frac{1}{\sigma \beta_{BS} - p_\perp^2 - i\epsilon} U (2\delta_i^k \delta_m^l - g_{im} g^{kl}) (i\partial_k - U_k) \tilde{\mathcal{F}}_l(\beta_B) \frac{p^m}{p_\perp^2} \\
 &\quad \left. + \tilde{\mathcal{F}}_i(\beta_B) \frac{\sigma \beta_{BS}}{p_\perp^2 (\sigma \beta_{BS} - p_\perp^2 - i\epsilon)} | x_\perp ) \mathcal{F}_j(\beta_B, y_\perp) \right\} + \mathcal{O}(\alpha_s^2)
 \end{aligned}$$

Light-cone limit  $\Rightarrow$  DGLAP

$$\frac{d}{d\eta} \alpha_s \mathcal{D}(x_B, \mathbf{0}_\perp, \eta) = \frac{\alpha_s}{\pi} N_c \int_{x_B}^1 \frac{dz'}{z'} \left[ \left( \frac{1}{1-z'} \right)_+ + \frac{1}{z'} - 2 + z'(1-z') \right] \alpha_s \mathcal{D}\left(\frac{x_B}{z'}, \mathbf{0}_\perp, \eta\right)$$

Light-cone limit  $\Rightarrow$  DGLAP

$$\frac{d}{d\eta} \alpha_s \mathcal{D}(x_B, \mathbf{0}_\perp, \eta) = \frac{\alpha_s}{\pi} N_c \int_{x_B}^1 \frac{dz'}{z'} \left[ \left( \frac{1}{1-z'} \right)_+ + \frac{1}{z'} - 2 + z'(1-z') \right] \alpha_s \mathcal{D}\left(\frac{x_B}{z'}, \mathbf{0}_\perp, \eta\right)$$

Low-x limit:  $p_\perp \sim (x-y)^{-2} \ll 1$  non-linear evolution equation

$$\begin{aligned} & \frac{d}{d\eta} \tilde{U}_i^a(x) U_j^a(y) \\ &= -\frac{\alpha}{2\pi^2} \text{Tr}\{(-i\partial_i^x + \tilde{U}_i^x) \left[ \int d^2z (\tilde{U}_x \tilde{U}_z^\dagger - 1) \frac{(x-y)^2}{(x-z)^2 (y-z)^2} (U_z U_y^\dagger - 1) \right] (i\overleftarrow{U}_j^y + U_j^y)\} \end{aligned}$$

with  $U_i = \partial_i U$  and  $\frac{(x-y)^2}{(x-z)^2 (y-z)^2}$  BK kernel



Light-cone limit  $\Rightarrow$  DGLAP

$$\frac{d}{d\eta} \alpha_s \mathcal{D}(x_B, \mathbf{0}_\perp, \eta) = \frac{\alpha_s}{\pi} N_c \int_{x_B}^1 \frac{dz'}{z'} \left[ \left( \frac{1}{1-z'} \right)_+ + \frac{1}{z'} - 2 + z'(1-z') \right] \alpha_s \mathcal{D}\left(\frac{x_B}{z'}, \mathbf{0}_\perp, \eta\right)$$

Low-x limit:  $p_\perp \sim (x-y)^{-2} \ll 1$  non-linear evolution equation

$$\begin{aligned} & \frac{d}{d\eta} \tilde{U}_i^a(x) U_j^a(y) \\ &= -\frac{\alpha}{2\pi^2} \text{Tr}\{(-i\partial_i^x + \tilde{U}_i^x) \left[ \int d^2z (\tilde{U}_x \tilde{U}_z^\dagger - 1) \frac{(x-y)^2}{(x-z)^2 (y-z)^2} (U_z U_y^\dagger - 1) \right] (i\overleftarrow{U}_j^y + U_j^y)\} \end{aligned}$$

with  $U_i = \partial_i U$  and  $\frac{(x-y)^2}{(x-z)^2 (y-z)^2}$  BK kernelDouble-Log:  $1 \gg \sigma \gg \frac{(x-y)_\perp^{-2}}{s}$  and  $\sigma x_{BS} \gg p_\perp^2 (x-y)_\perp^{-2}$   $\eta = \ln \sigma$ 

$$\frac{d}{d\eta} \mathcal{D}(x_B, z_\perp, \ln \sigma) = \frac{\alpha_s N_c}{\pi} \mathcal{D}(x_B, z_\perp, \ln \sigma) \int \frac{d^2 p_\perp}{p_\perp^2} [e^{i(p,z)_\perp} - 1]$$

Result is complicated and not unique. **Conformal Invariance may help.**

$$\begin{aligned}
 \frac{d}{d \ln \sigma} \tilde{\mathcal{F}}_i^a(\beta_B, x_\perp) \mathcal{F}_j^a(\beta_B, y_\perp) &= -\alpha_s \text{Tr} \left\{ \int d^2 k_\perp (x_\perp | \left\{ U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} (U k_k + p_k U) \frac{\sigma \beta_{BS} g_{\mu i} - 2k_\mu^\perp k_i}{\sigma \beta_{BS} + k_\perp^2} \right. \right. \\
 &- 2k_\mu^\perp g_{ik} U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} U - 2g_{\mu k} U^\dagger \frac{p_i}{\sigma \beta_{BS} + p_\perp^2} U + \left. \left. \frac{2k_\mu^\perp}{k_\perp^2} g_{ik} \right\} \tilde{\mathcal{F}}^k \left( \beta_B + \frac{k_\perp^2}{\sigma s} \right) | k_\perp \right) \\
 &\times (k_\perp | \mathcal{F}^l \left( \beta_B + \frac{k_\perp^2}{\sigma s} \right) \left\{ \frac{\sigma \beta_{BS} \delta_j^\mu - 2k_\perp^\mu k_j}{\sigma \beta_{BS} + k_\perp^2} (k_l U^\dagger + U^\dagger p_l) \frac{1}{\sigma \beta_{BS} + p_\perp^2} U \right. \\
 &\quad \left. - 2k_\perp^\mu g_{jl} U^\dagger \frac{1}{\sigma \beta_{BS} + p_\perp^2} U - 2\delta_l^\mu U^\dagger \frac{p_j}{\sigma \beta_{BS} + p_\perp^2} U + 2g_{jl} \frac{k_\perp^\mu}{k_\perp^2} \right\} | y_\perp ) \\
 &+ 2\tilde{\mathcal{F}}_i(\beta_B, x_\perp) (y_\perp | \frac{p^m}{p_\perp^2} \mathcal{F}_k(\beta_B) (i \overleftarrow{\partial}_l + U_l) (2\delta_m^k \delta_j^l - g_{jm} g^{kl}) U^\dagger \frac{1}{\sigma \beta_{BS} - p_\perp^2 + i\epsilon} U \\
 &\quad + \mathcal{F}_j(\beta_B) \frac{\sigma \beta_{BS}}{p_\perp^2 (\sigma \beta_{BS} - p_\perp^2 + i\epsilon)} | y_\perp ) \\
 &+ 2(x_\perp | - U^\dagger \frac{1}{\sigma \beta_{BS} - p_\perp^2 - i\epsilon} U (2\delta_i^k \delta_m^l - g_{im} g^{kl}) (i \partial_k - U_k) \tilde{\mathcal{F}}_l(\beta_B) \frac{p^m}{p_\perp^2} \\
 &\quad \left. + \tilde{\mathcal{F}}_i(\beta_B) \frac{\sigma \beta_{BS}}{p_\perp^2 (\sigma \beta_{BS} - p_\perp^2 - i\epsilon)} | x_\perp ) \mathcal{F}_j(\beta_B, y_\perp) \right\} + \mathcal{O}(\alpha_s^2)
 \end{aligned}$$


# Conformal properties of 6-point functions in $\mathcal{N}=4$ SYM

For TMD we need to study 6-point correlation functions in  $\mathcal{N}=4$  SYM theory

$$\mathcal{S} \equiv \frac{4\pi^2\sqrt{2}}{\sqrt{N_c^2-1}} \text{Tr}\{Z^2\} \quad (Z = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)) \text{ renorm-invariant chiral primary operator}$$

$\langle T\{\mathcal{S}(z_1)\mathcal{S}(z_2)\mathcal{S}(z_3)\mathcal{S}(z_4)\mathcal{S}(z_5)\mathcal{S}(z_6)\} \rangle$       Work in progress    I. Balitsky and G.A.C.

8 (out of 9) conformal ratios will contribute.




# Conformal properties of 6-point functions in $\mathcal{N}=4$ SYM

For TMD we need to study 6-point correlation functions in  $\mathcal{N}=4$  SYM theory

$$\mathcal{S} \equiv \frac{4\pi^2\sqrt{2}}{\sqrt{N_c^2-1}} \text{Tr}\{Z^2\} \quad (Z = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)) \text{ renorm-invariant chiral primary operator}$$

$\langle T\{\mathcal{S}(z_1)\mathcal{S}(z_2)\mathcal{S}(z_3)\mathcal{S}(z_4)\mathcal{S}(z_5)\mathcal{S}(z_6)\} \rangle$       Work in progress I. Balitsky and G.A.C.

8 (out of 9) conformal ratios will contribute.



Simpler case has been studied: 4-point correlation functions in the Regge limit:  
only two conformal ratios.

$$\hat{U}(x, y) \equiv 1 - \frac{1}{N_c} \text{tr}\{\hat{U}(x_\perp)\hat{U}^\dagger(y_\perp)\}$$

$$\frac{d}{d\eta}\hat{U}(x, y) = \frac{\alpha_s N_c}{2\pi^2} \int \frac{d^2z}{(x-z)^2(y-z)^2} \left\{ \hat{U}(x, z) + \hat{U}(z, y) - \hat{U}(x, y) - \hat{U}(x, z)\hat{U}(z, y) \right\}$$

- LLA for DIS in pQCD  $\Rightarrow$  BFKL
  - (LLA:  $\alpha_s \ll 1, \alpha_s \eta \sim 1$ ): describes proliferation of gluons.
- LLA for DIS in semi-classical-QCD  $\Rightarrow$  BK eqn
  - background field method: describes recombination process.

Formally, a light-like Wilson line

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] = \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} dx^+ A_+(x^+, x_\perp) \right\}$$

is invariant under inversion (with respect to the point with  $x^- = 0$ ).

Formally, a light-like Wilson line

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] = \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} dx^+ A_+(x^+, x_\perp) \right\}$$

is invariant under inversion (with respect to the point with  $x^- = 0$ ).

Indeed,

$(x^+, x_\perp)^2 = -x_\perp^2 \Rightarrow$  after the inversion  $x_\perp \rightarrow x_\perp/x_\perp^2$  and  $x^+ \rightarrow x^+/x_\perp^2$

# Conformal invariance of the BK equation

Formally, a light-like Wilson line

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] = \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} dx^+ A_+(x^+, x_\perp) \right\}$$

is invariant under inversion (with respect to the point with  $x^- = 0$ ).

Indeed,

$(x^+, x_\perp)^2 = -x_\perp^2 \Rightarrow$  after the inversion  $x_\perp \rightarrow x_\perp/x_\perp^2$  and  $x^+ \rightarrow x^+/x_\perp^2 \Rightarrow$

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] \rightarrow \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} d\frac{x^+}{x_\perp^2} A_+\left(\frac{x^+}{x_\perp^2}, \frac{x_\perp}{x_\perp^2}\right) \right\} = [\infty p_1 + \frac{x_\perp}{x_\perp^2}, -\infty p_1 + \frac{x_\perp}{x_\perp^2}]$$



# Conformal invariance of the BK equation

Formally, a light-like Wilson line

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] = \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} dx^+ A_+(x^+, x_\perp) \right\}$$

is invariant under inversion (with respect to the point with  $x^- = 0$ ).

Indeed,

$(x^+, x_\perp)^2 = -x_\perp^2 \Rightarrow$  after the inversion  $x_\perp \rightarrow x_\perp/x_\perp^2$  and  $x^+ \rightarrow x^+/x_\perp^2 \Rightarrow$

$$[\infty p_1 + x_\perp, -\infty p_1 + x_\perp] \rightarrow \text{Pexp} \left\{ ig \int_{-\infty}^{\infty} d\frac{x^+}{x_\perp^2} A_+\left(\frac{x^+}{x_\perp^2}, \frac{x_\perp}{x_\perp^2}\right) \right\} = [\infty p_1 + \frac{x_\perp}{x_\perp^2}, -\infty p_1 + \frac{x_\perp}{x_\perp^2}]$$

$\Rightarrow$  The dipole kernel is invariant under the inversion  $V(x_\perp) = U(x_\perp/x_\perp^2)$

$$\frac{d}{d\eta} \text{Tr}\{V_x V_y^\dagger\} = \frac{\alpha_s}{2\pi^2} \int \frac{d^2 z}{z^4} \frac{(x-y)^2 z^4}{(x-z)^2 (z-y)^2} [\text{Tr}\{V_x V_z^\dagger\} \text{Tr}\{V_z V_y^\dagger\} - N_c \text{Tr}\{V_x V_y^\dagger\}]$$

## SL(2,C) for Wilson lines

$$\hat{S}_- \equiv \frac{i}{2}(K^1 + iK^2), \quad \hat{S}_0 \equiv \frac{i}{2}(D + iM^{12}), \quad \hat{S}_+ \equiv \frac{i}{2}(P^1 - iP^2)$$

$$[\hat{S}_0, \hat{S}_\pm] = \pm \hat{S}_\pm, \quad \frac{1}{2}[\hat{S}_+, \hat{S}_-] = \hat{S}_0,$$

$$[\hat{S}_-, \hat{U}(z, \bar{z})] = z^2 \partial_z \hat{U}(z, \bar{z}), \quad [\hat{S}_0, \hat{U}(z, \bar{z})] = z \partial_z \hat{U}(z, \bar{z}), \quad [\hat{S}_+, \hat{U}(z, \bar{z})] = -\partial_z \hat{U}(z, \bar{z})$$

$$z \equiv z^1 + iz^2, \quad \bar{z} \equiv z^1 - iz^2, \quad U(z_\perp) = U(z, \bar{z})$$

## SL(2,C) for Wilson lines

$$\hat{S}_- \equiv \frac{i}{2}(K^1 + iK^2), \quad \hat{S}_0 \equiv \frac{i}{2}(D + iM^{12}), \quad \hat{S}_+ \equiv \frac{i}{2}(P^1 - iP^2)$$

$$[\hat{S}_0, \hat{S}_\pm] = \pm \hat{S}_\pm, \quad \frac{1}{2}[\hat{S}_+, \hat{S}_-] = \hat{S}_0,$$

$$[\hat{S}_-, \hat{U}(z, \bar{z})] = z^2 \partial_z \hat{U}(z, \bar{z}), \quad [\hat{S}_0, \hat{U}(z, \bar{z})] = z \partial_z \hat{U}(z, \bar{z}), \quad [\hat{S}_+, \hat{U}(z, \bar{z})] = -\partial_z \hat{U}(z, \bar{z})$$

$$z \equiv z^1 + iz^2, \quad \bar{z} \equiv z^1 - iz^2, \quad U(z_\perp) = U(z, \bar{z})$$

## Conformal invariance of the evolution kernel

$$\frac{d}{d\eta} [\hat{S}_-, \text{Tr}\{U_x U_y^\dagger\}] = \frac{\alpha_s N_c}{2\pi^2} \int dz K(x, y, z) [\hat{S}_-, \text{Tr}\{U_x U_y^\dagger\} \text{Tr}\{U_x U_y^\dagger\}]$$
$$\Rightarrow \left[ x^2 \frac{\partial}{\partial x} + y^2 \frac{\partial}{\partial y} + z^2 \frac{\partial}{\partial z} \right] K(x, y, z) = 0$$

In a conformal theory the amplitude of **4-point functions** depends on two conformal ratios which can be chosen as

$$R = \frac{(x - x')^2 (y - y')^2}{(x - y)^2 (x' - y')^2} \xrightarrow{\text{Regge limit}} 0$$

$$r = R \left[ 1 - \frac{(x - y')^2 (y - x')^2}{(x - x')^2 (y - y')^2} + \frac{1}{R} \right]^2 \xrightarrow{\text{Regge limit}} \text{fixed}$$

Cornalba (2007)

Provides general structure of the 4-point function in the regge limit as an integral over one real variable  $\nu$

$$\langle \{ \mathcal{S}(z_1^-, z_{1\perp}) \mathcal{S}(z_2^-, z_{2\perp}) \mathcal{S}(x^+, x_\perp) \mathcal{S}(y^+, y_\perp) \} \rangle = \int d\nu \Phi(r, \nu) F(\nu) R^{\frac{1}{2}\omega(\nu)}$$


$$\mathcal{S} \equiv \frac{4\pi^2\sqrt{2}}{\sqrt{N_c^2-1}} \text{Tr}\{Z^2\} \quad (Z = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)) \text{ renorm-invariant operator}$$

- $\omega(\nu) \equiv \omega(0, \nu)$  is the pomeron intercept.
- $F(\nu)$  is the “pomeron residue”.
- $\Phi(r, \nu)$  some function of  $\nu$  and the conformal ratio  $r$ .

Explicit calculation of the 4-point function in the Regge limit at NLL0




Factorization in rapidity




4-point function at NLO in the Regge limit using conformal composite dipole in Wilson line.

Factorization in rapidity



- $F(\nu)$  Pomeron residue (Impact factor) at NLO in  $\mathcal{N}=4$  SYM and QCD  
I. Balitsky and G.A.C. (2009)
- NLO Pomeron intercept  $\omega(\nu)$ 
  - QCD: Fadin-Lipatov (1998) and I. Balitsky and G.A.C (2007)
  - $\mathcal{N}=4$ : Lipatov-Kotikov (2000) and I. Balitsky and G.A.C. (2008)

## Factorization in rapidity



$$\begin{aligned}
 & \langle T\{\mathcal{S}(x)\mathcal{S}^\dagger(y)\mathcal{S}(x')\mathcal{S}^\dagger(y')\} \rangle \\
 &= \int d^2 z_{1\perp} d^2 z_{2\perp} d^2 z'_{1\perp} d^2 z'_{2\perp} \text{IF}^{a_0}(x, y; z_1, z_2) [\text{DD}]^{a_0, b_0}(z_1, z_2; z'_1, z'_2) \text{IF}^{b_0}(x', y'; z'_1, z'_2)
 \end{aligned}$$


in QCD: NLO  $\gamma^* \gamma^*$  cross section G.A.C. and Yu. Kovchegov (2015)



# Particle production in hadron-hadron collisions

## Rapidity Factorization

I. Balitsky and G.A.C. (2008),  
I. Balitsky and A. Tarasov (2015)



- Projectile fields:  $k^- < \sigma_a \Rightarrow$  Projectile TMD
- Central fields: coefficient functions
- Target fields:  $k^+ < \sigma_b \Rightarrow$  Target TMD

$$\begin{aligned}
 & \langle p_A, p_B | g^2 F_{\mu\nu}^a F^{a\mu\nu}(z_1) g^2 F_{\lambda\rho}^b F^{b\lambda\rho}(z_2) | p_A, p_B \rangle \\
 &= \frac{1}{N_c^2 - 1} \langle p_A | \tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) | p_A \rangle^{\sigma_A} \langle p_B | \mathcal{O}^{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) | p_B \rangle^{\sigma_B} + \dots
 \end{aligned}$$

## Gluon TMD operator

$$\mathcal{O}_{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) = \mathcal{F}_i^a(z_1) [z_1 - \infty n, z_2 - \infty n]^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^- = z_2^- = 0},$$

$$\tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) = \mathcal{F}_i^a(z_1) [z_1 - \infty n', z_2 - \infty n']^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^+ = z_2^+ = 0},$$

$$(\mathcal{F}^{i,a}(z_\perp, z^+))^\sigma \equiv g F^{-i,m}(z) [P e^{ig \int_{-\infty}^{z^+} dz^+ A^{-,\sigma}(up_1 + x_\perp)}]^{ma},$$

## Gluon TMD operator

$$\mathcal{O}_{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) = \mathcal{F}_i^a(z_1)[z_1 - \infty n, z_2 - \infty n]^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^- = z_2^- = 0},$$

$$\tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) = \mathcal{F}_i^a(z_1)[z_1 - \infty n', z_2 - \infty n']^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^+ = z_2^+ = 0},$$

$$(\mathcal{F}^{i,a}(z_\perp, z^+))^\sigma \equiv g F^{-i,m}(z) [\mathbf{P} e^{ig \int_{-\infty}^{z^+} dz^+ A^{-,\sigma}(up_1 + x_\perp)}]^{ma},$$

$$A_\mu^\sigma(x) = \int \frac{d^4 k}{16\pi^4} \theta\left(\frac{\sigma\sqrt{2}}{z_{12\perp}} - |k^+|\right) e^{-ik \cdot x} A_\mu(k)$$

$\frac{\sigma\sqrt{2}}{z_{12\perp}}$  **cutoff preserving conformal invariance**

$$[x, y] \equiv \mathbf{P} e^{ig \int du (x-y)^\mu A_\mu(ux + (1-u)y)}$$

## Gluon TMD operator

$$\mathcal{O}_{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) = \mathcal{F}_i^a(z_1)[z_1 - \infty n, z_2 - \infty n]^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^- = z_2^- = 0},$$

$$\tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) = \mathcal{F}_i^a(z_1)[z_1 - \infty n', z_2 - \infty n']^{ab} \mathcal{F}_j^b(z_2) \Big|_{z_1^+ = z_2^+ = 0},$$

TMDs are invariant under the inversion

$$\begin{aligned} \mathcal{F}_i^m(z_\perp, z^+) &= F^{-i,n}(z^+, z_\perp)[z^+, z^+ - \infty n]^{nm} \\ &\rightarrow F^{-i,n}\left(\frac{z^+}{z_\perp^2}, \frac{z_\perp}{z_\perp^2}\right) \left[\frac{z^+}{z_\perp^2}, \frac{z^+}{z_\perp^2} - \infty n\right]^{nm} = \mathcal{F}_i^m(z'_\perp, z'^+) \end{aligned}$$

We need the full conformal group of TMD operators.

Conformal group has 15 generators: Poincare + Dilatation + Special conformal transformation (inversion + shift + inversion)

$$i[M_{\mu\nu}, M_{\alpha\beta}] = g_{\mu\alpha}M_{\nu\beta} + g_{\nu\beta}M_{\mu\alpha} - g_{\mu\beta}M_{\nu\alpha} - g_{\nu\alpha}M_{\mu\beta}$$

$$i[M_{\alpha\beta}, P_{\mu}] = g_{\alpha\mu}P_{\beta} - g_{\beta\mu}P_{\alpha}$$

$$i[M_{\alpha\beta}, K_{\mu}] = g_{\alpha\mu}K_{\beta} - g_{\beta\mu}K_{\alpha}$$

$$i[D, P_{\mu}] = P_{\mu}, \quad i[D, K_{\mu}] = -K_{\mu}, \quad i[K_{\mu}, P_{\nu}] = 2(g_{\mu\nu}D + M_{\mu\nu})$$

Action on scalar field of canonical dimension  $\Delta$ ;

$$i[D, \Phi(x)] = (x^{\alpha}\partial_{\alpha} + \Delta)\Phi(x)$$

$$i[K^{\mu}, \Phi(x)] = (2x^{\mu}x^{\alpha}\partial_{\alpha} - x^2\partial^{\mu} + 2\Delta x^{\mu})\Phi(x)$$

quantum correction:  $\Delta \rightarrow \Delta + \text{anomalous}$

# Conformal invariance of TMD operator

Conformal  $SO(2,4)$  group has 15 generators

TMD operator transform covariantly under 11 generators:

## TMD Conformal group

$$P^i, P^-, M^{12}, M^{-i}, D, K^i, K^-, M^{-+}$$

Generators  $P^+, K^+, M^{+i}$  do not preserve the form of  $\mathcal{F}^{-j}$ .

## Action of conformal generators on TMD operators

$$-i\mathbf{P}^i \mathcal{F}^{-j}(x^+, x_\perp) = \partial^i \mathcal{F}^{-j}(x^+, x_\perp), \quad -i\mathbf{P}^- \mathcal{F}^{-j}(x^+, x_\perp) = \frac{\partial}{\partial x^+} \mathcal{F}^{-j}(x^+, x_\perp),$$

$$-i\mathbf{M}^{-i} \mathcal{F}^{-j}(x^+, x_\perp) = -x^i \frac{\partial}{\partial x^+} \mathcal{F}^{-j}(x^+, x_\perp),$$

$$-i\mathbf{D} \mathcal{F}^{-i}(x^+, x_\perp) = (x^+ \frac{\partial}{\partial x^+} + x^k \frac{\partial}{\partial x^k} + 2) \mathcal{F}^{-i}(x^+, x_\perp),$$

$$-i\mathbf{M}^{ij} \mathcal{F}^{-k}(x^+, x_\perp) = (x^i \partial^j - x^j \partial^i) \mathcal{F}^{-k}(x^+, x_\perp) + g^{ik} \mathcal{F}^{-j}(x^+, x_\perp) - g^{jk} \mathcal{F}^{-i}(x^+, x_\perp),$$

$$\begin{aligned} -\mathbf{K}^i \mathcal{F}^{-j}(x^+, x_\perp) &= 2x^i (x^+ \frac{\partial}{\partial x^+} + x^k \frac{\partial}{\partial x^k} + 2) \mathcal{F}^{-j}(x^+, x_\perp) + x_\perp^2 \frac{\partial}{\partial x^i} \mathcal{F}^{-j}(x^+, x_\perp) \\ &\quad - 2x^j \mathcal{F}^{-i}(x^+, x_\perp) + 2g^{ij} x_i \mathcal{F}^{-i}(x^+, x_\perp), \end{aligned}$$

$$-i\mathbf{K}^- \mathcal{F}^{-j}(x^+, x_\perp) = x_\perp^2 \frac{\partial}{\partial x^+} \mathcal{F}^{-j}(x^+, x_\perp),$$

$$-i\mathbf{M}^{+-} \mathcal{F}^{-i}(x^+, x_\perp) = (x^+ \frac{\partial}{\partial x^+} + 1) \mathcal{F}^{-i}(x^+, x_\perp)$$

## Conformal TMD group in the embedding formalism

Conformal TMD group in the 6-dim space;  $g^{-2,-2} = 1$ ,  $g^{-1,-1} = -1$

Conf. Transf. are Lorentz transf. of light-rays  $\left(\frac{1-x^2}{2}, \frac{1+x^2}{2}, x_\mu\right)$  in 6-dim space with metric  $(1, -1, 1, -1, -1, -1)$

$$L_{\mu\nu} \equiv M_{\mu\nu}, \quad L_{-2,\mu} \equiv \frac{1}{2}(P_\mu - K_\mu), \quad L_{-1,\mu} \equiv \frac{1}{2}(P_\mu + K_\mu), \quad L_{-2,-1} \equiv D$$

$$i[L_{ab}, L_{mn}] = g_{ma}L_{nb} + g_{nb}L_{ma} - g_{mb}L_{na} - g_{na}L_{mb}$$

Define

$$L_{mn} \equiv \mathbb{M}_{mn} \quad L_{-n} \equiv \mathbb{P}_n \quad L_{+-} \equiv \mathbb{D}$$

with  $m, n, l = -2, -1, 1, 2$ , get usual Poincare generators  $\mathbb{M}_{mn}, \mathbb{P}_n$  and the "Dilatation"  $\mathbb{D}$  in 4-dim sub-space orthogonal to the physical "+" and "-" directions




# Rapidity evolution at leading in Sudakov region

$$m_X^2 = q^2 \equiv Q^2 \quad \text{Sudakov region: } Q \gg q_\perp \gg 1\text{Gev}$$

Sudakov region in coord. space:  $z_{12\parallel}^2 \equiv 2z_{12}^- z_{12}^+ \ll z_{12\perp}^2$

Typical LO diagrams



To calculate these diagrams the approximation is:  $k^+ \gg \frac{z_{12}^+}{z_{12\perp}^2}$

**Sudakov evolution:** transverse separation between the gluon operators  $\mathcal{F}_i$  and  $\mathcal{F}_j$  does not change **while the longitudinal one increases.**

$$\mathcal{O}^{\sigma_2}(z_1^+, z_2^+) = \frac{\alpha_s N_c}{2\pi} \int_{\frac{\sigma_1 \sqrt{2}}{z_{12\perp}}^+}^{\frac{\sigma_2 \sqrt{2}}{z_{12\perp}}^+} \frac{dk^+}{k^+} K \mathcal{O}^{\sigma_1}(z_1^+, z_2^+)$$

where the kernel  $K$  is given by

$$\begin{aligned} K \mathcal{O}(z_1^+, z_2^+) &= \mathcal{O}(z_1^+, z_2^+) \int_{-\infty}^{z_1^+} \frac{dz^+}{z_2^+ - z^+} e^{-i \frac{z_{12\perp} \sigma}{\sqrt{2}(z_2 - z)^+}} + \mathcal{O}(z_1^+, z_2^+) \int_{-\infty}^{z_2^+} \frac{dz^+}{z_1^+ - z^+} e^{i \frac{z_{12\perp} \sigma}{\sqrt{2}(z_1 - z)^+}} \\ &- \int_{-\infty}^{z_1^+} dz^+ \frac{\mathcal{O}(z_1^+, z_2^+) - \mathcal{O}(z^+, z_2^+)}{z_1^+ - z^+} - \int_{-\infty}^{z_2^+} dz^+ \frac{\mathcal{O}(z_1^+, z_2^+) - \mathcal{O}(z_1^+, z^+)}{z_2^+ - z^+} \end{aligned}$$

To solve the evolution equation, perform Fourier transform

$$\begin{aligned}
 Ke^{-ik^-z_1^+ + ik'^-z_2^+} &= \left[ -2 \ln \sigma z_{12\perp} - \ln(ik^-) \right. \\
 &\quad \left. - \ln(-ik'^-) + \ln 2 - 4\gamma_E + O\left(\frac{z_{12}^+}{z_{12\perp}\sigma}\right) \right] e^{-ik^-z_1^+ + ik'^-z_2^+}
 \end{aligned}$$

Result ( $\bar{\alpha}_s = \frac{\alpha_s N_c}{4\pi}$ )

I. Balitsky and G.A.C (2019)

$$\begin{aligned}
 \mathcal{O}^{\sigma_2}(z_1^+, z_2^+) &= e^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1} [\ln \sigma_1 \sigma_2 + 4\gamma_E - \ln 2]} \int dz_1'^+ dz_2'^+ \mathcal{O}^{\sigma_1}(z_1'^+, z_2'^+) z_{12\perp}^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} \\
 &\times \frac{1}{4\pi^2} \left[ \frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_1^+ - z_1'^+ + i\epsilon)^{1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} + c.c.} \right] \left[ \frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_2^+ - z_2'^+ + i\epsilon)^{1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} + c.c.} \right]
 \end{aligned}$$

Result ( $\bar{\alpha}_s = \frac{\alpha_s N_c}{4\pi}$ )

I. Balitsky and G.A.C (2019)

$$\begin{aligned} \mathcal{O}^{\sigma_2}(z_1^+, z_2^+) &= e^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1} [\ln \sigma_1 \sigma_2 + 4\gamma_E - \ln 2]} \int dz_1'^+ dz_2'^+ \mathcal{O}^{\sigma_1}(z_1'^+, z_2'^+) z_{12\perp}^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} \\ &\times \frac{1}{4\pi^2} \left[ \frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_1^+ - z_1'^+ + i\epsilon)^{1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}}} + c.c. \right] \left[ \frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_2^+ - z_2'^+ + i\epsilon)^{1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}}} + c.c. \right] \end{aligned}$$

Result transform covariantly under the TMD-conformal group generators except the Lorentz boost  $M^{+-}$  which is the generator of the evolution equation: The Lorentz boost in  $z$  direction changes the cutoffs for the evolution.

Sudakov-region result is applicable in the region between:

$$\sigma_2 = \sigma_B = \frac{z_{12\perp}}{z_{12}^- \sqrt{2}} \quad \text{and} \quad \sigma_1 = \frac{z_{12}^+ \sqrt{2}}{z_{12\perp}}$$

# Conformal invariance of the TMD matrix element

Sudakov-region result is applicable in the region between:

$$\sigma_2 = \sigma_B = \frac{z_{12\perp}}{z_{12}^- \sqrt{2}} \quad \text{and} \quad \sigma_1 = \frac{z_{12}^+ \sqrt{2}}{z_{12\perp}}$$

Lorentz boost:  $z^+ \rightarrow \lambda z^+$ ,  $z^- \rightarrow \frac{1}{\lambda} z^-$

$$\blacksquare \langle p_B | \mathcal{O} | p_B \rangle \rightarrow \langle p_B | \mathcal{O} | p_B \rangle \exp\left\{4\lambda \bar{\alpha}_s \ln \frac{z_{12\parallel}^2}{z_{12\perp}^2}\right\} \quad \text{Target}$$

$$\blacksquare \langle p_A | \tilde{\mathcal{O}} | p_A \rangle \rightarrow \langle p_A | \tilde{\mathcal{O}} | p_A \rangle \exp\left\{-4\lambda \bar{\alpha}_s \ln \frac{z_{12\parallel}^2}{z_{12\perp}^2}\right\} \quad \text{Projectile}$$

So the amplitude is invariant:

$$\begin{aligned} & \langle p_A, p_B | g^2 F_{\mu\nu}^a F^{a\mu\nu}(z_1) g^2 F_{\lambda\rho}^b F^{b\lambda\rho}(z_2) | p_A, p_B \rangle \\ &= \frac{1}{N_c^2 - 1} \langle p_A | \tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) | p_A \rangle^{\sigma_A} \langle p_B | \mathcal{O}^{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) | p_B \rangle^{\sigma_B} \end{aligned}$$

# Comparing with conventional TMD analysis

Evolution for generalized TMD  $\xi = \frac{p_B - p'_B}{\sqrt{2s}}$

$$D^\sigma(x_B, \xi) = \int dz^+ e^{-ix_B \sqrt{\frac{s}{2}} z^+} \langle p'_B | \mathcal{O}^\sigma \left( -\frac{z^+}{2}, \frac{z^+}{2} \right) | p_B \rangle$$

From our result we get

$$\frac{D^{\sigma_2}(x, \xi)}{D^{\sigma_1}(x, \xi)} = e^{-2\bar{\alpha}_s \ln \frac{q_2^2}{q_1^2} [\ln \sigma_2 \sigma_1 (x^2 - \xi^2) s z_{12\perp}^2 + 4\gamma_E - 2 \ln 2]}$$

For usual TMD at  $\xi = 0$  with the limits of Sudakov evolution one obtains

$$\frac{D^{\sigma_2}(x, q_\perp)}{D^{\sigma_1}(x, q_\perp)} = e^{-2\bar{\alpha}_s \ln \frac{q_2^2}{q_1^2} [\ln \frac{q_2^2}{q_1^2} + 4\gamma_E - 2 \ln 2]}$$

- Coincides with usual one-loop evolution of TMDs up to replacement  $4\gamma_E - 2 \ln 2 \rightarrow 4\gamma_E - 4 \ln 2$ .
- constant depends on the way of cutting  $k^+$ -integration which should be coordinated with the cutoffs in the “coefficient function”  $\sigma(ff \rightarrow H)$ 
  - The discrepancy is just like using two different schemes for usual renormalization.

Sudakov:  $Q \gg q_{\perp}$ ; in coord. space  $(x - y)_{\perp}^2 \gg (x - y)_{\parallel}^2$

■  $x_B \sim 1$  and  $q_{\perp} \sim m_N$

- The relative energy between Wilson-line operators  $\mathcal{F}$  and target nucleon at the final point of the evolution is  $\sim m_N^2$  so one should use phenomenological models of TMDs with this low rapidity cutoff as a starting point of the evolution.

■  $x_B \ll 1$

- This relative energy is within  $\frac{q_{\perp}^2}{x_B s} > \sigma > \frac{m_N^2}{s}$  beyond Sudakov region into the low-x region: the TMD operator, known as Weiczsäcker-Williams distribution, will produce a of color dipoles as a result of the non-linear evolution. **The transition between Sudakov region and small-x region is described by rather complicated interpolation formula.** In coordinate space this means the study of the operator  $\mathcal{O}$  at  $z_{\parallel}^2 \sim z_{\perp}^2$ . **Conformal invariance may help us obtain the TMD evolution in that region.**



- The conformal TMD group has been obtained: it is made out of 11 generators of the full conformal group.
- We obtained conformal evolution of gluon and quark ( $N_c \rightarrow C_F$ ) TMDs in the Sudakov region.

## Outlook

- Conformal properties of TMD evolution in the small-x region.
- Conformal evolution for all  $x_B$ .
- The plan is to perform the calculation for 6 point function similarly to the one done for the 4 point function in  $\mathcal{N}=4$  and in in QCD.