

O3.401 Positron confinement exceeding 1s in a magnetic dipole trap

Wednesday, 10 July 2019 12:10 (15 minutes)

See the full abstract here <http://ocs.ciemat.es/EPS2019ABS/pdf/O3.401.pdf>

Magnetic dipole fields have excellent confinement properties for charged particle ensembles with an arbitrary degree of neutrality. One idea to create a magnetized low-energy electron-positron pair plasma pursued by the APEX/PAX collaboration relies on the initial formation of a non-neutral plasma in the confining magnetic field and hence the dipole geometry is a promising candidate to study such a system. While electron injection and confinement have been readily achieved in a magnetic dipole trap the positron counterpart is more challenging due to the limited positron flux provided even by world class positron sources such as NEPOMUC (Neutron-Induced Positron Source Munich).

In this contribution we present results from recent positron confinement experiments conducted with a supported magnetic dipole trap at the NEPOMUC facility. After suppressing parallel losses onto the magnet surface by applying a positive electrostatic bias to the magnet we have observed positron confinement in a magnetic dipole field in excess of 1 s. Supported by single-particle simulations we identify transport due to scattering off of neutrals as the dominant mechanism limiting confinement. The importance of the results for the electron-positron pair plasma experiment will be discussed.

pppo

Presenter: HORN-STANJA, J. (EPS 2019)

Session Classification: BSAP

Track Classification: BSAP