

# The Dark Matter Phase Diagram: Thermal Relics between keV and TeV

Raffaele Tito D'Agnolo - IPhT Saclay



$10^{-22}$  eV  $m_\chi$   $10^{48}$  GeV

$\mathcal{O}(10^{-50})$



$\mathcal{O}(10^{-50})$

$\alpha_{\text{self}}$

$\alpha_{\text{SM}}$


$\mathcal{O}(1)$



$\mathcal{O}(1)$

# LDM INTRO FROM 6 YEARS AGO

We have looked for WIMPs for decades...  
We have looked for SUSY for decades...



We have not found anything (yet)



Here is something new (and motivated?)

# THE UNUSUAL APPEAL OF THE WIMP

1. Insensitive to initial conditions (no assumptions on the UV)

$$T > m_{\text{DM}}$$

DM in  
Kinetic and  
Chemical Eq.  
with a bath at  
 $T = T_{\text{SM}}$

# THE UNUSUAL APPEAL OF THE WIMP

## 1. Insensitive to initial conditions (no assumptions on the UV)

$$T > m_{\text{DM}}$$

DM in  
Kinetic and  
Chemical Eq.  
with a bath at  
 $T = T_{\text{SM}}$

**A coupling in the dark sector [or non-standard cosmology] is needed to remove excess DM abundance**

Cannibal, Secluded, Forbidden  $v1$ , ELDER, Co-decaying, sub-MeV, Inflatable, asymmetric, WIMPs with GUTs, ...

**Potential Experimental Target: Decay to SM radiation**

$$\alpha_{\text{eff}} \gtrsim 10^{-20}$$

# THE UNUSUAL APPEAL OF THE WIMP

## 2. The coupling to the Standard Model fixes the relic density

$$T > m_{\text{DM}}$$

DM in  
Kinetic and  
Chemical Eq.  
with a bath at  
 $T = T_{\text{SM}}$

**A coupling in the dark sector is needed to remove excess DM abundance**

Cannibal, Secluded, Forbidden v1, ELDER, Co-decaying, sub-MeV, Inflatable, asymmetric, WIMPs with GUTs, ...

$$\alpha_{\text{eff}} \gtrsim 10^{-20}$$

**A coupling to the SM removes excess DM abundance**



WIMP [=WIMPlless, Semi-Annihilation, Inelastic, non-Abelian] + ?

**Irreducible Experimental Target: Relic Density**

$$\alpha_{\text{eff}} \gtrsim 10^{-7}$$

# WIMPs AND TARGETS

$\chi$  = Dark Matter





**Semi-Annihilation** [D'Eramo, Thaler '10], **Inelastic** [Smith, Weiner '01], **WIMPIless Miracle** [Feng, Kumar '08], **Heavy Mediators** [Bohem, Fayet '07], **Non-Abelian DM** [Buen-Abad, Marques-Tavares, Schmaltz '15], and many more with similar parametrics

# SCATTERING WITH A PARTNER

[**RTD**, Ruderman, Pappadopulo '17]


[**RTD**, Ruderman, Pappadopulo, Wang, '19]



**Resonant** [Feng, Smolinsky '17]  
Similar in this plane


# HOW IT WORKS: COSCATTERING

$\chi$  = Dark Matter




$$\frac{dn_{\chi}}{dt} + 3Hn_{\chi} = -\langle\sigma v\rangle [n_{\chi} - n_{\chi}^{eq}] n^{eq}$$

**WIMP**



**COSCATTERING**





$$\sim e^{-m_{\chi}/T_F}$$

# ANNIHILATIONS WITH A PARTNER

[Griest, Seckel '89]


[**RTD**, Mondino, Ruderman, Wang '18]




$\chi$  = Dark Matter

# HOW IT WORKS: COANNIHILATION


$\chi$  = Dark Matter



$$\frac{dn_{\chi}}{dt} + 3Hn_{\chi} = -\langle\sigma v\rangle \left[ n_{\chi}n_{\chi'} - n_{\chi}^{eq}n_{\chi'}^{eq} \right]$$



**COANNIHILATION**




$$\sim e^{-\frac{m_{\chi'} - m_{\chi}}{T_F}}$$

[Griest, Seckel '89], [**RTD**, Mondino, Ruderman, Wang '18]


# TOWARDS A PHASE DIAGRAM OF DARK MATTER

Coscattering, Coannihilation and WIMP phases are all continuously connected and there is more...



# TOWARDS A PHASE DIAGRAM OF DARK MATTER

Coscattering, Coannihilation and WIMP phases are all continuously connected and there is more...



# COSCATTERING



# INGREDIENTS BEYOND WIMPS

$\chi$  = Dark Matter

$\psi$  = Strongly interacting Dark Matter partner

# INGREDIENTS BEYOND WIMPs

$\chi$  = Dark Matter

$\psi$  = Strongly interacting Dark Matter partner

Examples: Gluino-Bino, Wino-Bino, ...

# INGREDIENTS BEYOND WIMPS

$\chi$  = Dark Matter

$\psi$  = Strongly interacting Dark Matter partner

$$m_\psi \gtrsim m_\chi$$


$$\Delta \equiv \frac{m_\psi - m_\chi}{m_\chi}$$

# A GENERAL REALIZATION OF THE MECHANISM


$x$

$\psi$

SM  
THERMAL BATH





# A GENERAL REALIZATION OF THE MECHANISM





# A GENERAL REALIZATION OF THE MECHANISM

$$\chi \leftrightarrow \psi$$



$$\chi, \psi \leftrightarrow \text{SM}$$




# BOLTZMANN "NDA"


Relevant timescale

$$n_\chi \langle \sigma v \rangle_F \approx H_F$$

$$n_\chi(T_F) \sim e^{-m_\chi/T_F}$$




$$\sim \delta^2 T_F^3 \sigma_0$$




$$\sim (T_F m_\psi)^{3/2} e^{-m_\psi/T_F} \sigma_0$$

# TIMESCALES OF FREEZE-OUT: COSCATTERING




# COSCATTERING: A REMINDER

$\chi = \text{Dark Matter}$




$$\frac{dn_\chi}{dt} + 3Hn_\chi = -\langle\sigma v\rangle [n_\chi - n_\chi^{eq}] n^{eq}$$

**WIMP**




**COSCATTERING**



$$\sim e^{-m_\chi/T_F}$$

# PHENOMENOLOGY




[RTD,  
Ruderman,  
Pappadopulo,  
Wang, '19]

# CONCLUSION

- Thermal Freeze-out has “built-in” exponential hierarchies that organize **DM phenomenology** in a rich **phase diagram**
- Take your favourite WIMP model and add a DM partner: the **couplings to SM** that give the right abundance are **exponentially reduced**
- Experimental lesson: **don't stop at thermal targets!**

# SARA: Superconducting Apparatus for Resonant Axion detection



W/ A. Berlin, S. Ellis, P. Schuster, N. Toro, K. Zhou | C. Nantista, J. Nielson, S. Tantawi

TEASER: VERY PRELIMINARY

# BEHIND THE WIMP

- Change the coupling
  - WIMPless, Heavy Mediators, Freeze-in, ...
- Change the kinematics of freeze-out
  - Forbidden channel DM, Coannihilation, Coscattering, Impeded DM, ...
- Change the thermal history of the DM sector
  - Cannibal DM, Inflatable DM, ...
- Change the interaction relevant for freeze-out
  - SIMP, ELDER, Coscattering, ...