

# Experimental proposal for MDM/ EDM of charm (beauty) baryons in LHCb

---

A. Merli

Università degli Studi di Milano

Workshop on electromagnetic dipole moments of unstable particles  
Milano, 3rd October 2019



# Electromagnetic dipole moments

$\delta$  = electric dipole moment (EDM)      $\mu$  = magnetic dipole moment (MDM)

- Classic systems

$$\delta = \int \mathbf{r} \rho(\mathbf{r}) d^3 r \quad \mu = \int \mathbf{r} \times \mathbf{j}(\mathbf{r}) d^3 r$$

- Quantum systems


$$\delta = d\mu_N \frac{S}{2} \quad \mu = g\mu_N \frac{S}{2}$$

- Hamiltonian

$$H = -\delta \cdot \mathbf{E} - \mu \cdot \mathbf{B}$$

Time reversal, Parity:

$$\begin{array}{l} \xrightarrow{T} \\ \xrightarrow{P} \end{array} \quad +\delta \cdot \mathbf{E} - \mu \cdot \mathbf{B}$$



The EDM violates  $T$  and  $P$  and via CPT theorem, violates  $CP$

# Physics motivation for EDM

---

- ▶ CP violation (CPV) is a necessary condition for baryogenesis:
- ▶ CPV in weak interactions via CKM mechanism in the SM is too small to explain the absence of antimatter in the Universe
- ▶ CPV in strong interactions allowed in the SM. Stringent experimental limit from neutron EDM



New CPV sources are expected to exist

# EDM as a possible solution for baryogenesis


- ▶ EDM of fundamental particles from the structure of quarks and gluons, and processes with photon and flavour-diagonal coupling
- ▶ A measurement of a heavy baryon EDM is directly sensitive to:



Charm EDM in Standard Model  $\sim 10^{-32}$  ecm

Charm EDM in new physics  $\sim 10^{-17}$  ecm

EPJC 77 (2017), 102



**EDM** observation = clear sign of **new physics**



# Physics motivation for MDM

---

- ▶ Experimental anchor points for tests of low-energy QCD models, related to non-perturbative QCD dynamics
- ▶ Test of quark substructure
- ▶ Measurement of MDM of particles and antiparticles would allow a test of CPT symmetry

# Channeling in bent crystals

- ▶ Potential well between crystal planes
- ▶ Incident positive charged particle can be trapped if parallel to crystal plane (within **few  $\mu\text{rad}$** )
- ▶ Well understood phenomenon (Lindhard 1965)
- ▶ Effect of the **bent crystals**:
  - ▶ **Steer** high energy particle beams
  - ▶ Induce **spin precession**



# Spin precession in bent crystals

- ▶ Firstly predicted by **Baryshevsky (1979)**
- ▶ Determine particle gyromagnetic factor from TBMT equation

V.L. Lyuboshits, Sov. J. Nucl. Phys. 31 (1980) 509

$\Phi$  = spin rotation angle  
 $\theta_C$  = crystal bending angle  $\sim 10^{-2}$  rad  
 $g$  = gyromagnetic factor  
 $\gamma$  = Lorentz boost  $\sim 4-5 \cdot 10^2$

$$\Phi = \frac{g-2}{2} \gamma \theta_C$$





Fig. 1. Spin rotation in a bent crystal.

- ▶ Before decay the baryons experience a huge electric field in the crystal


- ▶ **MDM** and **EDM** precession in the limit  $\gamma \gg 1, d \ll g - 2$

EPJC 77 (2017), 181

$$S_x = S_0 \frac{d}{g-2} (\cos \Phi - 1)$$



# Experimental proposal



$p$  extraction


Heavy baryon polarised production

Channeling  
Spin precession

Event reconstruction





# LHCb detector



# Simulation studies





- ▶ Tungsten (W) 5mm fixed target + bent crystal positioned at 116cm before the interaction point



- ▶ Use EPOS for fixed target minimum bias events, PYTHIA for baryons produced in pW hard collisions
- ▶ Signal reconstruction and background rejection studied using LHCb full simulation

# Detector occupancy


- ▶ Flux =  $1 \times 10^8$  p/s,  $v = 0.181$  in **W target**,  $v = 0.629$  in **Ge crystals**



- ▶ Occupancies for **fixed-target events** under control wrt generic **bb events** ( $v=7.6$ )


# Identification of signal events

- ▶ About  $10^{-4}$   $\Lambda_c^+$  produced in the target are **channeled** in the bent crystal




- ▶ Use PV to identify  $\Lambda_c^+$  produced in W target, and  $\Lambda_c^+$  vertex helps to identify decays outside of the crystal (max spin precession)
- ▶  $\Lambda_c^+$  angle determined by crystal bending angle, e.g.  $\theta_C = 15 \text{ mrad}$
- ▶ Channeled baryons have **high momentum**  $\gtrsim 1 \text{ TeV}/c$

# $\Lambda_c^+$ momentum distribution




- ▶ At production (top)
- ▶ After channeling and  $p > 800$  GeV/c

Si,  $L \sim 7$  cm,  $\theta_c \sim 14$  mrad





Ge,  $L \sim 5$  cm,  $\theta_c \sim 15$  mrad




# Reconstruction of signal events

- ▶ LHCb Upgrade performs well in reconstructing these events
- ▶  $\Lambda_c^+ \rightarrow pK^-\pi^+$  daughter particles ( $p > 300$  GeV) have reduced momentum resolution  $> 1\%$
- ▶ Invariant mass resolution 20 MeV is good enough for signal reconstruction and background rejection

Int. J. Mod. Phys. A 30, 1530022 (2015)




- ▶ Reconstruction independent on the proton flux



# Background rejection


- ▶ Rejection of unchanneled  $\Lambda_c^+$  produced in W target



- ▶ Background rejection  $10^{-7}$  level and signal efficiency 80%
- ▶ High momentum  $\Lambda_c^+$  most sensitive for EDM measurements


# Synergetic run with LHCb

- ▶ Synergetic running with LHCb feasible for small flux  $< 10^7$  p/s
- ▶ Simulated one PV in the target and  $v=7.6$  pp collisions
- ▶ The presence of the target doesn't impact the reconstruction of pp events





# Sensitivity on EDM



PoT: protons on target  
W target 5 mm thick

[EPJC 77 \(2017\), 828](#)

**S1**: configuration at LHCb

**S2**: dedicated experiment at LHC


Material of the crystal:

→ **Silicon**

→ **Germanium**

- ▶ Technique applies to all short-lived positive baryons
- ▶ Possibility to test new physics models

# Sensitivity on MDM



PoT: protons on target  
W target 5 mm thick

[EPJC 77 \(2017\), 828](#)

**S1**: configuration at LHCb

**S2**: dedicated experiment at LHC

Material of the crystal:

→ **Silicon**

→ **Germanium**

► First MDM measurements

# Conclusions

---

- ▶ Experimental **proposal for unique baryon EDM/MDM measurements** in LHCb was presented
- ▶ Those searches will extend the new physics discovery potential of LHC
- ▶ Synergetic runs with pp collisions feasible

# Acknowledgment


---

- ▶ Proponents within LHCb: S. Aiola, J. Fu, L. Henry, D. Marangotto, F. Martinez Vidal, A. Merli, N. Neri, P. Robbe, J. Ruiz Vidal
- ▶ References:
  - E. Bagli, L. Bandiera, G. Cavoto, V. Guidi, L. Henry, D. Marangotto, F. Martinez Vidal, A. Mazzolari, A. Merli, N. Neri, J. Ruiz Vidal, *Electromagnetic dipole moments of charged baryons with bent crystals at the LHC*, arXiv: 1708.08483 (2017), Eur. Phys. J. C **77** (2017) 828.
  - A.S. Fomin , A.Yu. Korchin, A. Stocchi, O.A. Bezshyyko, L. Burmistrov, S.P. Fomin, I.V. Kirillin, L. Massacrier , A. Natochii, P. Robbe, W. Scandale, N.F. Shul'ga, *Feasibility of measuring the magnetic dipole moments of the charm baryons at the LHC using bent crystals*, JHEP **1708** (2017) 120.
  - V. G. Baryshevsky, *On the search for the electric dipole moment of strange and charm baryons at LHC and parity violating (P) and time reversal (T) invariance violating spin rotation and dichroism in crystal*, arXiv:1708.09799 (2017).
  - L. Henry, D. Marangotto, F. Martinez Vidal, A. Merli, N. Neri, P. Robbe, J. Ruiz Vidal, CERN- LHCb-INT-2017-011, *Proposal to search for baryon EDMs with bent crystals at LHCb*.
  - F. J. Botella, L. M. Garcia Martin, D. Marangotto, F. Martinez Vidal, A. Merli, N. Neri, A. Oyanguren, J. Ruiz Vidal, *On the search for the electric dipole moment of strange and charm baryons at LHC*, Eur. Phys. J. C **77** (2017) 181.
  - V. G. Baryshevsky, *The possibility to measure the magnetic moments of short-lived particles (charm and beauty baryons) at LHC and FCC energies using the phenomenon of spin rotation in crystals*, Phys. Lett. **B757** (2016) 426.
  - L. Burmistrov, G. Calderini, Yu Ivanov, L. Massacrier, P. Robbe, W. Scandale, A. Stocchi, *Measurement of short living baryon magnetic moment using bent crystals at SPS and LHC*, CERN-SPSC-2016-030 ; SPSC-EOI-012.

---

# Back-up slides

# Channeling efficiency



- ▶ Channeling efficiency for  $\Lambda_c^+$  particles within Lindhard angle
- ▶ Total channelling efficiency: Lindhard angle, dechanneling,  $\Lambda_c^+$  decay flight:  $1 \cdot 10^{-5}$  (Si),  $4 \cdot 10^{-5}$  (Ge)

$$w(\theta_C, R) = \left(1 - \frac{R_c}{R}\right)^2 \exp\left(-\frac{\theta_C}{\theta_D \frac{R_c}{R} \left(1 - \frac{R_c}{R}\right)^2}\right)$$

- ▶ Parametrisation from Biryukov, Valery M. (et al.), *Crystal Channeling and Its Application at High-Energy Accelerators*, Springer Verlag (1997)

# Sensitivity to EDM/MDM

► Studies based on:

- $\Lambda_c^+$  from fixed-target (Pythia + EvtGen)
- Reconstruction, Decay flight efficiency (LHCb simulation)
- Channeling efficiency (parametrization)
- Fit to spin precession (pseudo experiments)

$$N_{\Lambda_c^+}^{\text{reco}} = N_{\Lambda_c^+} \mathcal{B}(\Lambda_c^+ \rightarrow f) \epsilon_{\text{CH}} \epsilon_{\text{DF}} \epsilon_{\text{det}}$$

$$\sigma(pp \rightarrow \Lambda_c^+ X) \approx 18.2 \mu\text{b}$$

$$|S_0| \approx 0.6$$

$$\epsilon_{\text{det}} \approx 20\% \quad \epsilon_{\text{DF}} \approx 10\%$$

$$\epsilon_{\text{ch}} \approx 10^{-4}$$

$$\frac{dN}{d\Omega} \propto 1 + \alpha_f S \cdot p$$


$$\alpha_{\Delta^{++}K^-} \approx -0.67$$

$$\sigma_d \approx \frac{g-2}{\alpha_f s_0 (\cos \Phi - 1)} \frac{1}{\sqrt{N_{\Lambda_c^+}^{\text{reco}}}}$$

$$\sigma_g \approx \frac{2}{\alpha_f s_0 \gamma \theta_C} \frac{1}{\sqrt{N_{\Lambda_c^+}^{\text{reco}}}}$$

# LHCb acceptance

- ▶ Channeled particles with crystals with bending angle  $< 14/15\text{mrad}$  has low reconstruction efficiency




- ▶ Dependence of the reconstruction efficiency over azimuthal angle due to the LHCb detector geometry



# Crystal optimisation

- ▶ Optimised sensitivity to EDM and MDM
- ▶ Channeling and reconstruction efficiency included



- ▶ Regions of minimal uncertainty of EDM (continuous line) and MDM (dotted line) defined as +20% uncertainty wrt the minimum (point marker)