



Muon g-2 Status

Mark Lancaster

Aim of experiment





If measure 1σ below BNL value



Motivation



arXiv:1908.07525



“.. it’s extremely sensitive to new physics. It’s still running, but if I were to put my money on something that would signal new physics, it’s the g-2 experiment at Fermilab. I think it’s really fascinating...”

$(g-2)_e$ vs $(g-2)_\mu$



arXiv.org > hep-ph > arXiv:1908.03607

High Energy Physics - Phenomenology

Explanation of electron and muon $g-2$ anomalies in the MSSM

Marcin Badziak, Kazuki Sakurai

(Submitted on 9 Aug 2019)

arXiv.org > hep-ph > arXiv:1907.08109

High Energy Physics - Phenomenology

$(g-2)_{\mu,e}$ and the ANITA anomalous events in a three-loop neutrino mass model

Mohammad Abdullah, Bhaskar Dutta, Sumit Ghosh, Tianjun Li



(Submitted on 18 Jul 2019)

arXiv.org > hep-ph > arXiv:1905.03789

High Energy Physics - Phenomenology

Combined explanations of $(g-2)_\mu$, $(g-2)_e$ and implications for a large muon $E\Gamma^M$

Andreas Crivellin, Martin Hoferichter




New physics in muon interactions ?




Physicists Finally Nail the Proton's Size, and Hope Dies


27 | A new measurement appears to have eliminated an anomaly that had captivated physicists for nearly a decade.



Experiment Overview



Experiment Overview




Methodology





140 ppb

$$a_\mu = \frac{\omega_a}{\tilde{\omega}_p} \frac{\mu_p}{\mu_e} \frac{m_\mu}{m_e} \frac{g_e}{2}$$

- 3 ppb : Hydrogen Maser
- 22 ppb : Muonium hyperfine splitting
- 0.00026 ppb : Electron g-2/QED






$$\frac{\omega_a}{\omega_p \otimes \rho(r)}$$





Monitoring the Field


- Fixed probes track field at top/bottom of vacuum chamber monitor field 24/7
- NMR trolley maps field where muons traverse every 2-3 days
- Digitizing FID signals





Absolute Field Calibration







Calorimeters




800 MSPS Digitizers



Straw Trackers





Commissioning Jun-17 → Mar-18



5 orders of magnitude improvement in muon flux
TDR envisaged 200 days of commissioning (June-17, Nov-17 → Mar-18)

Run-1





Raw data: x2 BNL **but** several different quad/kicker settings.
 Resulted in 7 datasets with approx. x1.4 BNL.


Run-2 aim: fewer datasets with constant conditions

Run-1 issues affecting integrated stats (& systematics/ease of analysis)

- kick was too low
- kicker had significant downtime
- significant number of quad sparks
- magnet downtime due to cryo purity issues



Such that fraction of days with $> 100\text{M e}^+$ was 57%.

The run-1/2 shutdown addressed these issues




Kicker ran very stably at 142 kV for 2 months.
Run-1 average was 124 kV.

Much reduced sparking
Still some cable issues above 145 kV




Run-1/2 e⁺ per fill




systematic runs

Run-2: 1.3 x Run-1

This despite a 19% reduction in intensity due to running the Li lens with 10% lower current



Run-1/2 : e⁺ per day




Run-2: 1.5 x Run-1 per day


When providing beam for more than 12 hrs in a day

Average running was 3.3% BNL/day

But when beam was available more than 12 hrs/day: 4.4% BNL/day




Impact of 5/14 running




5/14 running

Quad/Kicker Run-2 Uptime

Combined uptime of kicker/quads was 90% : far better than Run-1



DAQ Uptime



Reduced intensity per fill



	e+/fill	Effect	Factor
TDR	1100		
		Wedges	1.06
		Li Lens	0.81
		Kick (142 vs 155 kV)	0.84
		Quads (18 vs 28 kV)	0.92
		Actual beamline apertures	0.8
		TOTAL	0.53
RUN-2 Predicted			582
RUN-2 Actual			525 +/- 85*

*includes some systematic runs where rate was lower.

In physics running intensity is x0.5 that of TDR expectation.

Reduced uptime




	TDR	Run-2
g-2 systems	0.9	0.77 <i>0.92 (DAQ), 0.90 (kicker/quad), 0.93 (cryo)</i>
MI Cycles 1.4 vs 1.33 sec		0.95
Trolley Runs	0.94	0.94
Testbeam Users		0.91
Accelerator uptime*	0.85	0.82
TOTAL	0.72	0.5

TDR expectation: 1,100 e+/fill and 72% uptime → 825M/day

Predicted Run-2 : x 0.5 (e+/fill) and x 0.7 (uptime) → 290M/day

Run-2 actual : 286M/day (3.3% BNL/day).

A good week...




Anticipate Run-3 period to be more like this

Run-1 & Run-2 Integrated




Over x4 BNL before data-quality cuts (DQC)



Due to smoother running expect Run-2 DQC impact to be modest and so Run-2 analysis will be on ~ x1.8 BNL vs ~ x1.4 BNL in Run-1



Quick Analysis Highlights



Everything is blinded !




On average x2 better than BNL

Run-1 : Field Determination (ω_p)

Fixed probe \rightarrow Trolley \rightarrow Absolute (plunging probe)


Comparison of trolley and plunging probe measurements in two independent analyses

Quantity	Uncertainty (ppb)
Material Perturbation	10
Probe Pitch	5
Radiation Damping	1
Water Sample Oxygen Contamination	1
Water Sample Container Geometry	1
Water Diamagnetic Shielding	14
Water Magnetic Susceptibility	3
Water Temperature	16
TOTAL	24




Also two independent analysis doing interpolation from fixed probe measurements to the trolley measurements.

^3He cross calibration with H_2O probe



Sphere
 $1.24 \pm 2.29 \text{ Hz}$ ($20 \pm 37 \text{ ppb}$)



- Cross-calibration performed blind
- Analysis completed and unblinded
- Agreement with original BNL spherical probe and cylindrical plunging probe well within errors
- Difference also small relative to 540 ppb BNL error





Cylinder
 $0.92 \pm 2.23 \text{ Hz}$ ($15 \pm 36 \text{ ppb}$)

ω_a determination

$$N(t) = N_0 \cdot \Lambda(t) \cdot N_{1\text{CBO}}(t) \cdot N_{2\text{CBO}}(t) \cdot N_{VW}(t) \cdot N_{VO}(t) \cdot e^{-t/\tau} [1 + A_0 \cdot A_{1\text{CBO}}(t) \cdot \cos(\omega_a(R) \cdot t + \phi_0 + \phi_{1\text{CBO}}(t))]$$




Run-1 Analysis Status



Hardware blinding : x10 size of the BNL discrepancy wrt SM

Present datasets have stat. uncertainty 1/2 BNL



410 ppb (stat) Run-1 vs
460 ppb (BNL).

With Run-2 data approach
half the BNL stat.
uncertainty

Run-1 sub-datasets have statistical variations within expectations

Expectations for Run-3/4



- Accelerator uptime to be closer to 90% not 80% assuming overtime restrictions don't exist in Run-3/4.
- Modest g-2 DAQ improvements: x0.92 → x0.95.
- Optimisation of upstream wedges: x1.06 → x1.15
- Moves average BNL/day from 3.3% to 4.0%.
- Improved temperature control (critical for field & calorimeter gain)

Other potential improvements:

- install new inflector: x1.4
- increase quad / kicker voltage : x1.1
- faster switching PS (mitigates testbeam) : x1.05

In projections we are assuming 4% of BNL per day



When had POT for > 12 hrs a day in Run-2 **we had 4.2% BNL/day**

Improved temperature control for Run-3




Before

After




Large day/night oscillations
Significant overall rise over data taking period

Run-3/4



Possible Run-4 with μ^-

Run-4 could alternatively accumulate x8.5 the BNL μ^- sample
 e.g. if becomes systematics limited with μ^+ or μ^+ result $> 5\sigma$



Requires extra work in Run-3/4 shutdown

- Cryo pumps
- Kicker standoffs

New inflector would give x12 BNL μ^- sample.

Beam time is prioritised for Mu2e but initial part of Run-5 is likely g-2 running.

Ideas presently being pursued for future use of g-2 ring

- dedicated muon EDM experiment : new Si trackers in front of each calo
- $O(10 \text{ keV}) \nu_{\mu}$ mass measurement

HEPAP/P5 is summer of 2021 i.e. as Run-4 concludes.

Conclusions



- Run-1 and Run-2 were challenging: several technical issues had to be resolved
- Run-1 data will by itself surpass BNL precision
- Early analysis allowed us to quickly understand where we most needed to make improvements to control systematics for Run-2
- Run 2 (1.8×BNL) was taken under much more stable and improved conditions
- Expect even higher quality data in Runs 3 & 4 with 1×BNL collected per month
- (Blinded) independent analyses making good progress