




ATLAS flavour physics: recent results and prospects



Marcella Bona
(QMUL)
on behalf of the
ATLAS collaboration




8th Workshop on Theory, Phenomenology and Experiments
in Flavour Physics (FPCapri2022)
Anacapri, Italy
June 12th, 2022




B physics in ATLAS

- 25 fb⁻¹ in Run 1, and 139 fb⁻¹ in Run 2
- Has access to B, B_s, B_c, Λ_b, etc.
- Focus mostly on final states with muons
 - Typical trigger: di-muons with p_T thresholds at 4, 6 and 11 GeV
 - In 2018, a di-electron high-level trigger implemented and being analysed now




- **Properties of b-quark fragmentation to $B^\pm \rightarrow J/\psi K^\pm$ in Run 2**
 - arXiv:2108.11650, JHEP 12 (2021) 131
- **Study of $B_c^+ \rightarrow J/\psi D_s$ decays in Run 2**
 - arXiv:2203.01808, CERN-EP-2022-025
- **Rare and semi-rare decays:**
 - B to K* $\mu\mu$ angular analysis in Run 1 [JHEP 10 (2018) 047]
 - B_(s) to $\mu\mu$ in 2015-2016 Run 2 [JHEP 04 (2019) 098]
 - LHC combination B_(s) to $\mu\mu$ for 2020, [ATLAS-CONF-2020-049]
- **CP violating phase φ_s in $B_s^0 \rightarrow J/\psi\phi$ angular analysis:**
 - 2015-2017 Run 2 [Eur. Phys. J. C 81 (2021) 342]



Properties of b-quark fragmentation to $B^\pm \rightarrow J/\psi K^\pm$

Run2 result:

arXiv:2108.11650, JHEP 12 (2021) 131




Properties of b-quark fragmentation

- 139 fb⁻¹ of Run 2 data
- b-fragmentation functions provide:
 - Test of QCD at LHC energy; MC tunes
 - H → b \bar{b} and many other channels with b-jet signatures - dominant uncertainty
- We measure longitudinal (z) and transverse (p_T^{rel}) projections of the B[±] momentum to jet axis.

$$z = \frac{\vec{p}_J \cdot \vec{p}_B}{|\vec{p}_J|^2}; \quad p_T^{\text{rel}} = \frac{|\vec{p}_J \times \vec{p}_B|}{|\vec{p}_J|}$$



- B[±] mesons are associated to jets if they are within $\Delta R = 0.4$ from jet axis.
- B[±] invariant mass is used to extract differential cross section in each z or p_T^{rel} bins, for the lower and higher intervals of jet momenta:
50 GeV < p_T < 70 GeV
and p_T > 100 GeV.



Properties of b-quark fragmentation

arXiv:2108.11650
JHEP 12 (2021) 131

- Results for z distributions for $50 \text{ GeV} < p_T < 70 \text{ GeV}$ and $p_T > 100 \text{ GeV}$





- All Pythia fragmentation models give a decent description.
- Herwig7 (dipole) and, to some extent, Sherpa (cluster) visibly deviate from data.

Properties of b-quark fragmentation

arXiv:2108.11650
JHEP 12 (2021) 131

- Results for p_T^{rel} distributions for $50 \text{ GeV} < p_T < 70 \text{ GeV}$ and $p_T > 100 \text{ GeV}$



- All Pythia fragmentation models give a decent description.
- Both Herwig7 models and Sherpa (cluster) visibly deviate from data.



Study of $B_c^+ \rightarrow J/\psi D_s^{(*)}$ decays

Run2 result:

arXiv:2203.01808, CERN-EP-2022-025

Study of $B_c^+ \rightarrow J/\psi D_s^{(*)}$ decays

ArXiv:2203.01808
CERN-EP-2022-025


- Observed earlier by LHCb (PRD 87 (2013) 112012) and ATLAS (EPJC 76 (2016) 1) in Run 1.
- Using entire Run 2 dataset: aiming at more precise measurement of branching fractions and the final state polarisation
- Testing predictions of various theory models, e.g. pQCD calculation, relativistic potential models, sum rules calculations.

● D_s^+ and D_s^{*+} are reconstructed from their decays:

- $D_s^+ \rightarrow \varphi(K^+K^-)\pi^+$
- $D_s^{*+} \rightarrow D_s^+ \pi^0/\gamma$ (soft, not reco)
- Use $B_c^+ \rightarrow J/\psi\pi^+$ reference channel for BR measurement
- Fiducial range: $p_T(B_c^+) > 15$ GeV, $|\eta(B_c^+)| < 2.0$

Reference channel
with signal statistics



$$N(B_c^+ \rightarrow J/\psi\pi^+) = 8440^{+550}_{-470}$$



Study of $B_c^+ \rightarrow J/\psi D_s^{(*)}$ decays

ArXiv:2203.01808
CERN-EP-2022-025

- 2D fit to extract the signal parameters
- $m(J/\psi D_s^+)$ and the J/ψ helicity angle
- Both sensitive to polarisation of the final state particles J/ψ and D_s^+ in $B_c^+ \rightarrow J/\psi D_s^{*+}$ decay.



Left: fit to inv. mass $m(J/\psi D_s^+)$. Right: fit to $|\cos \theta'(\mu^+)|$, where $\theta'(\mu^+)$ is the helicity angle between μ^+ and D_s^+ momenta, in J/ψ rest frame.

Study of $B_c^+ \rightarrow J/\psi D_s^{(*)}$ decays

ArXiv:2203.01808
CERN-EP-2022-025


$$R_{D_s^+/\pi^+} \equiv \mathcal{B}_r(B_c^+ \rightarrow J/\psi D_s^+)/\mathcal{B}_r(B_c^+ \rightarrow J/\psi \pi^+) = 2.76 \pm 0.33(\text{stat.}) \pm 0.30(\text{syst.}) \pm 0.16(\text{BF})$$

$$R_{D_s^{*+}/\pi^+} \equiv \mathcal{B}_r(B_c^+ \rightarrow J/\psi D_s^{*+})/\mathcal{B}_r(B_c^+ \rightarrow J/\psi \pi^+) = 5.33 \pm 0.61(\text{stat.}) \pm 0.67(\text{syst.}) \pm 0.32(\text{BF})$$


$$R_{D_s^{*+}/D_s^+} \equiv \mathcal{B}_r(B_c^+ \rightarrow J/\psi D_s^+)/\mathcal{B}_r(B_c^+ \rightarrow J/\psi D_s^{*+}) = 1.93 \pm 0.24(\text{stat.}) \pm 0.10(\text{syst.})$$

$$B_c^+ \rightarrow J/\psi D_s^{*+} \text{ transvers polarisation fraction } \Gamma_{\pm\pm}/\Gamma = 0.70 \pm 0.10(\text{stat.}) \pm 0.04(\text{syst.})$$


- New results consistent with earlier measurements
- $R(D_s^{*+}/\pi^+)$ described well by the predictions



- $R(D_s^+/\pi^+)$ and $R(D_s^{*+}/D_s^+)$ predictions consistently deviate from data
- except QCD PM (PRD 61 (2000) 034012)
- $\Gamma_{\pm\pm}/\Gamma$ agrees with a naive spin-counting estimate of 2/3 and larger than predictions
- Hatched areas \rightarrow stat uncertainties; yellow bands \rightarrow total uncertainties.



NP searches and CP violation




CP violation in the SM and NP:

- $B_{(s)}$ systems are giving us a rather precise picture
- However there is some space for NP
- Could appear as new contributions in $\Delta F=2$ loop processes


$$A_q = C_{B_q} e^{2i\phi_{B_q}} A_q^{SM} e^{2i\phi_q^{SM}}$$

$$A_q = \left(1 + \frac{A_q^{NP}}{A_q^{SM}} e^{2i(\phi_q^{NP} - \phi_q^{SM})} \right) A_q^{SM} e^{2i\phi_q^{SM}}$$




The ratio of NP/SM amplitudes is:

- < 35% @68% prob. (45% @95%) in B_d mixing
- < 25% @68% prob. (35% @95%) in B_s mixing



[0707.0636 hep-ph]





Angular analysis on $B \rightarrow K^* \mu\mu$

Run1 result:

JHEP 10 (2018) 047, arXiv:1805.04000


HL-LHC prospects:

ATL-PHYS-PUB-2019-003

$K^* \mu\mu$ angular analysis

- Data collected in 2012 at 8 TeV with 20.3 fb^{-1} Run 1 data
- fold the angular distribution via trigonometric relations to reduce the number of free parameters
- Results are compatible with theoretical calculations & fits
- $P(P')$ parameters have reduced dependence on hadronic form factors.
- ATLAS gets deviations of about 2.5σ (2.7σ) from DHMV in $P'_4(P'_5)$ in $[4,6] \text{ GeV}^2$

q^2 [GeV^2]	P'_4	P'_5
[0.04, 2.0]	$0.31 \pm 0.40 \pm 0.20$	$0.67 \pm 0.26 \pm 0.16$
[2.0, 4.0]	$-0.76 \pm 0.31 \pm 0.21$	$-0.33 \pm 0.31 \pm 0.13$
[4.0, 6.0]	$0.64 \pm 0.33 \pm 0.18$	$0.26 \pm 0.35 \pm 0.18$
[0.04, 4.0]	$-0.30 \pm 0.24 \pm 0.17$	$0.32 \pm 0.21 \pm 0.11$
[1.1, 6.0]	$0.05 \pm 0.22 \pm 0.14$	$0.01 \pm 0.21 \pm 0.08$
[0.04, 6.0]	$0.05 \pm 0.20 \pm 0.14$	$0.27 \pm 0.19 \pm 0.06$




OPE and LHCb data fit: CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 QCD factorisation: DMVH: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

Projections for $K^* \mu\mu$ angular analysis at HL-LHC

- Extrapolation from signal/background yields in Run 1 and toy-MC simulations
- Accounting for improved performance of the ATLAS Upgraded tracking system
- Three trigger scenarios: high-yield, intermediate and low-statistics for signal.
- The precision on, for example, the P'_5 parameter expected to improve by factors of $\sim 9\times$, $\sim 8\times$, $\sim 5\times$ (for the three trigger scenarios) relative to Run 1



LHC phase	q^2 [GeV 2]	$\delta_{F_L}^{\text{tot}}$	$\delta_{P'_1}^{\text{tot}}$	$\delta_{P'_4}^{\text{tot}}$	$\delta_{P'_5}^{\text{tot}}$	$\delta_{P'_6}^{\text{tot}}$	$\delta_{P'_8}^{\text{tot}}$
Run 1	[0.04, 2.0]	0.11	0.31	0.45	0.31	0.21	0.51
	[2.0, 4.0]	0.12	0.61	0.37	0.34	0.34	0.57
	[4.0, 6.0]	0.18	0.50	0.38	0.39	0.30	0.43
HL-LHC $\mu 6\mu 6$	[0.04, 2.0]	0.010	0.027	0.037	0.037	0.019	0.046
	[2.0, 4.0]	0.008	0.093	0.040	0.038	0.040	0.070
	[4.0, 6.0]	0.016	0.083	0.032	0.047	0.033	0.041
HL-LHC $\mu 10\mu 6$	[0.04, 2.0]	0.011	0.037	0.046	0.040	0.023	0.055
	[2.0, 4.0]	0.011	0.103	0.047	0.042	0.044	0.075
	[4.0, 6.0]	0.018	0.100	0.040	0.053	0.038	0.052
HL-LHC $\mu 10\mu 10$	[0.04, 2.0]	0.018	0.065	0.076	0.059	0.041	0.093
	[2.0, 4.0]	0.017	0.15	0.074	0.068	0.059	0.100
	[4.0, 6.0]	0.026	0.17	0.074	0.082	0.063	0.090



ATL-PHYS-PUB-2019-003

Brewing in ATLAS...

- In 2018, a di-electron high-level trigger implemented and being analysed now
- Aiming at $R(K^*)$ measurement
- Angular analysis on di-muon final state on the whole 139 fb^{-1} Run-2 data also ongoing
- ATLAS potentially can do
 - $R(K)$, $R(\phi)$, $R(pK) = \text{BR}(\Lambda_b \rightarrow pK\mu\mu) / \text{BR}(\Lambda_b \rightarrow pKe e)$
- Stay tuned



rare B decays $B_{(s)} \rightarrow \mu^+ \mu^-$

Run1 result:

EPJ C76 (2016) 513, arXiv:1604.04263

Run2 result on 2015-2016 data:



JHEP 04 (2019) 098, arXiv:1812.03017

LHC combination:

ATLAS-CONF-2020-049

HL-LHC prospects:

ATL-PHYS-PUB-2018-005






JHEP 04 (2019) 098, arXiv:1812.03017

rare B decays $B_{(s)} \rightarrow \mu^+ \mu^-$

Run: 302137
Event: 1093131714
2016-06-16 01:48:30 CEST



Motivations

- Flavour Changing Neutral Currents (FCNC), CKM and helicity suppressed.
- SM prediction with small theoretical uncertainties of order 6-8%
- Perfect for indirect new physics searches: virtual new particles in the loop
 - both enhancement and suppression effects are possible

ATLAS analysis on 2015-2016 Run 2 data


- 36.2 fb⁻¹ dataset of 2015-2016 data taking:
 - effectively 26.3 fb⁻¹ for B → μμ
- Trigger: higher thresholds [4-6 GeV] than in Run 1,
 - L_{xy} > 0 request at trigger level

JHEP 04 (2019) 098
arXiv:1812.03017

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = \frac{N_{d(s)}}{\epsilon_{\mu^+ \mu^-}} \times \frac{\epsilon_{J/\psi K^+}}{N_{J/\psi K^+}} \times \frac{f_u}{f_{d(s)}} \times [\mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)]$$


Normalisation B yield extraction

- unbinned maximum likelihood fit of the invariant mass $m_{J/\psi K} \rightarrow m_{\mu\mu K}$



Backgrounds and control samples


- combinatorial background: μ 's from other b quarks
 - BDT classifier with 15 variables
- partially reconstructed B decays:
 - Same Vertex (SV): $B \rightarrow \mu\mu X$ decays
 - Same Side (SS): $b \rightarrow c\mu\nu \rightarrow s(d)\mu\mu\nu$
 - B_c decays: like $B_c \rightarrow J/\psi \mu\nu$
- semileptonic B and B_S decays: μ and charged hadron
- peaking background from hadronic $B_{(S)}$ decays:
 - B decays to two hadrons h (K/ π): $B_{(S)}^0 \rightarrow hh'$



Tight muon-ID against hadron misID

- negligible misidentification of protons ($< 0.01\%$)
- misidentification is 0.08% (0.10%) for K (π).

peaking-background events: 2.7 ± 1.3




Efficiency ratio $\epsilon_{\mu\mu} / \epsilon_{J/\psi K}$

- from MC and systematic from data-MC discrepancies
- For B_S^0 : 2.7% correction for lifetime difference of the B_S^0 mass eigenstates

Source	Contribution (%)
Statistical	0.8
BDT Input Variables	3.2
Kaon Tracking Efficiency	1.5
Muon trigger and reconstruction	1.0
Kinematic Reweighting (DDW)	0.8
Pile-up Reweighting	0.6

Signal yield extraction

- unbinned maximum likelihood fit to the dimuon mass simultaneously in 4 BDT bins
 - 18% signal efficiency each bin
 - signals, B to hh: 3 double Gaussians
 - continuum: first order polynomial
 - partially reconstructed B: exponential
 - semi-leptonic: exponential




Run 2 results and combinations with Run 1

- yields unconstrained:
 - $N_S = 80 \pm 22$ and $N_d = -12 \pm 20$
 - expected from the SM:
 - $N_S = 91 \pm$ and $N_d = 10$

Neyman Contours for Run 2:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.21^{+0.96+0.49}_{-0.91-0.30}) \times 10^{-9} = (3.2^{+1.1}_{-1.0}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.3 \times 10^{-10} \text{ @ 95\% CL}$$



Run 1 + Run 2 (2015+2016):
Compatible with SM at 2.4σ

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.8^{+0.8}_{-0.7}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 2.1 \times 10^{-10}$$

LHC combination from Summer 2020

- Combination from binned two-dimensional profile likelihoods
- Independent systematics, except for ratio of fragmentation fractions f_d/f_s ,
 - f_d/f_s profiled separately and its uncertainty included in one likelihood.

Latest LHCb result not included



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.69^{+0.37}_{-0.35}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 1.9 \times 10^{-10} \text{ at } 95\% \text{ CL.}$$

Prospect on $B_{(s)} \rightarrow \mu^+ \mu^-$ at ATLAS


- HL-LHC \rightarrow 3 trigger scenarios: with thresholds $(p_{T}^{\mu 1}, p_{T}^{\mu 2})$:
 - Conservative: (10 GeV, 10 GeV) \rightarrow $\times 15$ Run 1
 - Intermediate: (6 GeV, 10 GeV) \rightarrow $\times 60$ Run 1
 - High-yield: (6 GeV, 6 GeV) \rightarrow $\times 75$ Run 1

	$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$		$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-)$	
	stat [10^{-10}]	stat + syst [10^{-10}]	stat [10^{-10}]	stat + syst [10^{-10}]
Run 2	7.0	8.3	1.42	1.43
HL-LHC: Conservative	3.2	5.5	0.53	0.54
HL-LHC: Intermediate	1.9	4.7	0.30	0.31
HL-LHC: High-yield	1.8	4.6	0.27	0.28



ATL-PHYS-PUB-2018-005

Brewing in ATLAS...



- B_s lifetime analysis ongoing in the dimuon final state
- Branching ratio analysis on 2017+2018 to get to the whole 139 fb^{-1} Run-2 data ongoing
- Stay tuned

CP violation parameters from time-dependent angular analysis on $B_s \rightarrow J/\psi\phi$

Run1 result:


JHEP 08 (2016) 147, arXiv:1601.03297

Run2 result with 2015-2017 data:

Eur. Phys. J. C 81 (2021) 342, arXiv:2001.07115


HL-LHC prospects:

ATL-PHYS-PUB-2018-041



Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

- Parameters of the B_s system:
 - Decay width difference $\Delta\Gamma_s = \Gamma_L - \Gamma_H = 0.087 \pm 0.021 \text{ ps}^{-1}$ (SM) [*arXiv:1102.4274*]
 - CPV phase $\varphi_s \rightarrow$ weak phase between mixing and $b \rightarrow ccs$ decay
 - $\varphi_s = -2\beta_s = 0.0370 \pm 0.0010$ (SM) [*Utfit18*] with $\beta_s = \arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$
 - Golden mode: penguin diagrams contribute either with the same weak phase (λ^2) or they are CKM suppressed (λ^4)**
- Pseudoscalar to vector–vector decay
 - \rightarrow mixed CP-odd and CP-even ($L = 0, 1$ or 2).
 - Also K^+K^- pairs in S-wave \rightarrow CP-odd.
- Angular analysis: differential decay rate depends on amplitudes $A_0, A_{\parallel}, A_{\perp}, A_S$ (and interferences) and angles $\theta_T, \psi_T, \varphi_T$.





ATLAS Run-2 result

- 80.5 fb^{-1} of 13 TeV data (Run 2, 2015-2017)
- J/ψ trigger with muon p_T of 4 or 6 GeV
- Measurement of the proper decay time $t = L_{xy} m_B / p_T^B$
- Flavour tagging to identify the flavour of the b quark

ATLAS $B_s \rightarrow J/\psi\phi$ analysis: flavour tagging

- Flavour tagging to identify the flavour of the b quark:
 - opposite-side tagging (OST) using p_T -weighted charge of tracks in cone around muons / electrons / b jets
 - Calibrated on self-tagged $B^\pm \rightarrow J/\psi K^\pm$ events
 - Tag probabilities included in the B_s fit
 - Dilution $D(Q_x)$ and tagging power T_x defined as:

$$Q_x = \frac{\sum_i^{N \text{ tracks}} q_i \cdot (p_{Ti})^\kappa}{\sum_i^{N \text{ tracks}} (p_{Ti})^\kappa}$$








$$D(Q_x) = 2P(B|Q_x) - 1$$

$$T_x = \sum_i \epsilon_{xi} \cdot (2P(B|Q_{xi}) - 1)^2$$

Tag method	ϵ_x [%]	D_x [%]	T_x [%]
Tight muon	4.50 ± 0.01	43.8 ± 0.2	0.862 ± 0.009
Electron	1.57 ± 0.01	41.8 ± 0.2	0.274 ± 0.004
Low- p_T muon	3.12 ± 0.01	29.9 ± 0.2	0.278 ± 0.006
Jet	12.04 ± 0.02	16.6 ± 0.1	0.334 ± 0.006
Total	21.23 ± 0.03	28.7 ± 0.1	1.75 ± 0.01

ATLAS $B_s \rightarrow J/\psi\phi$ analysis: ML fit

- Unbinned maximum-likelihood fit
- B_s properties: mass m_B (and its error), proper decay time t , proper decay time error σ_t , tagging probability $P(B|Q_x)$
- Transversity angles: $\Omega(\theta_T, \psi_T, \phi_T)$
- Physical parameters: $\Delta\Gamma_s, \varphi_s, \Gamma_s, |A_0(0)|^2, |A_{\parallel}(0)|^2, \delta_{\parallel}, \delta_{\perp}, |A_s(0)|^2$ and δ_s
- Systematics:
 - Lifetime model: varying p_T bins and signal fraction
 - Backgrounds: $B_d / \Lambda_b /$ angular models varied / p_T bins varied
 - Tagging: variation of the parameterisation / recalibration from MC samples / pile-up effects



ATLAS $B_s \rightarrow J/\psi\phi$ analysis: Run-2 results



ATLAS Run-2 result on 80.5 fb^{-1} of 2015-2017 data

Run 2 only (80.5 fb^{-1}):

Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.081	0.041	0.022
$\Delta\Gamma_s$ [ps^{-1}]	0.0607	0.0047	0.0043
Γ_s [ps^{-1}]	0.6687	0.0015	0.0022
$ A_{\parallel}(0) ^2$	0.2213	0.0019	0.0023
$ A_0(0) ^2$	0.5131	0.0013	0.0038
$ A_S(0) ^2$	0.0321	0.0033	0.0046
$\delta_{\perp} - \delta_S$ [rad]	-0.25	0.05	0.04
Solution (a)			
δ_{\perp} [rad]	3.12	0.11	0.06
δ_{\parallel} [rad]	3.35	0.05	0.09
Solution (b)			
δ_{\perp} [rad]	2.91	0.11	0.06
δ_{\parallel} [rad]	2.94	0.05	0.09

Two solutions in $\delta_{\parallel} - \delta_{\perp}$ plane,
negligible impact on other parameters

Run 1 (19.2 fb^{-1}) & Run 2 (80.5 fb^{-1}):




ATLAS $B_s \rightarrow J/\psi\phi$ analysis: Run1+2 combination

ATLAS Run 1 & Run 2 combined
(19.2 fb⁻¹ + 80.5 fb⁻¹)

Parameter	Value	Solution (a)	
		Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.087	0.036	0.021
$\Delta\Gamma_s$ [ps ⁻¹]	0.0657	0.0043	0.0037
Γ_s [ps ⁻¹]	0.6703	0.0014	0.0018
$ A_{\parallel}(0) ^2$	0.2220	0.0017	0.0021
$ A_0(0) ^2$	0.5152	0.0012	0.0034
$ A_S ^2$	0.0343	0.0031	0.0045
δ_{\perp} [rad]	3.22	0.10	0.05
δ_{\parallel} [rad]	3.36	0.05	0.09
$\delta_{\perp} - \delta_S$ [rad]	-0.24	0.05	0.04

Comparison with CMS & LHCb:



$$\phi_s = -0.087 \pm 0.036 \text{ (stat)} \pm 0.021 \text{ (syst) rad}$$



$$\Delta\Gamma_s = 0.0657 \pm 0.0043 \text{ (stat)} \pm 0.0037 \text{ (syst) ps}^{-1}$$

- ϕ_s result consistent with results from CMS, LHCb and SM
- Competitive single measurement of $\Delta\Gamma_s$, Γ_s and helicity parameters
- Still to add 60 fb⁻¹ of 2018 data

$B_s \rightarrow J/\psi\phi$ results: HFLAV average


HFLAV average for PDG21:
 $\phi_s = -0.050 \pm 0.019$ rad

Because of tensions, errors on Γ_s and $\Delta\Gamma_s$ scaled by 2.5 and 1.77



ATLAS $B_s \rightarrow J/\psi\phi$ results: HL-LHC projections

- Updated tracking (ITk): proper decay time resolution improved by 21% w.r.t. Run 2
- Three trigger scenarios for thresholds
- Improvements w.r.t. Run 1:
 - ϕ_s stat: better by ~9x to 20x
 - uncertainty on ϕ_s at least as the theory error
 - $\Delta\Gamma_s$ stat: better by ~4x to 10x



ATL-PHYS-PUB-2018-041

Brewing in ATLAS...

- Time-dependent analysis ongoing on the 2018 to get to the whole 139 fb^{-1} Run-2 data ongoing
- Includes λ and Δm_s among the fit parameters
- LHC combination group in stand-by for the updated analyses
- Stay tuned

Conclusions

- ATLAS is competitive in B physics
 - Thanks to accumulated statistical samples
 - Thanks to some detector performance (tracking)
 - Perfect example the angular analysis of the golden mode $B_s \rightarrow J/\psi\phi$
 - **Working on the updates of all analyses to full Run-2 statistics and preparing for Run 3**



back-up slides


Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

- Systematics:
 - Lifetime model*: varying p_T bins and signal fraction
 - Backgrounds*: $B_d / \Lambda_b /$ angular models varied / p_T bins varied
 - Tagging*: variation of the parameterisation / recalibration from MC samples / pile-up effects

	ϕ_s [10^{-3} rad]	$\Delta\Gamma_s$ [10^{-3} ps $^{-1}$]	Γ_s [10^{-3} ps $^{-1}$]	$ A_{\parallel}(0) ^2$ [10^{-3}]	$ A_0(0) ^2$ [10^{-3}]	$ A_S(0) ^2$ [10^{-3}]	δ_{\perp} [10^{-3} rad]	δ_{\parallel} [10^{-3} rad]	$\delta_{\perp} - \delta_S$ [10^{-3} rad]
Tagging	19	0.4	0.3	0.2	0.2	1.1	17	19	2.3
ID alignment	0.8	0.2	0.5	< 0.1	< 0.1	< 0.1	11	7.2	< 0.1
Acceptance	0.5	0.3	< 0.1	1.0	0.9	2.9	37	64	8.6
Time efficiency	0.2	0.2	0.5	< 0.1	< 0.1	0.1	3.0	5.7	0.5
Best candidate selection	0.4	1.6	1.3	0.1	1.0	0.5	2.3	7.0	7.4
Background angles model:									
Choice of fit function	2.5	< 0.1	0.3	1.1	< 0.1	0.6	12	0.9	1.1
Choice of p_T bins	1.3	0.5	< 0.1	0.4	0.5	1.2	1.5	7.2	1.0
Choice of mass window	9.3	3.3	0.2	0.4	0.8	0.9	17	8.6	6.0
Choice of sidebands intervals	0.4	0.1	0.1	0.3	0.3	1.3	4.4	7.4	2.3
Dedicated backgrounds:									
B_d^0	2.6	1.1	< 0.1	0.2	3.1	1.5	10	23	2.1
Λ_b	1.6	0.3	0.2	0.5	1.2	1.8	14	30	0.8
Alternate Δm_s	1.0	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	15	4.0	< 0.1
Fit model:									
Time res. sig frac	1.4	1.1	0.5	0.5	0.6	0.8	12	30	0.4
Time res. p_T bins	0.7	0.5	0.8	0.1	0.1	0.1	2.2	14	0.7
S-wave phase	0.3	< 0.1	< 0.1	< 0.1	< 0.1	0.2	8.0	15	37
Fit bias	5.7	1.3	1.2	1.3	0.4	1.1	3.3	19	0.3
Total	22	4.3	2.2	2.3	3.8	4.6	55	88	39

Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

$$\frac{d^4\Gamma}{dt d\Omega} = \sum_{k=1}^{10} \mathcal{O}^{(k)}(t) g^{(k)}(\theta_T, \psi_T, \phi_T),$$



k	$\mathcal{O}^{(k)}(t)$	$g^{(k)}(\theta_T, \psi_T, \phi_T)$
1	$\frac{1}{2} A_0(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$2 \cos^2 \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$
2	$\frac{1}{2} A_{ }(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T (1 - \sin^2 \theta_T \sin^2 \phi_T)$
3	$\frac{1}{2} A_{\perp}(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T \sin^2 \theta_T$
4	$\frac{1}{2} A_0(0) A_{ }(0) \cos \delta_{ }$ $\left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin^2 \theta_T \sin 2\phi_T$
5	$ A_{ }(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos(\delta_{\perp} - \delta_{ }) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin(\delta_{\perp} - \delta_{ }) \cos(\Delta m_s t) - \cos(\delta_{\perp} - \delta_{ }) \cos \phi_s \sin(\Delta m_s t)) \right]$	$-\sin^2 \psi_T \sin 2\theta_T \sin \phi_T$
6	$ A_0(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos \delta_{\perp} \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin \delta_{\perp} \cos(\Delta m_s t) - \cos \delta_{\perp} \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin 2\theta_T \cos \phi_T$
7	$\frac{1}{2} A_S(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{2}{3} (1 - \sin^2 \theta_T \cos^2 \phi_T)$
8	$\alpha A_S(0) A_{ }(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \sin(\delta_{ } - \delta_S) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos(\delta_{ } - \delta_S) \cos(\Delta m_s t) - \sin(\delta_{ } - \delta_S) \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin^2 \theta_T \sin 2\phi_T$
9	$\frac{1}{2}\alpha A_S(0) A_{\perp}(0) \sin(\delta_{\perp} - \delta_S)$ $\left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin 2\theta_T \cos \phi_T$
10	$\alpha A_0(0) A_S(0) \left[\frac{1}{2}(e^{-\Gamma_H^{(s)} t} - e^{-\Gamma_L^{(s)} t}) \sin \delta_S \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos \delta_S \cos(\Delta m_s t) + \sin \delta_S \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{4}{3} \sqrt{3} \cos \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$





TD angular analysis of $B_s \rightarrow J/\psi\phi$

ATLAS Run-1 result:



- 14.3 fb⁻¹ of ATLAS data from 2012 at 8 TeV
 - Results:
 - $\phi_s = -0.090 \pm 0.078$ (stat) ± 0.041 (syst) rad
 - $\Delta\Gamma_s = 0.085 \pm 0.011$ (stat) ± 0.007 (syst) ps⁻¹
- [JHEP 08 (2016) 147]

- Agrees with SM
- Consistent with other experiments
- Consistent with previous analysis, using 2011 data at 7 TeV
 - [Phys. Rev. D 90, 052007 (2014)]
- A Best Linear Unbiased Estimate (BLUE) combination used to combine 7 and 8 TeV measurements




Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

ATLAS combined Run-1 result:



Motivations and predictions


- Decays of B^0 and B_s^0 into two leptons have to proceed through Flavour Changing Neutral Currents (FCNC)
 - forbidden at tree level in the SM
- In addition, they are CKM and helicity suppressed.
- Within the SM, they can be calculated with small theoretical uncertainties of order 6-8%



meson type	Lepton type		
	e	μ	τ
B^0	$(2.48 \pm 0.21)10^{-15}$	$(1.06 \pm 0.09)10^{-10}$	$(2.22 \pm 0.19)10^{-8}$
B_s^0	$(8.54 \pm 0.55)10^{-14}$	$(3.65 \pm 0.23)10^{-9}$	$(7.73 \pm 0.49)10^{-7}$

Bobeth et al.,
 PRL 112 (2104)
 101801
 [includes NLO EM
 and NNLO QCD
 corrections]

- Perfect ground for indirect new physics searches:
 - virtual new particles can contribute to the loop
 - both enhancement and suppression effects are possible



ATLAS analysis on 2015-2016 Run 2 data

JHEP 04 (2019) 098, arXiv:1812.03017

- 36.2/fb dataset of 2015-2016 data taking:
 - effectively 26.3/fb for $B \rightarrow \mu\mu$
 - 15.1/fb for $B \rightarrow J/\psi\Phi$ and $B \rightarrow J/\psi K$
- Trigger: higher thresholds [4-6 GeV] than in Run1,
 - $L_{xy} > 0$ request at trigger level

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = \frac{N_{d(s)}}{\epsilon_{\mu^+ \mu^-}} \times \frac{\epsilon_{J/\psi K^+}}{N_{J/\psi K^+}} \times \frac{f_u}{f_{d(s)}} \times [\mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)]$$



- correction for the different hadronisation probabilities for $B_{(s)}^0$ and B^0 vs B^\pm
- include the B^\pm and J/ψ branching fractions
- correction for the efficiencies of the two channels
- normalisation yield and efficiency ratio define the factor:

$$\mathcal{D}_{\text{norm}} = N_{J/\psi K^+} \left(\frac{\epsilon_{\mu^+ \mu^-}}{\epsilon_{J/\psi K^+}} \right)$$

Background contributions

In order of relative magnitude:


- combinatorial background:
 - two real muons from different b quarks
- partially reconstructed B decays:
 - two real muons
 - Same Vertex (SV): $B \rightarrow mmX$ decays
 - Same Side (SS): semileptonic decay cascades ($b \rightarrow cmn \rightarrow s(d)mmnn$)
 - B_c decays: like $B_c \rightarrow J/\psi mn$
 - all these accumulate at low values of the dimuon invariant mass
- semileptonic B and B_s decays:
 - one real muon and a charged hadron.
- peaking background from charmless hadronic $B_{(s)}$ decays:
 - B decays into two hadrons h (kaons and pions): $B^0_{(s)} \rightarrow hh'$
 - smaller component, but overlays with the signal in dimuon invariant mass



Tight muon-ID against hadron misidentification


- mis-identification reduced by 0.39^2 using standard 'tight' ATLAS selections
- studied on simulated samples
- validated on control regions
- negligible misidentification of protons ($< 0.01\%$)
- misidentification is 0.08% (0.10%) for K(p).

peaking-background events: 2.7 ± 1.3



BDT against combinatorial bkg

- MVA classifier to discriminate from signal
- trained and tested on mass sidebands
 - divided in 3 subsets
 - 3 independent BDTs
 - compatible performance
- 15 variables related to properties of B candidates, muons from the B decay, other tracks from the same collision and to pile-up vertices.



Normalisation B yield extraction



- unbinned maximum likelihood fit of the invariant mass $m_{J/\psi K} \rightarrow m_{mmK}$
- cross-checked with raw relative yield of J/ψp over J/ψK ratio
 $r_{p/K} = (3.71 \pm 0.09)\%$

$$D_{\text{norm}} = N_{J/\psi K^+} \left(\frac{\epsilon_{\mu^+ \mu^-}}{\epsilon_{J/\psi K^+}} \right)$$

Efficiency ratio $\epsilon_{\mu\mu}/\epsilon_{J/\psi K}$





- efficiency ratio from MC
- systematic from data-MC discrepancies
- For B^0_s : 2.7% correction for lifetime difference of the B^0_s mass eigenstates

Source	Contribution (%)
Statistical	0.8
BDT Input Variables	3.2
Kaon Tracking Efficiency	1.5
Muon trigger and reconstruction	1.0
Kinematic Reweighting (DDW)	0.8
Pile-up Reweighting	0.6




Signal yield extraction

- signal yields extracted with a unbinned maximum likelihood fit to the dimuon mass
- fit performed simultaneously in four BDT bins
- 18% signal efficiency








- signals, B to hh: 3 double Gaussians
- continuum: first order polynomial
- partially reconstructed B: exponential
- semi-leptonic: exponential



Signal yield extraction

- yields unconstrained:
 - $N_S = 80 \pm 22$ and $N_d = -12 \pm 20$
- expected from the SM:
 - $N_S = 91 \pm$ and $N_d = 10$



- consistent with Standard Model predictions
- likelihood maximum:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = \left(3.21_{-0.83-0.31}^{+0.90+0.48} \right) \times 10^{-9}$$



$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = \left(-1.3_{-1.9-0.8}^{+2.2+0.7} \right) \times 10^{-10}$$

Combination of Run 1 and Run 2 results

Neyman Contours yield for Run 2:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.21^{+0.96+0.49}_{-0.91-0.30}) \times 10^{-9} = (3.2^{+1.1}_{-1.0}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.3 \times 10^{-10} \text{ @ 95\% CL}$$



Run 1 + Run 2 (2015+2016) combination:
Compatible with SM at 2.4σ



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.8^{+0.8}_{-0.7}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 2.1 \times 10^{-10}$$

Angular analysis on $B \rightarrow K^* \mu \mu$

JHEP 10 (2018) 047, arXiv:1805.04000

- FCNC b to s transition with a BR $\sim 1.1 \cdot 10^{-6}$
- Angular distribution of the 4 particles in the final state sensitive to new physics for the interference of NP and SM diagrams



- Decay described by three angles (q_L, q_K, ϕ) and the di-muon mass squared $q^2 \rightarrow$ angular distribution in bins of q^2 as function of q_L, q_K and ϕ .

$$\frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{d \cos \theta_\ell d \cos \theta_K d\phi dq^2} = \frac{9}{32\pi} \left[\frac{3(1-F_L)}{4} \sin^2 \theta_K + F_L \cos^2 \theta_K + \frac{1-F_L}{4} \sin^2 \theta_K \cos 2\theta_\ell - F_L \cos^2 \theta_K \cos 2\theta_\ell + S_3 \sin^2 \theta_K \sin^2 \theta_\ell \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + S_6 \sin^2 \theta_K \cos \theta_\ell + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_\ell \sin 2\phi \right].$$



- The S parameters are translated into the $P^{(\prime)}$ parameters via

$$P_1 = \frac{2S_3}{1-F_L} \quad P'_{i=4,5,6,8} = \frac{S_{j=4,5,7,8}}{\sqrt{F_L(1-F_L)}}$$

with reduced dependence on the hadronic form factors.

Angular analysis on $B \rightarrow K^* \mu \mu$

- another way to look at FCNC: $b \rightarrow s$ transition with a BR $\sim 1.1 \cdot 10^{-6}$
- angular distribution of the 4 particles in the final state sensitive to new physics for the interference of NP and SM diagrams
- allows measuring a large set of angular parameters sensitive to Wilson coefficients $C^{(i)}_7, C^{(i)}_9, C^{(i)}_{10}, C^{(i)}_{S,P}$



- decay described by three angles (θ_L, θ_K, ϕ) and the di-muon mass squared $q^2 \rightarrow$ the angular distribution is analysed in finite bins of q^2 as a function of θ_L, θ_K and ϕ .
- LHCb reports a 3.4σ deviation from the SM.

JHEP 02 (2016) 104
arXiv:1512.04442

Angular analysis on $B \rightarrow K^* \mu \mu$

- B^0 flavour eigenstate can be identified through the $K^* \rightarrow K^- \pi^+$ decay
- angular distribution given by:

$$\frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{d\cos\theta_\ell d\cos\theta_K d\phi dq^2} = \frac{9}{32\pi} \left[\frac{3(1-F_L)}{4} \sin^2\theta_K + F_L \cos^2\theta_K + \frac{1-F_L}{4} \sin^2\theta_K \cos 2\theta_\ell \right. \\ \left. - F_L \cos^2\theta_K \cos 2\theta_\ell + S_3 \sin^2\theta_K \sin^2\theta_\ell \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi \right. \\ \left. + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + S_6 \sin^2\theta_K \cos \theta_\ell \right. \\ \left. + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi \right. \\ \left. + S_9 \sin^2\theta_K \sin^2\theta_\ell \sin 2\phi \right].$$

- the S parameters are translated into the $P^{(\prime)}$ parameters via

$$P_1 = \frac{2S_3}{1-F_L} \quad P'_{i=4,5,6,8} = \frac{S_{j=4,5,7,8}}{\sqrt{F_L(1-F_L)}}$$





- the $P^{(\prime)}$ parameters are expected to have a reduced dependence on the hadronic form factors.
- ATLAS and CMS need to fold the angular distribution via trigonometric relations to reduce the number of free parameters

Analysis strategy for $B \rightarrow K^* \mu\mu$

- Data collected in 2012 at 8 TeV with 20.3 fb⁻¹ Run 1 data
- Measured in 6 (overlapping) bins of q^2 in the range [0.04, 6] GeV²
- 4 sets of fits for three parameters (F_L , S_3 and S_j with $j=4,5,7,8$)
- Selection of triggers with muon p_T thresholds starting at 4 GeV
- K^* tagged by the kaon sign:
 - dilution from mistag probability included in $(1-2\langle w \rangle)$:
 - $\langle w \rangle \sim 10.9(1)\%$ with small dependence on q^2
- 787 events selected with $q^2 < 6$ GeV²
- Extended unbinned maximum likelihood fits in each of the fit variants in each q^2 bin:
 - two step fit procedure: first fit the invariant mass distribution
 - then add to the fit the angular distributions to extract the F_L and $S(P)$ parameters
- Signal shape studies from control samples $K^* J/\psi$ and $K^* \psi(2S)$

Fit projections




- fit $m(K^*_{\mu\mu})$, $\cos\theta_L$, $\cos\theta_K$ and ϕ to isolate signal and extract parameters of interest.




- Data shown for $[0.04, 2.0] \text{ GeV}^2$
- projections for the S_5 fit.
- Approx 106-128 signal events in 2 GeV^2 q^2 bin.
- Similar results for the other q^2 bins and other fit variants.

Angular analysis results

- Results are compatible with theoretical calculations & fits:




q^2 [GeV ²]	P_1	P'_4	P'_5
[0.04, 2.0]	$-0.05 \pm 0.30 \pm 0.08$	$0.31 \pm 0.40 \pm 0.20$	$0.67 \pm 0.26 \pm 0.16$
[2.0, 4.0]	$-0.78 \pm 0.51 \pm 0.34$	$-0.76 \pm 0.31 \pm 0.21$	$-0.33 \pm 0.31 \pm 0.13$
[4.0, 6.0]	$0.14 \pm 0.43 \pm 0.26$	$0.64 \pm 0.33 \pm 0.18$	$0.26 \pm 0.35 \pm 0.18$
[0.04, 4.0]	$-0.22 \pm 0.26 \pm 0.16$	$-0.30 \pm 0.24 \pm 0.17$	$0.32 \pm 0.21 \pm 0.11$
[1.1, 6.0]	$-0.17 \pm 0.31 \pm 0.13$	$0.05 \pm 0.22 \pm 0.14$	$0.01 \pm 0.21 \pm 0.08$
[0.04, 6.0]	$-0.15 \pm 0.23 \pm 0.10$	$0.05 \pm 0.20 \pm 0.14$	$0.27 \pm 0.19 \pm 0.06$



OPE and LHCb data fit: CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 QCD factorisation: DMVH: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

Angular analysis results

- ATLAS gets deviations of about 2.5σ (2.7σ) from DHMV in P'_4 (P'_5) in $[4,6] \text{ GeV}^2$



CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 DHMV: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.



TD angular analysis of $B_s \rightarrow J/\psi\phi$

ATL-PHYS-PUB-2018-041

Period	L_{int} [fb^{-1}]	N_{sig}	f_{sig}	Tag Power [%]	$\sigma(\tau)$ [ps]	$\delta_{\phi_s}^{\text{stat}}$ [rad] measured (extrapolated)	$\delta_{\Delta\Gamma_s}^{\text{stat}}$ [ps^{-1}] measured (extrapolated)
2012	14.3	73693	0.20	1.49	0.091	0.082	0.013
2011	4.9	22690	0.17	1.45	0.100	0.25 (0.22)	0.021 (0.023)
HL-LHC	3000					$\delta_{\phi_s}^{\text{stat}}$ [rad] extrapolated	
Trigger $\mu 6\mu 6$		$9.72 \cdot 10^6$	0.17	1.49	0.048	0.004	0.0011
Trigger $\mu 10\mu 6$		$5.93 \cdot 10^6$	0.17	1.49	0.044	0.005	0.0014
Trigger $\mu 10\mu 10$		$1.75 \cdot 10^6$	0.15	1.49	0.038	0.009	0.003

Around and outside B physics...


Test of the universality of τ and μ in W decays from $t\bar{t}$ events

Muons from W and muons from tau distinguished using the lifetime of the τ , through the muon transverse impact parameter, and differences in the muon transverse momentum spectra.

Tag and probe approach: tag leptons to select the events, probe muon from prompt decay, $W \rightarrow \mu\nu_\mu$, or via intermediate τ , $W \rightarrow \tau\nu_\tau \rightarrow \mu\nu_\mu\nu_\tau\nu_\tau$.

Di-leptonic $t\bar{t}$ events with either one electron and one muon ($e\text{-}\mu$ channel), or two muons ($\mu\text{-}\mu$ channel).

[arXiv:2007.14040](https://arxiv.org/abs/2007.14040) [hep-ex]



$$R(\tau/\mu) = 0.992 \pm 0.013 [\pm 0.007 \text{ (stat)} \pm 0.011 \text{ (syst)}]$$

CP violation in the SM and NP:

- $B_{(s)}$ systems are giving us a rather precise picture
- However there is some space for NP
- Could appear as new contributions in $\Delta\Gamma^{\circ}$ loop processes

$$A_q = C_{B_q} e^{2i\phi_{B_q}} A_q^{SM} e^{2i\phi_{B_q}^{\circ}}$$

$$A_q = \left(1 + \frac{A_q^{NP}}{A_q^{SM}} e^{2i(\phi_q^{NP} - \phi_q^{SM})} \right) A_q^{SM} e^{2i\phi_q^{SM}}$$

The ratio of NP/SM amplitudes need to be:

