



Contribution ID: 54

Type: **not specified**

Dark Photon Limits: a Cookbook

Wednesday, 16 June 2021 18:20 (20 minutes)

The dark photon is a massive hypothetical particle that interacts with the Standard Model by kinetically mixing with the visible photon. Due to the similarity with the electromagnetic signals generated by axions, many putative bounds on dark photon signals are simply reinterpretations of historical bounds set by axion haloscopes. However, the dark photon has a property that the axion does not: an intrinsic polarization. Due to the rotation of the Earth, accurately incorporating this polarization into dark photon analyses is nontrivial, and highly experiment-dependent. Several well-known searches for axions employ techniques for testing signals that preclude their ability to set exclusion limits on dark photons, and hence should not be reinterpreted as such. Most experiments do not have a straight forward reinterpretation for polarized dark photons. On the other hand, we find that if one does account for the dark photon's polarization, and the rotation of the Earth, the theoretical sensitivity to the dark photon's kinetic mixing parameter can be improved by over an order of magnitude. Here, we detail the strategies that would need to be taken to properly optimise a dark photon search. These include a judiciously choosing the orientation of the experiment, as well as strategically timing any repeated measurements and splitting measurements into multiple parts. Such strategies have significant impact on limits without additional time or cost.

Speaker

Alexander Millar

Primary authors: MILLAR, Alexander (Stockholm University); Dr CAPUTO, Andrea (Tel-Aviv University); Dr O'HARE, Ciaran (The University of Sydney); Dr VITAGLIANO, Edoardo (UCLA)

Presenter: MILLAR, Alexander (Stockholm University)

Session Classification: Session 12