

# Probing gluon TMDs

—

## overview



Daniël Boer  
IWHSS 2020  
November 17, 2020



/ university of  
groningen

# Outline


- Parallels between quark and gluon TMD investigations
- Process dependence of gluon TMDs
  - sign change relation for gluon Sivers TMDs
  - small-x limit & Wilson loop matrix elements
- Probes of unpolarized gluon TMDs
- Probes of linearly polarized gluon TMDs

# Parallels between quark and gluon TMD investigations

# Typical TMD processes


Semi-inclusive DIS is a process sensitive to the transverse momentum of quarks

$$e p \rightarrow e' h X$$



D-meson pair production is sensitive to transverse momentum of gluons

$$e p \rightarrow e' D \bar{D} X$$



# Gluons TMDs

# The transverse momentum dependent gluon correlator:

$$\Gamma_g^{\mu\nu}(x, p_T) \propto \langle P | F^{+\nu}(0) \mathcal{U} F^{+\mu}(\xi^-, \xi_T) \mathcal{U}' | P \rangle$$

## For unpolarized protons:

$$\Gamma_U^{\mu\nu}(x, \mathbf{p}_T) = \frac{x}{2} \left\{ -g_T^{\mu\nu} f_1^g(x, \mathbf{p}_T^2) + \left( \frac{p_T^\mu p_T^\nu}{M_p^2} + g_T^{\mu\nu} \frac{\mathbf{p}_T^2}{2M_p^2} \right) h_1^{\perp g}(x, \mathbf{p}_T^2) \right\}$$

unpolarized gluon TMD      linearly polarized gluon TMD

Gluons inside *unpolarized* protons can be polarized!

[Mulders, Rodrigues, 2001]

## For transversely polarized protons:


$$\Gamma_T^{\mu\nu}(x, \mathbf{p}_T) = \frac{x}{2} \left\{ g_T^{\mu\nu} \frac{\epsilon_T^{\rho\sigma} p_{T\rho} S_{T\sigma}}{M_p} f_{1T}^{\perp g}(x, \mathbf{p}_T^2) + \dots \right\}$$

Perhaps surprisingly, these TMDs have not been extracted from experiments yet

# Entering the gluon TMD era


## Higgs p<sub>T</sub> distribution

$$pp \rightarrow H(\rightarrow \gamma\gamma) + X$$



[Gutierrez-Reyes, Leal-Gomez, Scimemi, Vladimirov, 2019]

## J/ψ pair production



[Scarpa et al., 2020]

## Sivers asymmetry in high-p<sub>T</sub> hadron pair production

$$A^{Siv} = -0.23 \pm 0.08 \text{ (stat)} \pm 0.05 \text{ (syst)} \text{ at } \langle x_g \rangle = 0.15$$

[COMPASS Collab., 2017]

## GPM studies of $A_N^{\pi, D}$

[D'Alesio, Murgia, Pisano, 2015;  
& Taels, 2017]

# Parallels between quarks and gluons

$$\Phi_U(x, \mathbf{k}) = \frac{1}{2} \left[ \not{n} f_1(x, \mathbf{k}^2) + \frac{\sigma_{\mu\nu} k_T^\mu \bar{n}^\nu}{M} h_1^\perp(x, \mathbf{k}^2) \right],$$

$$\Phi_L(x, \mathbf{k}) = \frac{1}{2} \left[ \gamma^5 \not{n} S_L g_1(x, \mathbf{k}^2) + \frac{i\sigma_{\mu\nu} \gamma^5 \bar{n}^\mu k_T^\nu S_L}{M} h_{1L}^\perp(x, \mathbf{k}^2) \right],$$

$$\begin{aligned} \Phi_T(x, \mathbf{k}) = & \frac{1}{2} \left[ \frac{\not{n} \epsilon_T^{S_T k_T}}{M} f_{1T}^\perp(x, \mathbf{k}^2) + \frac{\gamma^5 \not{n} \mathbf{k} \cdot \mathbf{S}_T}{M} g_{1T}(x, \mathbf{k}^2) \right. \\ & \left. + i\sigma_{\mu\nu} \gamma^5 \bar{n}^\mu S_T^\nu h_1(x, \mathbf{k}^2) - \frac{i\sigma_{\mu\nu} \gamma^5 \bar{n}^\mu k_T^\nu S_{T\rho}}{M^2} h_{1T}^\perp(x, \mathbf{k}^2) \right] \end{aligned}$$

$$\Gamma_U^{ij}(x, \mathbf{k}) = x \left[ \delta_T^{ij} f_1(x, \mathbf{k}^2) + \frac{k_T^{ij}}{M^2} h_1^\perp(x, \mathbf{k}^2) \right],$$


$$\Gamma_L^{ij}(x, \mathbf{k}) = x \left[ i\epsilon_T^{ij} S_L g_1(x, \mathbf{k}^2) + \frac{\epsilon_T^{\{i} \alpha k_T^{j\}\alpha} S_L}{2M^2} h_{1L}^\perp(x, \mathbf{k}^2) \right],$$

$$\begin{aligned} \Gamma_T^{ij}(x, \mathbf{k}) = & x \left[ \frac{\delta_T^{ij} \epsilon_T^{S_T k_T}}{M} f_{1T}^\perp(x, \mathbf{k}^2) + \frac{i\epsilon_T^{ij} \mathbf{k} \cdot \mathbf{S}_T}{M} g_{1T}(x, \mathbf{k}^2) \right. \\ & \left. - \frac{\epsilon_T^{k_T \{i} S_T^{j\}} + \epsilon_T^{S_T \{i} k_T^{j\}}}{4M} h_1(x, \mathbf{k}^2) - \frac{\epsilon_T^{\{i} \alpha k_T^{j\}\alpha S_T}}{2M^3} h_{1T}^\perp(x, \mathbf{k}^2) \right] \end{aligned}$$

For quarks the BM & Sivers TMDs are T-odd and the h-type functions are chiral-odd

For gluons  $h_{1\perp}$  is T-even and  $h_1$  is  $k_T$ -odd, T-odd and unrelated to transversity

# Parallels between SIDIS and HQ pair production



The heavy quarks will not be exactly back-to-back in the transverse plane:

$$K_{\perp} = (K_{Q\perp} - K_{\bar{Q}\perp})/2$$

$$q_T = K_{Q\perp} + K_{\bar{Q}\perp}$$

$$|q_T| \ll |K_{\perp}|$$

$\phi_T, \phi_{\perp}$  are the angles of  $q_T, K_{\perp}$

Linear gluon polarization shows up as a  $\cos 2\phi_T$  or  $\cos 2(\phi_T - \phi_{\perp})$  distribution

Despite the differences in properties of some of the quark and gluon TMDs, the asymmetries they lead to are analogous for SIDIS and HQ pair production

There is a “Collins” asymmetry without a Collins function, but it does probe  $h_{1g}$  which is not transversity however

# Parallels between SIDIS and HQ pair production

LO asymmetries in HQ pair production:

[Boer, Pisano, Mulders, Zhou, 2016]

$$|\langle \cos 2\phi_T \rangle| = \left| \frac{\int d\phi_\perp d\phi_T \cos 2\phi_T d\sigma}{\int d\phi_\perp d\phi_T d\sigma} \right| = \frac{|\mathbf{q}_T|^2 |B_0^U|}{2 A_0^U} = \frac{|\mathbf{q}_T|^2}{2M^2} \frac{|h_1^{\perp g}(x, \mathbf{p}_T^2)|}{f_1^g(x, \mathbf{p}_T^2)} \frac{|\mathcal{B}_0^{eg \rightarrow eQ\bar{Q}}|}{\mathcal{A}_0^{eg \rightarrow eQ\bar{Q}}}$$

$$A_N^{\sin(\phi_S - \phi_T)} = \frac{|\mathbf{q}_T|}{M_p} \frac{A_0^T}{A_0^U} = \frac{|\mathbf{q}_T|}{M_p} \frac{f_{1T}^{\perp g}(x, \mathbf{q}_T^2)}{f_1^g(x, \mathbf{q}_T^2)}$$

$$A_N^{\sin(\phi_S + \phi_T)} = \frac{|\mathbf{q}_T|}{M_p} \frac{{B'_0}^T}{A_0^U} = \frac{2(1-y) \mathcal{B}_{0T}^{\gamma^* g \rightarrow Q\bar{Q}}}{[1 + (1-y)^2] \mathcal{A}_{U+L}^{\gamma^* g \rightarrow Q\bar{Q}} - y^2 \mathcal{A}_L^{\gamma^* g \rightarrow Q\bar{Q}}} \frac{|\mathbf{q}_T|}{M_p} \frac{h_1^g(x, \mathbf{q}_T^2)}{f_1^g(x, \mathbf{q}_T^2)}$$

$$A_N^{\sin(\phi_S - 3\phi_T)} = -\frac{|\mathbf{q}_T|^3}{M_p^3} \frac{B_0^T}{2A_0^U} = -\frac{2(1-y) \mathcal{B}_{0T}^{\gamma^* g \rightarrow Q\bar{Q}}}{[1 + (1-y)^2] \mathcal{A}_{U+L}^{\gamma^* g \rightarrow Q\bar{Q}} - y^2 \mathcal{A}_L^{\gamma^* g \rightarrow Q\bar{Q}}} \frac{|\mathbf{q}_T|^3}{2M_p^3} \frac{h_{1T}^{\perp g}(x, \mathbf{q}_T^2)}{f_1^g(x, \mathbf{q}_T^2)}$$


SIDIS - Fragmentation functions

HQ pairs - calculable amplitudes

# Quarkonium production

$e p \rightarrow e' Q X$  with  $Q$  either a  $J/\psi$  or a  $\Upsilon$  meson

[Godbole, Misra, Mukherjee, Rawoot, 2012/3; Godbole, Kaushik, Misra, Rawoot, 2015; Mukherjee, Rajesh, 2017; Rajesh, Kishore, Mukherjee, 2018]



A  $\cos(2\phi_T)$  asymmetry probes  $h_1^{\perp g}$

$$\langle \cos 2\phi_T \rangle = \frac{(1-y) \mathcal{B}_T^{\gamma^* g \rightarrow Q}}{[1 + (1-y)^2] \mathcal{A}_{U+L}^{\gamma^* g \rightarrow Q} - y^2 \mathcal{A}_L^{\gamma^* g \rightarrow Q}} \times \frac{\mathbf{q}_T^2}{2M_p^2} \frac{h_1^{\perp g}(x, \mathbf{q}_T^2)}{f_1^g(x, \mathbf{q}_T^2)}.$$

In LO NRQCD the prefactor of the asymmetry depends on  $y$ ,  $Q$ ,  $M_Q$  and on two quite uncertain Color Octet (CO) Long Distance Matrix Elements (LDMEs)


One can cancel out the CO LDMEs by considering ratios with spin asymmetries

[Bacchetta, Boer, Pisano, Taels, 2018]

# Quarkonium production in ep

$e p^\uparrow \rightarrow e' Q X$  with  $Q$  either a  $J/\psi$  or a  $\Upsilon$  meson

[Godbole, Misra, Mukherjee, Rawoot, 2012/3; Godbole, Kaushik, Misra, Rawoot, 2015; Mukherjee, Rajesh, 2017; Rajesh, Kishore, Mukherjee, 2018]



Using LO NRQCD the Sivers asymmetry is:

$$A^{\sin(\phi_S - \phi_T)} = \frac{|\mathbf{q}_T|}{M_p} \frac{f_{1T}^{\perp g}(x, \mathbf{q}_T^2)}{f_1^g(x, \mathbf{q}_T^2)}$$

Other asymmetries depend on the quite uncertain CO NRQCD LDMEs, but one can consider ratios of asymmetries to cancel them out

[Bacchetta, Boer, Pisano, Taelis, 2018]

$$\frac{A^{\cos 2\phi_T}}{A^{\sin(\phi_S + \phi_T)}} = \frac{\mathbf{q}_T^2}{M_p^2} \frac{h_1^{\perp g}(x, \mathbf{q}_T^2)}{h_1^g(x, \mathbf{q}_T^2)}$$

$$\frac{A^{\cos 2\phi_T}}{A^{\sin(\phi_S - 3\phi_T)}} = -\frac{1}{2} \frac{h_1^{\perp g}(x, \mathbf{q}_T^2)}{h_{1T}^{\perp g}(x, \mathbf{q}_T^2)}$$

$$\frac{A^{\sin(\phi_S - 3\phi_T)}}{A^{\sin(\phi_S + \phi_T)}} = -\frac{\mathbf{q}_T^2}{2M_p^2} \frac{h_{1T}^{\perp g}(x, \mathbf{q}_T^2)}{h_1^g(x, \mathbf{q}_T^2)}$$

# Process dependence of gluon TMDs


# Gauge links

semi-inclusive DIS

$$e p \rightarrow e' h X$$

$$k \approx xP + k_T$$


$$P^\mu \approx P^+$$



$$\mathcal{U}_C[0, \xi] = \mathcal{P} \exp \left( -ig \int_{C[0, \xi]} ds_\mu A^\mu(s) \right)$$

$$\xi = [0^+, \xi^-, \xi_T]$$

Drell-Yan:



The path  $C$  depends on whether the color interactions are with an incoming or outgoing color charge, yielding different paths for different processes

[Collins & Soper, 1983; Boer & Mulders, 2000]

# Process dependence of gluon TMDs

This has observable effects, as was first noted for quark Sivers asymmetries  
[Brodsky, Hwang & Schmidt, 2002; Collins, 2002; Belitsky, Ji & Yuan, 2003]

$$f_{1T}^{\perp q[\text{SIDIS}]}(x, k_T^2) = -f_{1T}^{\perp q[\text{DY}]}(x, k_T^2) \quad [\text{Collins '02}]$$

A similar sign change relation for gluon Sivers functions holds, but due to the appearance of two gauge links, there are more possibilities

$$\Gamma_g^{\mu\nu[\mathcal{U},\mathcal{U}']}(x, k_T) \equiv \text{F.T.} \langle P | \text{Tr}_c \left[ F^{+\nu}(0) \mathcal{U}_{[0,\xi]} F^{+\mu}(\xi) \mathcal{U}'_{[\xi,0]} \right] | P \rangle$$

For most gluon TMDs there are only 2 link combinations of interest:  $[+,+]$  &  $[+,-]$

$[-,-]$  &  $[-,+]$  are related to them by parity and time reversal

More complicated links arise in processes where TMD factorization is questionable

The gauge link dependence even affects unpolarized gluon TMDs  
[Dominguez, Marquet, Xiao, Yuan, 2011]

# Sign change relation for gluon Sivers TMD


$$e p^\uparrow \rightarrow e' Q \bar{Q} X \quad \gamma^* g \rightarrow Q \bar{Q} \text{ probes } [+,-]$$

$$p^\uparrow p \rightarrow \gamma \gamma X$$

Qiu, Schlegel, Vogelsang, 2011

In the kinematic regime where pair rapidity is central, one effectively selects the subprocess:

$$g g \rightarrow \gamma \gamma \text{ probes } [-,-]$$




$$f_{1T}^\perp g [e p^\uparrow \rightarrow e' Q \bar{Q} X](x, p_T^2) = - f_{1T}^\perp g [p^\uparrow p \rightarrow \gamma \gamma X](x, p_T^2)$$

Boer, Mulders, Pisano, J. Zhou, 2016

# Asymmetries in heavy quark pair production

Sivers asymmetry in D-meson pair production at EIC:



$$f_{1T}^{\perp g} [+ , +]$$

Assumes gluon Sivers TMD that is 10% of the positivity bound

Luminosity of  $10 \text{ fb}^{-1}$

Zheng, Aschenauer, Lee, Xiao, Jin, 2018

$$\langle x_B \rangle = 0.0012$$

The  $[+,+]$  Sivers TMD lacks the  $1/x$  growth of the unpolarized gluon TMD (at least in the perturbative  $k_T$  regime), hence 10% at  $x=0.001$  may be too optimistic

Boer, Echevarria, Mulders, J. Zhou, PRL 2016

Jet pair production more promising, but receives contributions from quark TMDs

# High- $p_T$ hadron pairs

COMPASS measured the gluon Sivers asymmetry in high- $p_T$  hadron pair production in muon-deuteron and muon-proton scattering

The combined result for PGF:

$$A^{Siv} = -0.23 \pm 0.08 \text{ (stat)} \pm 0.05 \text{ (syst) at } \langle x_g \rangle = 0.15$$

$$f_{1T}^{\perp g [+,+]}$$

[C.Adolph et al., PLB 2017]

Imposed requirement:  $p_{1T} > 0.7 \text{ GeV}/c$  and  $p_{2T} > 0.4 \text{ GeV}/c$

The bulk of the data was for  $p_{1T} < 1.7 \text{ GeV}/c$  and  $p_{2T} < 1.1 \text{ GeV}/c$

The gluon contribution has been estimated using MC

There is significant contribution and asymmetry from QCD Compton


EIC projections are promising for gluon Sivers TMDs 5% of the positivity bound

Zheng, Aschenauer, Lee, Xiao, Jin, 2018

# Gluon Sivers effect in double $\gamma$ production

$$f_{1T}^{\perp g} [-,-]$$

$p^\uparrow p \rightarrow \gamma\gamma X$



$\sqrt{s}=500 \text{ GeV}$ ,  $p_T \gamma \geq 1 \text{ GeV}$ , integrated over  $4 < Q^2 < 30 \text{ GeV}^2$ ,  $0 \leq q_T \leq 1 \text{ GeV}$

At photon pair rapidity  $y < 3$  gluon Sivers dominates and  $\max(d\sigma_{TU}/d\sigma_{UU}) \sim 30\text{-}50\%$

Asymmetry may be large but di-photon rate is not at RHIC, so also very challenging

Same may apply to  $J/\psi \gamma$  and  $J/\psi J/\psi$  pair production in  $p^\uparrow p$  collisions

# f and d type gluon Sivers TMD

$$e p^\uparrow \rightarrow e' Q \bar{Q} X$$

$$\gamma^* g \rightarrow Q \bar{Q} \text{ probes } [+,+]$$

$$p^\uparrow p \rightarrow \gamma \text{ jet } X$$

In the kinematic regime where gluons in the polarized proton dominate, one effectively selects the subprocess:

$$g q \rightarrow \gamma q \text{ probes } [+,-]$$



These processes probe 2 distinct, *independent* gluon Sivers functions

Related to the antisymmetric ( $f^{abc}$ ) and symmetric ( $d^{abc}$ ) color structures

Bomhof, Mulders, 2007; Buffing, Mukherjee, Mulders, 2013

Conclusion: gluon Sivers TMD studies using different processes can be related or complementary

# Gluon Sivers effect in $\gamma$ jet production


[Bacchetta, Bomhof, D'Alesio, Mulders, Murgia, 2007]

$$f_{1T}^{\perp g} [+, -]$$

$p^\uparrow p \rightarrow \gamma \text{ jet } X$

maximum contribution from the gluon Sivers function (absolute value)

Prediction for the azimuthal moment at  $\sqrt{s}=200$  GeV and  $p_{T\gamma} \geq 1$  GeV



$$M_N^{\gamma j}(\eta_\gamma, \eta_j, x_\perp) = \frac{\int d\phi_j d\phi_\gamma \frac{2|\mathbf{K}_{\gamma\perp}|}{M} \sin(\delta\phi) \cos(\phi_\gamma) \frac{d\sigma}{d\phi_j d\phi_\gamma}}{\int d\phi_j d\phi_\gamma \frac{d\sigma}{d\phi_j d\phi_\gamma}}$$

# Gluon Sivers effect at small x

Selection of processes that probe the f-type or d-type Sivers gluon TMD:

	$e p^\uparrow \rightarrow e' Q \bar{Q} X$ $e p^\uparrow \rightarrow e' j_1 j_2 X$	$p^\uparrow A \rightarrow h X$ ( $x_F < 0$ )	$p^\uparrow A \rightarrow \gamma^{(*)} \text{jet } X$	$p^\uparrow p \rightarrow \gamma \gamma X$ $p^\uparrow p \rightarrow J/\psi \gamma X$ $p^\uparrow p \rightarrow J/\psi J/\psi X$
$f_{1T}^{\perp g [+,+]}$	✓	✗	✗	✓
$f_{1T}^{\perp g [+,-]}$	✗	✓	✓	✗

The d-type Sivers gluon TMD can also be probed in  $p^\uparrow A \rightarrow h X$  in the backward region

Small x in the polarized proton and large A will enhance gluon-gluon scattering

The *leading twist*  $[+,-]$  correlator becomes in the small-x limit:

$$\Gamma^{[+,-] ij}(x, \mathbf{k}_T) \xrightarrow{x \rightarrow 0} \frac{k_T^i k_T^j}{2\pi L} \Gamma_0^{[\square]}(\mathbf{k}_T) \quad \text{a single Wilson loop matrix element}$$

# d-type gluon Sivers effect

The d-type gluon Sivers function  $f_{1T}^{\perp g [+, -]}$  at small  $x$  is part of:

$$\Gamma_{(d)}^{(T-\text{odd})} \equiv (\Gamma^{[+, -]} - \Gamma^{[-, +]}) \propto \text{F.T.} \langle P, S_T | \text{Tr} \left[ U^{[\square]}(0_T, y_T) - U^{[\square]\dagger}(0_T, y_T) \right] |P, S_T \rangle$$

Boer, Echevarria, Mulders, J. Zhou, PRL 2016

At small  $x$  it can be identified with the *spin-dependent odderon* [J. Zhou, 2013]


It is the only relevant contribution to  $A_N$  in backward ( $x_F < 0$ ) charged hadron production in  $p^\uparrow A$

$A_N$  is not a TMD factorizing process, but at small  $x$  one can apply a hybrid factorization (at least at one-loop order)

[Chirilli, Xiao, Yuan, 2012]

As the odderon is C-odd, for gg-dominated scattering one should select final states that are not C-even, hence charged hadron production (as opposed to jets or  $\pi^0$ )


$p^\uparrow p \rightarrow h^\pm X$  at  $x_F < 0$



BRAHMS, 2008  $\sqrt{s} = 62.4$  GeV  
low  $p_T$ , up to roughly 1.2 GeV  
where gg channel dominates

PHENIX, 2017  
 $\sqrt{s} = 200$  GeV  
 $p_T$  between 1.25 and 7 GeV

Interesting process to study e.g. at NICA



# Probes of unpolarized gluon TMDs

## WW vs DP

For most processes of interest there are 2 relevant unpolarized gluon distributions

Dominguez, Marquet, Xiao, Yuan, 2011

$$xG^{(1)}(x, k_\perp) = 2 \int \frac{d\xi^- d\xi_\perp}{(2\pi)^3 P^+} e^{ixP^+ \xi^- - ik_\perp \cdot \xi_\perp} \langle P | \text{Tr} [F^{+i}(\xi^-, \xi_\perp) \mathcal{U}^{[+]^\dagger} F^{+i}(0) \mathcal{U}^{[+]}] | P \rangle \quad [+, +]$$

$$xG^{(2)}(x, k_\perp) = 2 \int \frac{d\xi^- d\xi_\perp}{(2\pi)^3 P^+} e^{ixP^+ \xi^- - ik_\perp \cdot \xi_\perp} \langle P | \text{Tr} [F^{+i}(\xi^-, \xi_\perp) \mathcal{U}^{[-]^\dagger} F^{+i}(0) \mathcal{U}^{[+]}] | P \rangle \quad [+, -]$$

For unpolarized gluons  $[+, +] = [-, -]$  and  $[+, -] = [-, +]$

At small  $x$  the two correspond to the Weizsäcker-Williams (WW) and dipole (DP) distributions, which are generally different in magnitude and width:

$$xG^{(1)}(x, k_\perp) = -\frac{2}{\alpha_S} \int \frac{d^2 v}{(2\pi)^2} \frac{d^2 v'}{(2\pi)^2} e^{-ik_\perp \cdot (v-v')} \langle \text{Tr} [\partial_i U(v)] U^\dagger(v') [\partial_i U(v')] U^\dagger(v) \rangle_{x_g} \quad \text{WW}$$


$$xG^{(2)}(x, q_\perp) = \frac{q_\perp^2 N_c}{2\pi^2 \alpha_s} S_\perp \int \frac{d^2 r_\perp}{(2\pi)^2} e^{-iq_\perp \cdot r_\perp} \frac{1}{N_c} \langle \text{Tr} U(0) U^\dagger(r_\perp) \rangle_{x_g} \quad \text{DP}$$

Different processes probe one or the other or a mixture, so this can be tested

# Processes for unpolarized gluon TMDs

Selection of processes that probe the WW or DP unpolarized gluon TMD:

	$pA \rightarrow \gamma \text{jet } X$	$e p \rightarrow e' Q \bar{Q} X$ $e p \rightarrow e' j_1 j_2 X$	$pp \rightarrow \eta_{c,b} X$ $pp \rightarrow H X$	$pp \rightarrow J/\psi \gamma X$ $pp \rightarrow \Upsilon \gamma X$
$f_1^g [+,+]$ (WW)	×	✓	✓	✓
$f_1^g [+,-]$ (DP)	✓	✗	✗	✗



$pp \rightarrow Q \gamma X$

a good process at LHC  
to extract

$f_1^g [+,+]$

[Den Dunnen, Lansberg, Pisano, Schlegel, 2014]

# Processes for unpolarized gluon TMDs

Heavy quarks are generally promising to exploit, but not in all processes:

$pp \rightarrow J/\psi X$  or  $\Upsilon X$  Color Singlet (CS) vector quarkonium production from 2 gluons is forbidden by Landau-Yang theorem, while Color Octet (CO) production involves a more complicated link structure

For C-even (pseudo-)scalar quarkonium production  $gg \rightarrow CS$  is leading contribution

$pp \rightarrow \eta_c X$  or  $pp \rightarrow \chi_c X$  could be studied at NICA

In LO NRQCD the differential cross sections in pp and pA are:

$$\frac{d\sigma(\eta_Q)}{dy d^2\mathbf{q}_T} = \frac{2}{9} \frac{\pi^3 \alpha_s^2}{M^3 s} \langle 0 | \mathcal{O}_1^{\eta_Q}(^1S_0) | 0 \rangle \mathcal{C} [f_1^g f_1^g] [1 - R(\mathbf{q}_T^2)]$$

$$\frac{d\sigma(\chi_{Q0})}{dy d^2\mathbf{q}_T} = \frac{8}{3} \frac{\pi^3 \alpha_s^2}{M^5 s} \langle 0 | \mathcal{O}_1^{\chi_{Q0}}(^3P_0) | 0 \rangle \mathcal{C} [f_1^g f_1^g] [1 + R(\mathbf{q}_T^2)]$$

$$\frac{d\sigma(\chi_{Q2})}{dy d^2\mathbf{q}_T} = \frac{32}{9} \frac{\pi^3 \alpha_s^2}{M^5 s} \langle 0 | \mathcal{O}_1^{\chi_{Q2}}(^3P_2) | 0 \rangle \mathcal{C} [f_1^g f_1^g]$$

[Boer, Pisano, 2012]

$R(\mathbf{q}_T^2)$  is the contribution from the linearly polarized gluon TMD  $h_{I^\perp g}$

# Probes of linearly polarized gluons TMDs

# Probes of linear gluon polarization

For linearly polarized gluons  $[+,+] = [-,-]$  and  $[+,-] = [-,+]$

Processes that probe the linearly polarized gluon TMD:

	$pp \rightarrow \gamma\gamma X$	$pA \rightarrow \gamma^* \text{jet } X$	$ep \rightarrow e' Q\bar{Q} X$ $ep \rightarrow e' j_1 j_2 X$	$pp \rightarrow \eta_{c,b} X$ $pp \rightarrow H X$	$pp \rightarrow J/\psi \gamma X$ $pp \rightarrow \Upsilon \gamma X$
$h_1^{\perp g [+,+]} (\text{WW})$	✓	✗	✓	✓	✓
$h_1^{\perp g [+,-]} (\text{DP})$	✗	✓	✗	✗	✗

1% level at RHIC

Qiu, Schlegel, Vogelsang, 2011

5% level at RHIC

Boer, Mulders, J. Zhou, Y. Zhou, 2017

10% level at EIC

Boer, Brodsky, Pisano, Mulders, 2011;  
Dumitru, Lappi, Skokov, 2015;  
Boer, Pisano, Mulders, J. Zhou, 2016;  
Efremov, Ivanov, Teryaev, 2018

10% level for  $\eta_Q$  and

1% level for Higgs at LHC

Boer & den Dunnen, 2014;  
Echevarria, Kasemets,  
Mulders, Pisano, 2015

Higgs and  $0^{\pm\pm}$  quarkonium production uses the angular *independent*  $p_T$  distribution

All other suggestions use angular modulations


$pp \rightarrow Q\bar{Q} X$  [Akçakaya, Schäfer, Zhou, 2013; Pisano, Boer, Brodsky, Buffing, Mulders, 2013]

TMD factorization is a concern here [Catani, Grazzini, Torre, 2015]

# Open heavy quark electro-production

Unpolarized open heavy quark production at EIC allows to probe  $h_1^{\perp g}(x, p_T^2)$

$$ep \rightarrow e' Q \bar{Q} X$$



no convolution!

[Boer, Brodsky, Mulders & Pisano, 2010]



The individual transverse momenta have to be large but their sum has to be small

Linearly polarized gluons lead to  $\cos 2\phi_T$  or  $\cos 2(\phi_T - \phi_\perp)$  angular modulation

$h_{1\perp g}$  appears by itself, so its sign can be determined and the effects could be significant, especially towards smaller  $x$  as it follows the fast growth of  $f_{1g}$

# Asymmetries in heavy quark pair production

$h_1^{\perp g}$  expected to keep up with growth of the unpolarized gluons TMD as  $x \rightarrow 0$



small  $x$   
**MV model**

$|K_\perp| = 10 \text{ GeV}$   
 $z = 0.5$   
 $y = 0.3$


Sizable asymmetries at EIC  
[Boer, Pisano, Mulders, Zhou, 2016]

# Dijet production at EIC

$h_{1\perp g}$  ( $WW$ ) is also accessible in dijet production at EIC


[Metz, Zhou 2011; Pisano, Boer, Brodsky, Buffing, Mulders, 2013; Boer, Pisano, Mulders, Zhou, 2016]

Linear gluon polarization shows itself through a  $\cos 2\phi$  distribution (“ $v_2$ ”)



Large effects are found

Dumitru, Lappi, Skokov, 2015




Jets are reasonable proxies for outgoing quarks concerning the  $q_T$  distribution

Dumitru, Skokov, Ullrich, 2018

# Quarkonium production in pp

$\text{pp} \rightarrow \eta_c X$  or  $\text{pp} \rightarrow \chi_c X$  allow to probe the linearly polarized gluon TMD


The relative contribution to the diff. cross section w.r.t. the unpolarized gluons:




[Boer, Den Dunnen, 2014]

# Bottomonium production in pp

The range of predictions for C-even bottomonium production:



Boer & den Dunnen, 2014



Echevarria, Kasemets, Mulders, Pisano, 2015

Very large theoretical uncertainties (from the nonperturbative part of the TMDs), even more for charmonium production, but contribution of 20% or more is expected

# $\gamma^*$ -jet production

$h_{1\perp g}$  is power suppressed in  $p\bar{p} \rightarrow \gamma$  jet  $X$

[Boer, Mulders, Pisano, 2008]

It is not power suppressed in  $p\bar{p} \rightarrow \gamma^* \text{jet } X$  if  $Q^2 \sim P_{\perp,\text{jet}}^2$

[Boer, Mulders, Zhou & Zhou, 2017]

Consider  $Q^2 \sim P_{\perp,\text{jet}}^2$  also to avoid a three-scale problem

	$p\bar{p} \rightarrow \gamma\gamma X$	$pA \rightarrow \gamma^* \text{jet } X$	$e p \rightarrow e' Q \bar{Q} X$ $e p \rightarrow e' j_1 j_2 X$	$p\bar{p} \rightarrow \eta_{c,b} X$ $p\bar{p} \rightarrow H X$	$p\bar{p} \rightarrow J/\psi \gamma X$ $p\bar{p} \rightarrow \Upsilon \gamma X$
$h_1^{\perp g [+,+]} (\text{WW})$	✓	✗	✓	✓	✓
$h_1^{\perp g [+,-]} (\text{DP})$	✗	✓	✗	✗	✗


$p\bar{p} \rightarrow \gamma^* \text{jet } X$  offers a unique opportunity to study the DP linear gluon polarization

At high gluon density (large  $A$  and/or small  $x$ ) the DP linear gluon polarization is expected to become maximal, as was first shown in the MV model for the CGC

$$x h_{1,DP}^{\perp g}(x, k_\perp) = 2x f_{1,DP}^g(x, k_\perp)$$

[Metz & Jian Zhou, 2011; Boer, Cotogno, van Daal, Mulders, Signori & Ya-Jin Zhou, 2016]

# Sudakov suppression of linear gluon polarization





Despite the DP linear gluon polarization becoming maximal at small  $x$ , there is amplitude and Sudakov suppression of the  $\cos(2\varphi)$  asymmetry in  $pA \rightarrow \gamma^* \text{ jet } X$ :

~5% asymmetry at RHIC

[Boer, Mulders, Jian Zhou & Ya-Jin Zhou, 2017]

It becomes effectively power suppressed as  $Q \sim P_\perp$  increases from 6 to 90 GeV

# Linear gluon polarization in di- $J/\Psi$ production



At LHC  $h_{I^\perp g}$  can be probed in  $p p \rightarrow J/\psi J/\psi X$

Estimated to lead to 1-5% level azimuthal modulations

[Scarpa, Boer, Echevarria, Lansberg, Pisano, Schlegel, 2019]

# Conclusions

# Conclusions

- Gluon TMDs measurements generally require higher energy collisions, less inclusive observables (particle pair correlations) and several processes (process dependence)
- Even the unpolarized gluon TMDs have not been extracted yet, but there are plenty of future opportunities
- For the linearly polarized gluon TMDs small  $x$  will be beneficial, for the f-type gluon Sivers TMD small  $x$  is not favorable
- The f-type  $[+,+]$  gluon Sivers TMD enters di-hadron production in SIDIS and satisfies the sign-change relation
- The d-type  $[+,-]$  gluon Sivers TMD at small  $x$  corresponds to the spin-dependent odderon, a C-odd Wilson loop matrix element that fully determines  $A_N$  at negative  $x_F$

# Main opportunities

$f_1^g [+,+]$	$pp \rightarrow \gamma J/\psi X$ $pp \rightarrow \gamma \Upsilon X$	LHC LHC
$f_1^g [+,-]$	$pp \rightarrow \gamma \text{jet} X$	LHC & RHIC
$h_1^{\perp g} [+,+]$	$e p \rightarrow e' Q \bar{Q} X$ $e p \rightarrow e' \text{jet jet} X$ $pp \rightarrow \eta_{c,b} X$ $pp \rightarrow H X$	EIC EIC LHC & NICA LHC
$h_1^{\perp g} [+,-]$	$pp \rightarrow \gamma^* \text{jet} X$	LHC & RHIC
$f_{1T}^{\perp g} [+,+]$	$e p^\uparrow \rightarrow e' Q \bar{Q} X$ $e p^\uparrow \rightarrow e' \text{jet jet} X$	EIC EIC
$f_{1T}^{\perp g} [-,-]$	$p^\uparrow p \rightarrow \gamma \gamma X$	RHIC
$f_{1T}^{\perp g} [+,-]$	$p^\uparrow A \rightarrow \gamma^{(*)} \text{jet} X$ $p^\uparrow A \rightarrow h X (x_F < 0)$	RHIC RHIC & NICA