



Contribution ID: 238

Type: not specified

## Confining vs. conformal scenario for $SU(2)$ with 2 adjoint fermions. Mesonic spectrum.

*Tuesday, 15 June 2010 09:10 (20 minutes)*

The Minimal Walking Technicolor model, based on the  $SU(2)$  gauge group with two Dirac adjoint fermions, is expected to lie close to the lower boundary of the conformal window. As such, it is believed to possess a dynamics different enough from QCD to be a viable candidate for a Technicolor theory. We present a detailed analysis of the mesonic spectrum of this model in which we aim to reach the chiral limit while controlling the systematic errors. Comparing our data to the expected behaviors for a QCD-like theory and for an IR conformal theory, we show that the latter scenario is favored.

**Please, insert your presentation type (talk, poster)**

talk

**Primary author:** PICA, Claudio (CP3-Origins, SDU)

**Co-authors:** Dr PATELLA, Agostino (Swansea University); Dr RAGO, Antonio (Wuppertal University); Dr LUCINI, Biagio (Swansea University); Dr DEL DEBBIO, Luigi (University of Edinburgh)

**Presenter:** PICA, Claudio (CP3-Origins, SDU)

**Session Classification:** Parallel 13: Applications beyond QCD

**Track Classification:** Applications beyond QCD