



Contribution ID: 147

Type: **not specified**

## Glueballs in the radiative J/psi decays

*Tuesday, 15 June 2010 08:30 (20 minutes)*

In the quenched approximation, glueball-vector-current-Jpsi three point functions are calculated on an anisotropic lattice with very high statistics. Using the relevant form factors derived from these three point functions, we also give a prediction of the radiative decay widths of J/psi to scalar and pseudoscalar glueballs.

### Please, insert your presentation type (talk, poster)

talk

**Primary author:** CHEN, Ying (Institute of High Energy Physics, Chinese Academy of Sciences, China)

**Co-authors:** LIU, Chuan (School of Physics, Peking University, China); ZHANG, Jian-Bo (Department of Physics, Zhejiang University, China); MA, Jian-Ping (Institute of Theoretical Physics, Chinese Academy of Sciences, China); GUI, Long-Cheng (Institute of High Energy Physics, Chinese Academy of Sciences, China); LIU, Yu-Bin (School of Physics, Nankai University, China); ZHANG, Yuan-Jiang (Institute of High Energy Physics, Chinese Academy of Sciences, China)

**Presenter:** CHEN, Ying (Institute of High Energy Physics, Chinese Academy of Sciences, China)

**Session Classification:** Parallel 21: Hadron spectroscopy

**Track Classification:** Hadron spectroscopy