



Contribution ID: 148

Type: not specified

Lattice QCD study of baryon-baryon interactions in the $(S,I)=(-2,0)$ system using the coupled-channel formalism

Friday, 18 June 2010 15:30 (20 minutes)

We investigate baryon-baryon interactions with strangeness $S=-2$ and isospin $I=0$ system from Lattice QCD. The study of $S=-2$ system opens a gate of multi-strangeness hadronic world and provides the unified understanding of YN and YY interactions.

A satisfactory description of YN and YY interaction is not yet obtained with use of phenomenological meson exchange model due to the lack of the direct measurement of hyperon-hyperon scattering to determine many free parameters.

It is important to understand these interactions directly from QCD.

In order to solve this system, we prepare three types of baryon-baryon operators (Λ - Λ , N - Ξ and Σ - Σ) and construct three operators diagonalizing the 3×3 correlation matrix.

Combining of these sink operators with the diagonalized source operators, we obtain nine effective Bethe-Salpeter wave functions.

The 3×3 potential matrix is calculated by solving the coupled-channel Schroedinger equation.

The flavor $SU(3)$ breaking effects of the potential matrix are also discussed by comparing with the results of $SU(3)$ limit in the same calculation.

Our numerical results are obtained from 2+1 flavor QCD gauge configurations with $m_{\pi} = 870$ MeV and $m_{\pi}/m_K = 0.95$, provided by the CP-PACS/JLQCD Collaborations.

Please, insert your presentation type (talk, poster)

talk

Primary author: SASAKI, Kenji (University of Tsukuba)

Presenter: SASAKI, Kenji (University of Tsukuba)

Session Classification: Parallel 49: Hadronic structure and interactions

Track Classification: Hadronic structure and interactions